

Spivak Notes and Solutions

Max Chien

February 2024

Contents

1	Euclidean Space	3
1.1	Vector Properties of Euclidean Space	3
1.2	Topology of Euclidean Space	5
1.3	Functions and Continuity	8
2	Differentiation	12
2.1	Basic Definitions	12
2.2	Basic Theorems	13
2.3	Partial Derivatives	19
2.4	Derivatives	20
2.5	Inverse Functions	24
2.6	Implicit Functions	28
3	Integration	31
3.1	Basic Definitions	31
3.2	Measure Zero and Content Zero	34
3.3	Integrable Functions	36
3.4	Fubini's Theorem	41
3.5	Partitions of Unity	45
3.6	Change of Variables	53
4	Integration on Chains	60
4.1	Algebraic Preliminaries	60
A	Solutions to Selected Exercises	72
A.1	Chapter 1 Exercises	72
A.2	Chapter 2 Exercises	87
A.3	Chapter 3 Exercises	130
	Definitions	169

Introduction

This document contains notes taken as personal self-study in Summer 2024 of the book *Calculus on Manifolds*, by Michael Spivak. The notes closely follow the structure of Spivak's text. The appendix contains solutions for selected exercises out of the book.

Chapter 1

Euclidean Space

1.1 Vector Properties of Euclidean Space

In this course, we study functions over Euclidean space. We will assume knowledge of most of the basic properties of the real numbers, and will only briefly introduce the basic properties of Euclidean space.

Definition 1.1

Euclidean n -space, denoted \mathbb{R}^n , is the set of n -tuples

$$(x_1, x_2, \dots, x_n)$$

such that $x_i \in \mathbb{R}$ for each i .

Euclidean space is intended to align with the “standard” notions of space. That is, \mathbb{R}^1 is often referred to as the line, \mathbb{R}^2 as the plane, and \mathbb{R}^3 as space. Moreover, from linear algebra we can see that \mathbb{R}^n can be considered as an n -dimensional vector space over \mathbb{R} , with addition and scalar multiplication defined coordinate-wise, so elements of \mathbb{R}^n will alternately be called points or vectors. In fact, it is the canonical representative of n dimensional vector spaces over \mathbb{R} , further justifying its study. We denote by 0 or $\mathbf{0}$ the vector $(0, 0, \dots, 0)$.

Moreover, \mathbb{R}^n is an example of a *normed* vector space. Specifically, we have

Definition 1.2

Given a vector $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, define the **norm** of x , denoted $|x|$, by

$$|x| := \sqrt{x_1^2 + \dots + x_n^2}$$

Note that for $n = 1$, the norm aligns with the standard absolute value of real numbers. Briefly, we can verify that the norm as defined here indeed satisfies the definition of a norm on a vector space:

Proposition 1.1

Let $x, y \in \mathbb{R}^n$, and $a \in \mathbb{R}$ be arbitrary. Then we have:

- $|x| \geq 0$, with $|x| = 0$ if and only if $x = 0$.
- $|\sum_{i=1}^n x_i y_i| \leq |x||y|$, with equality if and only if x, y are linearly dependent.
- $|x + y| \leq |x| + |y|$.
- $|ax| = |a||x|$

Beyond being a normed vector space, Euclidean space is also an inner product space. We can define the inner product as follows:

Definition 1.3

Given two vectors $x, y \in \mathbb{R}^n$, define the **inner product** of x and y , denoted $\langle x, y \rangle$, as

$$\langle x, y \rangle := \sum_{i=1}^n x_i y_i$$

Similarly, we can verify that this inner product satisfies the definitions of an inner product:

Proposition 1.2

Let $x, x_1, x_2, y, y_1, y_2 \in \mathbb{R}^n$ and $a \in \mathbb{R}$ be arbitrary. Then we have:

- $\langle x, y \rangle = \langle y, x \rangle$ (Symmetric)
- $a \langle x, y \rangle = \langle ax, y \rangle = \langle x, ay \rangle$ (Bilinear)
 $\langle x_1 + x_2, y \rangle = \langle x_1, y \rangle + \langle x_2, y \rangle$
 $\langle x, y_1 + y_2 \rangle = \langle x, y_1 \rangle + \langle x, y_2 \rangle$
- $\langle x, x \rangle \geq 0$, with $\langle x, x \rangle = 0$ if and only if $x = 0$. (Positive definite)

Moreover, given our definitions of the norm and inner product, we can also identify further properties:

Proposition 1.3

Let $x, y \in \mathbb{R}^n$ be arbitrary. Then we have:

- $\langle x, y \rangle \leq |x||y|$ (Cauchy-Schwarz Inequality)
- $|x| = \sqrt{\langle x, x \rangle}$
- $\langle x, y \rangle = \frac{|x+y|^2 - |x-y|^2}{4}$ (Polarization Identity)

Definition 1.4

The **standard basis** of \mathbb{R}^n is given by $\{e_1, \dots, e_n\}$, where $(e_i)_j = \delta_{ij}$, so that e_i has a 1 in the i th coordinate and 0 everywhere else.

Definition 1.5

Let $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a linear transformation. Then denote by $[T]$ the $n \times m$ matrix such that $T(x) = [T]x$ for each $x \in \mathbb{R}^m$. In particular, the i th column of $[T]$ is given by $T(e_i)$.

If $x = (x_1, \dots, x_m) \in \mathbb{R}^m$ and $y = (y_1, \dots, y_n) \in \mathbb{R}^n$, then let us adopt the convention that (x, y) is the concatenation $(x_1, \dots, x_m, y_1, \dots, y_n) \in \mathbb{R}^{m+n}$.

1.2 Topology of Euclidean Space

In many results in single variable analysis, we make use of open and closed intervals, denoted $[a, b]$ and (a, b) . The analog of these intervals in \mathbb{R}^n is the notion of a *rectangle* or *k-cell*.

Definition 1.6

Let $A \subseteq \mathbb{R}^m$, $B \subseteq \mathbb{R}^n$. Then define the **Cartesian product** of A and B as $A \times B = \{(a, b) \in \mathbb{R}^{m+n} \mid a \in A, b \in B\}$. Since this operation is associative, denote by $A_1 \times A_2 \times \dots \times A_i$ the product of any number of sets.

Definition 1.7

A **closed rectangle**, closed box, or **closed k-cell** is a set of the form $[a_1, b_1] \times \dots \times [a_n, b_n] \subseteq \mathbb{R}^k$. An **open rectangle**, open box, or **open k-cell** is a set of the form $(a_1, b_1) \times \dots \times (a_n, b_n) \subseteq \mathbb{R}^k$.

Then similarly to how we use open intervals to define a topology on \mathbb{R} , we can use open boxes to define a topology on \mathbb{R}^n :

Definition 1.8

A set $U \subseteq \mathbb{R}^n$ is **open** if, for every point $x \in U$, there is some open box $B(x) \subseteq U$ such that $x \in B(x)$. A set $C \subseteq \mathbb{R}^n$ is **closed** if $\mathbb{R}^n \setminus C$ is open.

Remark

Note that because every open box has an open ball inside, and because every open ball has an open box inside, the topology defined by open boxes on \mathbb{R}^n is the same topology defined by open balls on \mathbb{R}^n . Thus, for $x \in \mathbb{R}^n$ and $r \in \mathbb{R}$, denote by $B_r(x)$ the **open n -ball** with center x and radius r . That is, $B_r(x) := \{y \in \mathbb{R}^n : |x - y| < r\}$. When the dimension is ambiguous, denote this ${}_n B_r(x)$. Then we can alternately use open balls and open boxes as the definition of an open set, depending on which is more convenient.

Definition 1.9

If $A \subseteq \mathbb{R}^n$, then the **interior** of A is the set of points contained in an open rectangle entirely in A .

$$\text{int } A := \{x \in \mathbb{R}^n : \text{there exists an open rectangle } B \text{ s.t. } x \in B \subseteq A\}$$

Define the **exterior** of A to be the set of points contained in an open rectangle entirely in $\mathbb{R}^n \setminus A$.

$$\text{ext } A := \{x \in \mathbb{R}^n : \text{there exists an open rectangle } B \text{ s.t. } x \in B \subseteq \mathbb{R}^n \setminus A\}$$

Define the **boundary** of A to be the set of points where all open rectangles contain points of both A and $\mathbb{R}^n \setminus A$.

$$\partial A := \{x \in \mathbb{R}^n : \forall \text{ open rectangles } B, x \in B \implies B \cap A \neq \emptyset, B \cap \mathbb{R}^n \setminus A \neq \emptyset\}$$

Proposition 1.4

Every set of finitely many points in \mathbb{R}^n is closed.

Proof. Let $C \subseteq \mathbb{R}^n$ be a finite set. Let $x \in \mathbb{R}^n \setminus C$ be arbitrary. Then for each point $y \in C$, $x \neq y$, so $d(x, y) > 0$. Then since there are only finitely many points in C , the quantity $d' = \min\{d(x, y) | y \in C\}$ is defined and greater than 0. So we can define an open ball with radius $d'/2$, which does not contain any points in C . Thus we have an open ball containing x that is a subset of $\mathbb{R}^n \setminus C$. So $\mathbb{R}^n \setminus C$ is open and thus C is closed. \square

Definition 1.10

An **open cover** of a set A is a collection \mathcal{O} of open sets such that for any $x \in A$, $x \in U$ for some $U \in \mathcal{O}$. A **subcover** of \mathcal{O} is a subcollection of \mathcal{O} which is also a cover for A .

Definition 1.11

A set K is **compact** if for any open cover \mathcal{O} of K , there exists a finite subcover \mathcal{U} of \mathcal{O} .

In particular, we can derive certain theorems to identify compact sets.

Theorem 1.5: Heine-Borel Theorem

The closed interval $[a, b]$ is compact.

Proof. Let \mathcal{U} be some open cover of $[a, b]$. Then consider the set

$$A = \{x \in [a, b] : [a, x] \text{ is covered by a finite number of sets in } \mathcal{U}\}$$

The goal is to prove that $b \in A$. First, consider $\alpha = \sup A$ (since this set is bounded above and nonempty). We have $\alpha \leq b$, so $\alpha \in [a, b]$ and thus $\alpha \in U_1$ for some $U_1 \in \mathcal{U}$. Since U_1 is open and α is the supremum of A , there is some $a \leq x < \alpha$ with $x \in U_1$. Then we have $x \in A$, so some finite number of open sets in \mathcal{U} cover $[a, x]$, and U_1 covers $[x, \alpha]$, so a finite number of sets cover $[a, \alpha]$ and thus $\alpha \in A$.

Now suppose $\alpha < b$. Then there is some $y \in U_1$ such that $\alpha < y < b$. But if $[a, \alpha]$ is covered by a finite number of open sets, then so is $[a, y]$, so $y \in A$, contradicting $\alpha = \sup A$. So we must have $\alpha = b$, completing the proof. \square

Note that if $B \subseteq \mathbb{R}^m$ is compact and $x \in \mathbb{R}^n$, then the set $\{x\} \times B$ is clearly compact. Moreover, given any cover of $\{x\} \times B$, the finite subcovers have a “minimum width”:

Theorem 1.6

If $B \subseteq \mathbb{R}^m$ is compact and $x \in \mathbb{R}^n$, then given any open cover \mathcal{U} of $\{x\} \times B$, there is some open set $U \subseteq \mathbb{R}^n$ such that $U \times B$ is covered by a finite number of sets in \mathcal{U} .

Proof. Take some finite subcover \mathcal{U}' of \mathcal{U} . Then we just need to find a set U such that $U \times B$ is covered by \mathcal{U}' .

For each $y \in B$, (x, y) is in some open set $O \in \mathcal{U}'$, so there is an open box $U_x \times V_y$ such that $(x, y) \in U_x \times V_y \subseteq O$. Then consider the collection $(V_y)_{y \in B}$. This set covers B , which is compact, so we can pick a finite number V_1, \dots, V_k . Let $U = \bigcap U_i$. Then for any $(x_1, y_1) \in U \times B$, we have $y_1 \in V_i$ for some $1 \leq i \leq k$, and $x_1 \in U_i$, so $x_1 \in U_i \times V_i \subseteq O'$ for some $O' \in \mathcal{U}'$. Thus \mathcal{U}' covers $U \times B$. \square

Corollary

If $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$ are compact, then $A \times B \subseteq \mathbb{R}^{n+m}$ is compact.

Proof. Let \mathcal{O} be some open cover of $A \times B$. Then for each $x \in A$, \mathcal{O} covers $\{x\} \times B$, so there is some U_x such that a finite subcover O_{1x}, \dots, O_{kx} covers $U_x \times B$ and $x \in U_x$. Then the collection $(U_x)_{x \in A}$ covers A , so there is a finite subcover U_{x_1}, \dots, U_{x_j} that covers A . Then the sets $O_{1x_1}, \dots, O_{kx_1}, \dots, O_{1x_j}, \dots, O_{k'x_j}$ form a finite subcover of \mathcal{O} that covers $A \times B$. So $A \times B$ is compact. \square

Corollary

A product $A_1 \times \dots \times A_k$ is compact if each A_i is. A closed rectangle is compact.

Proof. Induct on k using the previous corollary. \square

This gives an important result which allows us to work with compactness much more easily in \mathbb{R}^n (though it is not necessarily true for other topological vector spaces).

Theorem 1.7

A set $K \subseteq \mathbb{R}^n$ is compact if and only if it is closed and bounded.

Proof. (\implies) Suppose $K \subseteq \mathbb{R}^n$ is compact. The collection of open rectangles $(i-1, i+1) \times (j-1, j+1) \dots \times (k-1, k+1)$ for $i, j, \dots, k \in \mathbb{Z}$ covers \mathbb{R}^n , so it covers K . Then a finite number of these boxes covers K , so it is bounded.

(\impliedby) Suppose $K \subseteq \mathbb{R}^n$ is closed and bounded. Then there exists a closed rectangle B with $K \subseteq B$. From the previous corollary, we know that B is compact. Then take some cover of K , $\mathcal{O} = \{O_1, \dots\}$. Now let \mathcal{U} consist of all the sets in \mathcal{O} , as well as the set $\mathbb{R}^n \setminus K$ (which is open since K is closed). \mathcal{U} covers \mathbb{R}^n , so it covers B . Then we can take a finite subcollection \mathcal{U}' of \mathcal{U} . Then \mathcal{U}' covers B as well as K , and in order to create a subcollection of \mathcal{O} , we simply remove $\mathbb{R}^n \setminus K$ if it is in \mathcal{U}' to get \mathcal{O}' . So K is compact. \square

1.3 Functions and Continuity

In this section, we study **vector valued functions**, which are functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, or more generally, $f : A \rightarrow B$ for some $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$. We briefly list a few definitions related to these functions that should be familiar to the reader.

Definition 1.12

If $f : A \rightarrow B$, then the **image** of $C \subseteq A$ is $f(C) = \{f(x) : x \in C\}$. The **preimage** of $D \subseteq B$ is $f^{-1}(D) = \{y \in A : f(y) \in D\}$.

Definition 1.13

If $f : A \rightarrow \mathbb{R}^m$ and $g : B \rightarrow \mathbb{R}^n$ with $B \subseteq \mathbb{R}^m$, then the **composition** is defined as $(g \circ f)(x) = g(f(x))$, with domain $A \cap f^{-1}(B)$.

Definition 1.14

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is one-to-one, then the **inverse** of f is the function $f^{-1} : f(\mathbb{R}^n) \rightarrow \mathbb{R}^n$ which takes $x \in f(\mathbb{R}^n)$ to the unique $y \in \mathbb{R}^n$ such that $f(y) = x$.

In addition to studying a vector valued function f , we can also study the *component functions* which encode its behavior on each axis individually.

Definition 1.15

If $f : A \rightarrow \mathbb{R}^m$, then f defines m **component functions** f^1, f^2, \dots, f^m such that $f(x) = (f^1(x), \dots, f^m(x))$. Similarly, for any functions $g_1, \dots, g_m : A \rightarrow \mathbb{R}$, we denote by (g_1, \dots, g_m) the function $f : A \rightarrow \mathbb{R}^m$ which satisfies $f(x) = (g_1(x), \dots, g_m(x))$.

Note that the above definition implies that we can write $f = (f^1, \dots, f^m)$.

Definition 1.16

Let $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the identity function. Then $\pi = (\pi^1, \dots, \pi^n)$. Then π^i is called the i -th **projection function**, such that $\pi^i(x)$ gives the i th coordinate of x .

With the above out of the way, we now turn our attention to limits of functions, which will prove important as we continue our study of multivariate calculus.

Definition 1.17

We write $\lim_{x \rightarrow a} f(x) = b$ (the **functional limit**) if, for any $\varepsilon > 0$ there exists $\delta = \delta(\varepsilon) > 0$ such that whenever $0 < |x - a| < \delta$, we have $|f(x) - b| < \varepsilon$.

Just as the above definition is reproduced from single-variable analysis (with the exception of generalizing the notion of distance in \mathbb{R}^n), we have an analogous definition of continuity:

Definition 1.18

A function $f : A \rightarrow \mathbb{R}^m$ is **continuous** at a point $a \in A$ if $\lim_{x \rightarrow a} f(x) = f(a)$. If f is continuous at each $a \in A$, we simply say that f is continuous.

Alternatively, we can utilize the topological nature of \mathbb{R}^n , which we discussed in the last section, to characterize continuity using the topological definition instead.

Proposition 1.8

A function $f : A \rightarrow \mathbb{R}^m$ for $A \subseteq \mathbb{R}^n$ is continuous if and only if for every open set $U \subseteq \mathbb{R}^m$, there is an open set $V \subseteq \mathbb{R}^n$ such that $f^{-1}(U) = V \cap A$.

Proof. (\implies) Suppose f is continuous. Then let $U \subseteq \mathbb{R}^m$. For each point $x \in f^{-1}(U)$, $f(x) \in U$ which is open. Thus, there is an open ball $B_{\varepsilon_x}(f(x)) \subseteq U$, there is an open ball $B_{\varepsilon_x}(f(x)) \subseteq U$, and a corresponding open ball $B_{\delta_x}(x) \subseteq f^{-1}(B_{\varepsilon_x}(f(x)))$. Then the set $V = \bigcup_{x \in f^{-1}(U)} B_{\delta_x}(x)$ is an open set.

Moreover, by construction, for any point $y \in V \cap A$, $y \in B_{\delta_x}(x)$ for some x , implying that $f(y) \in B_{\varepsilon_x}(f(x)) \subseteq U$ (which is defined since $y \in A$). So $V \cap A \subseteq f^{-1}(U)$. For any point $x \in f^{-1}(U)$, $x \in B_{\delta_x}(x)$, so $x \in V$. Moreover, any point in $f^{-1}(U)$ is in the domain of f ,

so $x \in V \cap A$, and thus $f^{-1}(U) = V \cap A$.

(\Leftarrow) Suppose every open set $U \subseteq \mathbb{R}^m$ has an associated open set $V \subseteq \mathbb{R}^n$ such that $f^{-1}(U) = V \cap A$. Then pick a point $a \in A$, and let $\varepsilon > 0$ be arbitrary. Then the open ball $B_\varepsilon(f(a))$ has an associated open set V . Moreover, $a \in B_\varepsilon(f(a)) \implies a \in V \cap A \implies a \in V$, so there exists an open ball $B_\delta(a) \subseteq V$. Then for any $x \in A$ with $|x-a| < \delta$, $x \in B_\delta(a) \subseteq V$, so $x \in f^{-1}(B_\varepsilon(f(a)))$, and thus $f(x) \in B_\varepsilon(f(a))$. So $\lim_{x \rightarrow a} f(x) = f(a)$. \square

When $A = \mathbb{R}^n$, this condition can be phrased as saying “the preimage of every open set is open.” Analogously, a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous if and only if the preimage of every closed set is closed. Note that it is not necessarily true that the *image* of every open set is open. For instance, the function $f(x) = x^2$ maps the open set \mathbb{R} to the set $[0, \infty)$, which is not open. However, this condition does imply that for any open set which is not also closed (the only examples are \emptyset and \mathbb{R}^n), the image is not closed. Thus, continuity allows us to infer openness *backward* through the function.

In contrast, compactness is passed *forward* through continuous functions, which is another reason that it is useful for our study.

Theorem 1.9

If $f : A \rightarrow \mathbb{R}^m$ is continuous and $A \subseteq \mathbb{R}^n$ is compact, then $f(A)$ is compact.

Proof. Pick an open cover \mathcal{O} of $f(A)$. Then by the proposition, for each open set $O \in \mathcal{O}$ there exists an open set $U \subseteq \mathbb{R}^n$ such that $U \cap A = f^{-1}(O)$. Then the collection \mathcal{U} of all such U covers A , so we pick a finite number U_1, \dots, U_n . Then the finite cover O_1, \dots, O_n cover $f(A)$. So $f(A)$ is compact. \square

One disadvantage of these definitions of continuity is that they are binary in nature: a function is either continuous or discontinuous at a certain point. The following definition allows us to measure how discontinuous a function is at a certain point.

Definition 1.19

Let $f : A \rightarrow \mathbb{R}^m$ with $A \subseteq \mathbb{R}^n$ bounded, and let $a \in A$. Define

$$M(f, a, \delta) = \sup\{f(x) : x \in A, |x-a| < \delta\}, m(f, a, \delta) = \inf\{f(y) : y \in A, |y-a| < \delta\}$$

Then the **oscillation** of f at a , denoted $o(f, a)$, is defined as

$$o(f, a) = \lim_{\delta \rightarrow 0} [M(f, a, \delta) - m(f, a, \delta)]$$

which always converges since it decreases as $\delta \rightarrow 0$ and is bounded below by 0.

In agreement with the intuition for $o(f, a)$ as measuring the discontinuity of f at a , we have the following theorem:

Theorem 1.10

A function $f : A \rightarrow \mathbb{R}^m$ with $A \subseteq \mathbb{R}^n$ bounded is continuous at $a \in A$ if and only if $o(f, a) = 0$.

Proof. (\implies) Suppose f is continuous at a . Let $\varepsilon > 0$ be arbitrary. Then there exists $\delta > 0$ such that for any $x \in A$ with $|x - a| < \delta$, we have

$$|f(x) - f(a)| < \varepsilon/2 \implies f(a) - \frac{\varepsilon}{2} < f(x) < f(a) + \frac{\varepsilon}{2}$$

Then $M(f, a, \delta) - m(f, a, \delta) < \varepsilon$. So $o(f, a) < \varepsilon$ for every positive ε , and of course $o(f, a) \geq 0$, so $o(f, a) = 0$.

(\impliedby) Suppose $o(f, a) = 0$. Then let $\varepsilon > 0$ be arbitrary. Since $\lim_{\delta \rightarrow 0} [M(f, a, \delta) - m(f, a, \delta)] = 0$, we can pick δ such that $M(f, a, \delta) - m(f, a, \delta) < \varepsilon$. Then for any $x \in A$ with $|x - a| < \delta$,

$$f(x) \leq M(f, a, \delta) < \varepsilon + m(f, a, \delta) < \varepsilon + f(a)$$

Similarly, $f(x) \geq f(a) - \varepsilon$. So $|f(x) - f(a)| < \varepsilon$. Thus f is continuous at a . \square

Proposition 1.11

Let $A \subseteq \mathbb{R}^n$ be closed, and let $f : A \rightarrow \mathbb{R}^m$ be bounded. For $\varepsilon > 0$, the set $O_\varepsilon = \{x \in A : o(f, x) \geq \varepsilon\}$ is closed.

Proof. We wish to show that $\mathbb{R}^n \setminus O_\varepsilon$ is open. Pick a point $y \in \mathbb{R}^n \setminus O_\varepsilon$. If $y \notin A$, then $y \in \mathbb{R}^n \setminus A$ open so there exists an open rectangle $B \subseteq \mathbb{R}^n \setminus A \subseteq \mathbb{R}^n \setminus O_\varepsilon$ such that $y \in B$.

If $y \in A$, then $o(f, y) < \varepsilon$. Then there exists $B_\delta(y)$ with $M(f, y, \delta) - m(f, y, \delta) < \varepsilon$. I claim that any point $z \in B_\delta(y)$ has $o(f, z) < \varepsilon$. Pick δ' small enough that $B_{\delta'}(z) \subseteq B_\delta(y)$. Then $M(f, z, \delta') \leq M(f, y, \delta)$ and $m(f, z, \delta') \geq m(f, z, \delta)$, so $M(f, z, \delta') - m(f, z, \delta') \leq M(f, y, \delta) - m(f, y, \delta) < \varepsilon$. So $o(f, z) < \varepsilon$, and thus $B_\delta(y) \subseteq \mathbb{R}^n \setminus O_\varepsilon$, so $\mathbb{R}^n \setminus O_\varepsilon$ is closed and O_ε is open. \square

Chapter 2

Differentiation

2.1 Basic Definitions

We now turn our attention to the first major topic of this book; namely, the generalization of differentiation to functions of the form $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$. To do so, first recall that $f : \mathbb{R} \rightarrow \mathbb{R}$ is differentiable at $a \in \mathbb{R}$ if there exists a number $f'(a)$ such that

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} = f'(a)$$

We cannot directly use this formula to define vector valued differentiation. First, the quotient would not even make sense when dividing vectors, and even if absolute value bars are taken, it would often be the case that this limit does not exist. However, we can rearrange this equation as

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - f'(a)h}{h} = 0$$

In other words, our new condition is that there is a linear transformation $\lambda(h) = f'(a)(h)$ such that

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - \lambda(h)}{h} = 0$$

Conceptually, this is the statement that f is approximated well near a by $f(a) + \lambda$. This interpretation extends nicely to higher dimensions:

Definition 2.1

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **differentiable** at $a \in \mathbb{R}^n$ if there exists a linear transformation $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = 0$$

In this case, λ is denoted $Df(a)$ and is called the **derivative** of f at a .

To justify uniqueness, we prove the following.

Proposition 2.1

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$ then there exists a unique linear transformation $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = 0$$

Proof. Existence follows from the definition of differentiability. Suppose that λ, μ are two linear transformations which satisfy the above. Then we have

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{|\lambda(h) - \mu(h)|}{|h|} &= \lim_{h \rightarrow 0} \frac{|\lambda(h) + f(a) - f(a+h) - \mu(h) - f(a) + f(a+h)|}{|h|} \\ &\leq \lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} + \lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \mu(h)|}{|h|} \\ &= 0 \end{aligned}$$

Picking any $x \neq 0 \in \mathbb{R}^n$, and any $t \neq 0$,

$$\frac{|\lambda(x) - \mu(x)|}{|x|} = \frac{t}{t} \frac{|\lambda(x) - \mu(x)|}{|x|} = \frac{|\lambda(tx) - \mu(tx)|}{|tx|}$$

But we just showed that

$$\lim_{t \rightarrow 0} \frac{|\lambda(tx) - \mu(tx)|}{|tx|} = 0$$

and $\frac{|\lambda(x) - \mu(x)|}{|x|}$ is constant so it must be 0. Thus

$$\frac{|\lambda(x) - \mu(x)|}{|x|} = 0 \implies \lambda = \mu \quad \square$$

We also are often interested in the matrix form of $Df(a)$, so we give it a special notation.

Definition 2.2

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable, then the **Jacobian matrix** of f is the $m \times n$ matrix

$$f'(a) := [Df(a)]$$

Lastly, we note that although many of the theorems presented in this chapter will assume that f is defined on all of \mathbb{R}^n , it is often only necessary that f is defined on an open set containing a , so we lose little generality.

2.2 Basic Theorems

As in single variable analysis, the $\varepsilon - \delta$ definition of continuity is often quite cumbersome to work with in practice. Thus, we present a number of theorems in this section which will allow us to easily prove differentiability and calculate derivatives.

Theorem 2.2: Chain Rule

Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at a , and suppose $g : \mathbb{R}^m \rightarrow \mathbb{R}^p$ is differentiable at $f(a)$. Then $g \circ f : \mathbb{R}^n \rightarrow \mathbb{R}^p$ is differentiable at a with derivative given by

$$D(g \circ f)(a) = Dg(f(a)) \circ Df(a)$$

which can also be written

$$(g \circ f)'(a) = g'(f(a)) \cdot f'(a)$$

Remark

When $n = m = p = 1$, this reduces to the single variable form of the chain rule.

Proof. Here, it will be more convenient to work with the errors of these functions relative to their derivatives:

$$\begin{cases} \varphi(x) := f(x) - f(a) - Df(a)(x - a) \\ \psi(x) := g(x) - g(f(a)) - Dg(f(a))(x - a) \\ \rho(x) := g(f(x)) - g(f(a)) - Dg(f(a))(Df(a)(x - a)) \end{cases}$$

By the definition of the derivatives, we know that

$$\lim_{x \rightarrow a} \frac{|\varphi(x)|}{|x - a|} = 0$$

and

$$\lim_{x \rightarrow f(a)} \frac{|\psi(x)|}{|x - f(a)|} = 0$$

We want to show that

$$\lim_{x \rightarrow a} \frac{|\rho(x)|}{|x - a|} = 0$$

Expanding and using linearity, we have

$$\begin{aligned} \rho(x) &= g(f(x)) - g(f(a)) - Dg(f(a))(Df(a)(x - a)) \\ &= g(f(x)) - g(f(a)) - Dg(f(a))(f(x) - f(a) - \varphi(x)) \\ &= g(f(x)) - g(f(a)) - Dg(f(a))(f(x) - f(a)) + Dg(f(a))(\varphi(x)) \\ &= \psi(f(x)) + Dg(f(a))(\varphi(x)) \end{aligned}$$

Let $\varepsilon > 0$ be arbitrary. Then there exists $\delta > 0$ such that whenever $|f(x) - f(a)| < \varepsilon$,

$$|\psi(f(x))| < \varepsilon |f(x) - f(a)|$$

Since f is continuous, there exists $\delta' > 0$ such that whenever $|x - a| < \delta'$, $|f(x) - f(a)| < \varepsilon$. Then whenever $|x - a| < \delta'$,

$$\begin{aligned} |\psi(f(x))| &< \varepsilon |f(x) - f(a)| \\ &= \varepsilon |\varphi(x) + Df(a)(x - a)| \\ &\leq \varepsilon |\varphi(x)| + \varepsilon |Df(a)(x - a)| \end{aligned}$$

By Exercise 1-10, there exists M_1 such that

$$|Df(a)(x - a)| \leq M_1|x - a|$$

so we have

$$|\psi(f(x))| \leq \varepsilon(|\varphi(x)| + M_1|x - a|)$$

Thus

$$0 \leq \frac{|\psi(f(x))|}{|x - a|} \leq \varepsilon \frac{|\varphi(x)|}{|x - a|} + \varepsilon M_1$$

so

$$0 \leq \lim_{x \rightarrow a} \frac{|\psi(f(x))|}{|x - a|} \leq \varepsilon \lim_{x \rightarrow a} \frac{|\varphi(x)|}{|x - a|} + \varepsilon M_1 = \varepsilon M_1$$

for all $\varepsilon > 0$, and thus we have

$$\lim_{x \rightarrow a} \frac{|\psi(f(x))|}{|x - a|} = 0$$

For the second term,

$$\lim_{x \rightarrow a} \frac{|Dg(f(a))(\varphi(x))|}{|x - a|} = \lim_{x \rightarrow a} \frac{|Dg(f(a))(\varphi(x))|}{|\varphi(x)|} \frac{|\varphi(x)|}{|x - a|}$$

Since $Dg(f(a))$ is linear, Exercise 1-10 tells us that there exists $M > 0$ such that for any h

$$\frac{|Dg(f(a))h|}{|h|} < M$$

so the first factor is bounded, and the second goes to zero, so we have

$$\lim_{x \rightarrow a} \frac{|Dg(f(a))(\varphi(x))|}{|x - a|} = 0$$

and thus

$$\lim_{x \rightarrow a} \frac{|\rho(x)|}{|x - a|} = 0$$

which implies that

$$D(g \circ f)(a) = Dg(f(a)) \circ Df(a)$$

□

Theorem 2.3

1. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a constant function, then

$$Df(a) = 0$$

2. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear transformation, then

$$Df(a) = f$$

3. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then f is differentiable at $a \in \mathbb{R}^n$ if and only if each component function f^i is, and in this case

$$Df(a) = (Df^1(a), \dots, Df^m(a))$$

In matrix form, $f'(a)$ is an $m \times n$ matrix with $(f^i)'(a)$ as its i th row.

4. Let $s : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the sum function, defined by $s(x, y) = x + y$. Then

$$Ds(a, b) = s$$

5. Let $p : \mathbb{R}^2 \rightarrow \mathbb{R}$ be the product function, defined by $p(x, y) = xy$. Then

$$Dp(a, b)(x, y) = bx + ay$$

1. *Proof.* Suppose f is constant. Let $a \in \mathbb{R}^n$ be arbitrary. Then

$$\lim_{x \rightarrow a} \frac{|f(x) - f(a) - 0|}{|x - a|} = \lim_{x \rightarrow a} 0 = 0$$

so $Df(a) = 0$. □

2. *Proof.* Suppose f is linear. Let $a \in \mathbb{R}^n$ be arbitrary. Then

$$\lim_{x \rightarrow a} \frac{|f(x) - f(a) - f(x - a)|}{|x - a|} = \lim_{x \rightarrow a} \frac{|f(x - a) - f(x - a)|}{|x - a|} = 0$$

so $Df(a) = f$. □

3. *Proof.* (\implies) Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$. Then any component function $f^i : \mathbb{R}^n \rightarrow \mathbb{R}$ is simply the composition $\pi^i \circ f$, where π^i is the i th projection function. π^i is linear, so by part 2 of this theorem it is also differentiable, and the chain rule tells us that $f^i = \pi^i \circ f$ is also differentiable.

(\impliedby) Now suppose each component function is differentiable at $a \in \mathbb{R}^n$, and define

$$\lambda = (Df^1(a), \dots, Df^m(a))$$

Then the function $f(a + h) - f(a) - \lambda(h)$ has components

$$(f^1(a + h) - f^1(a) - Df^1(a)(h), \dots, f^m(a + h) - f^m(a) - Df^m(a)(h))$$

so that

$$|f(a+h) - f(a) - \lambda(h)| \leq \sum_{i=1}^m |f^i(a+h) - f^i(a) - Df^i(a)(h)|$$

and thus

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} \leq \sum_{i=1}^m \lim_{h \rightarrow 0} \frac{|f^i(a+h) - f^i(a) - Df^i(a)(h)|}{|h|} = 0$$

so that

$$Df(a) = (Df^1(a), \dots, Df^m(a)) \quad \square$$

4. *Proof.* s is linear, so this follows from part 2. □

5. *Proof.* Let $\lambda(x, y) = bx + ay$. Then

$$\begin{aligned} \lim_{(h,k) \rightarrow \mathbf{0}} \frac{|p(a+h, b+k) - p(a, b) - \lambda(h, k)|}{|(h, k)|} &= \lim_{(h,k) \rightarrow \mathbf{0}} \frac{|hk|}{|(h, k)|} \\ &\leq \lim_{(h,k) \rightarrow \mathbf{0}} \frac{h^2 + k^2}{\sqrt{h^2 + k^2}} \\ &= \lim_{(h,k) \rightarrow \mathbf{0}} \sqrt{h^2 + k^2} \\ &= 0 \end{aligned} \quad \square$$

Using the sum and product functions, we can now prove the multivariate equivalent of the sum and product rules from single variable analysis.

Theorem 2.4

If $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ are differentiable at $a \in \mathbb{R}^n$, then

$$D(f+g)(a) = Df(a) + Dg(a)$$

and

$$D(fg)(a) = g(a)Df(a) + f(a)Dg(a)$$

If $g(a) \neq 0$, then

$$D(f/g)(a) = \frac{g(a)Df(a) - f(a)Dg(a)}{[g(a)]^2}$$

Proof. Note that we can express sums and products of (\mathbb{R} -valued functions) as compositions of the functions with the functions $s, p : \mathbb{R}^2 \rightarrow \mathbb{R}$ from the previous theorem.

Specifically, $f+g = s \circ (f, g)$. Then

$$\begin{aligned} D(f+g)(a) &= D(s \circ (f, g))(a) \\ &= Ds(f(a), g(a)) \circ D(f, g)(a) \\ &= s \circ (Df(a), Dg(a)) \\ &= Df(a) + Dg(a) \end{aligned}$$

Similarly, $fg = p \circ (f, g)$, Then

$$\begin{aligned}
 D(fg)(a) &= D(p \circ (f, g))(a) \\
 &= Dp(f(a), g(a)) \circ D(f, g)(a) \\
 &= Dp(f(a), g(a)) \circ (Df(a), Dg(a)) \\
 &= g(a)Df(a) + f(a)Dg(a)
 \end{aligned}$$

Finally, let $h : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ be defined by $x \mapsto 1/x$. Since we know $g(a) \neq 0$, then we have $f/g = f * (h \circ g)$. We also know from single variable calculus that $Dh(x) = -\frac{1}{x^2}$. Using the product rule we just derived, we have

$$\begin{aligned}
 D(f/g)(a) &= D(f * (h \circ g))(a) \\
 &= (h \circ g)(a)Df(a) + f(a)D(h \circ g)(a) \\
 &= \frac{Df(a)}{g(a)} + f(a)Dh(g(a))Dg(a) \\
 &= \frac{g(a)Df(a)}{[g(a)]^2} - \frac{f(a)Dg(a)}{[g(a)]^2} \\
 &= \frac{g(a)Df(a) - f(a)Dg(a)}{[g(a)]^2}
 \end{aligned}$$

□

The above theorems allow us, at least in theory, to differentiate vector-valued functions which have components given by sums, products, and quotients of the input components, as well as of single-variable differentiable functions and compositions thereof. However, using the rules above is not always the most convenient in practice.

Example 2.1

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \sin(xy^2) = \sin \circ (\pi^1 \cdot [\pi^2]^2)$$

Then we have

$$\begin{aligned}
 f'(a, b) &= \sin'(ab^2)(\pi^1 \cdot [\pi^2]^2)'(a, b) \\
 &= \cos(ab^2)[b^2(\pi^1)'(a, b) + a([\pi^2]^2)'(a, b)] \\
 &= \cos(ab^2)[b^2\pi^1 + a(2\pi^2(a, b))(\pi^2)'(a, b)] \\
 &= \cos(ab^2)[b^2\pi^1 + 2ab\pi^2] \\
 &= \cos(ab^2) \cdot (b^2, 2ab) \\
 &= (b^2 \cos(ab^2), 2ab \cos(ab^2))
 \end{aligned}$$

2.3 Partial Derivatives

Although the results of the previous section are helpful in assuring us of differentiability of functions, the application of those theorems is often not very efficient, as can be seen in the example at the end of the previous section. Thus, we instead develop the theory of partial derivatives, which will allow us to differentiate these functions much more quickly.

Definition 2.3

If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\vec{a} \in \mathbb{R}^n$, then the i -th **partial derivative** of f at \vec{a} , if it exists, is the limit

$$D_i f(\vec{a}) = \lim_{h \rightarrow 0} \frac{f(\vec{a} + h e_i) - f(\vec{a})}{h}$$

In other words, the i th partial derivative is the single variable derivative of the function $g_i(x) = f(a_1, \dots, x, \dots, a_n)$ which is produced by treating all the variables except the i th as constant.

Example 2.2

Let $f(x, y) = \sin(xy^2)$. Then by treating y as constant,

$$D_1 f(x, y) = y^2 \sin(xy^2)$$

and treating x as constant,

$$D_2 f(x, y) = 2xy \sin(xy^2)$$

Example 2.3

Let $f(x, y) = x^y$. Then treating y as constant,

$$D_1 f(x, y) = yx^{y-1}$$

Treating x as constant,

$$D_2 f(x, y) = x^y \ln x$$

Assuming that $D_i f$ exists at all points in \mathbb{R}^n , we obtain another function $\mathbb{R}^n \rightarrow \mathbb{R}$, and thus we can attempt to take another partial derivative of this function. The notation for repeated partial differentiation is "inside out," that is,

$$D_j(D_i f)(x) = D_{i,j} f(x)$$

However, the order of mixed partial derivatives is irrelevant for many common functions:

Theorem 2.5

If $D_{i,j}f$ and $D_{j,i}f$ are continuous in an open set containing \vec{a} , then

$$D_{i,j}f(\vec{a}) = D_{j,i}f(\vec{a})$$

Proof. This proof is Exercise 3-28. □

By repeatedly taking mixed partial derivatives of higher orders, we can continue to apply this theorem. In particular, if each partial derivative of f of each order is continuous, then f is said to be C^∞ . In this case, the order of partial differentiation is always irrelevant.

Theorem 2.6

Let $A \subseteq \mathbb{R}^n$. If $f : A \rightarrow \mathbb{R}$ attains a maximum (or minimum) at a point $\vec{a} \in \text{int } A$ and $D_i f(\vec{a})$ exists, then $D_i f(\vec{a}) = 0$.

Proof. Let $g_i : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$g_i(x) = (a_1, \dots, x, \dots, a_n)$$

Then g_i is defined in an open interval around a_i , and attains a maximum there, so $g_i'(a_i) = 0$, and thus $D_i f(\vec{a}) = g_i'(a_i) = 0$. □

As in single variable calculus, the above theorem only gives candidate extremal points. Moreover, we still have to check boundary points separately. However, when in single variable calculus this was only a problem of evaluating a function at 2 points, in multivariable calculus, the boundary may not be discrete at all.

2.4 Derivatives

By computing some partial derivatives of functions and comparing them to their derivatives, the reader may observe a correspondence between the two. Of course, this correspondence, which allows for the easy computation of derivatives, was our original motivation for studying partial derivatives. Thus we are retroactively justified in this study, and this correspondence can be summarized in the following theorem:

Theorem 2.7

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $\vec{a} \in \mathbb{R}^n$, then $D_j f^i(\vec{a})$ exists for $1 \leq i \leq m$, $1 \leq j \leq n$, and $f'(\vec{a})$ is the $m \times n$ matrix where $[f'(\vec{a})]_{ij} = D_j f^i(\vec{a})$.

Proof. We only need to prove this for the case $m = 1$, since we already know that the i th row of $f'(\vec{a})$ is given by $(f^i)'(a_i)$.

Fix j , and let $h : \mathbb{R} \rightarrow \mathbb{R}^n$ be defined by $h(t) = \vec{a} + te_j$. Then $D_j f(\vec{a}) = D(f \circ h)(0)$. By the chain rule,

$$\begin{aligned} D_j f(\vec{a}) &= (f \circ h)'(0) \\ &= f'(h(0))h'(0) \\ &= f'(\vec{a}) \begin{bmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 1 \end{bmatrix} \end{aligned}$$

The right side of this equation is the j th entry of $f'(\vec{a})$, showing that $D_j f(a)$ exists. This extends easily for all m . \square

While the converse of this theorem is false, we can add another condition to make it true.

Definition 2.4

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$, then f is called **continuously differentiable** at a if all $D_j f^i(x)$ exist in an open set containing a and if each function $D_j f^i$ is continuous at a .

Theorem 2.8

If $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuously differentiable at a , then $Df(a)$ exists.

Proof. Suppose f is continuously differentiable at \vec{a} . Then each $D_j f^i(\vec{a})$ exists. Define $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by

$$\lambda(x_1, \dots, x_n) = \left(\sum_{j=1}^n D_j f^1(\vec{a})x_j, \dots, \sum_{j=1}^n D_j f^m(\vec{a})x_j \right)$$

Then we have

$$\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(\vec{a} + \vec{h}) - f(\vec{a}) - \lambda(\vec{h})|}{|\vec{h}|} \leq \sum_{i=1}^m \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f^i(\vec{a} + \vec{h}) - f^i(\vec{a}) - \sum_{j=1}^n D_j f^i(\vec{a})h_j|}{|\vec{h}|}$$

Thus it is sufficient to consider the case $m = 1$. When $\vec{h} = (h_1, \dots, h_n)$, define $[\vec{h}]^k := (h_1, \dots, h_k, 0, \dots, 0) \in \mathbb{R}^n$. Then we can telescope:

$$f(\vec{a} + \vec{h}) - f(\vec{a}) = \sum_{k=1}^n f(\vec{a} + [\vec{h}]^k) - f(\vec{a} + [\vec{h}]^{k-1})$$

So

$$f(\vec{a} + \vec{h}) - f(\vec{a}) - \sum_{j=1}^n D_j f(\vec{a}) h_j = \sum_{j=1}^n \left[f(\vec{a} + [\vec{h}]^j) - f(\vec{a} + [\vec{h}]^{j-1}) - D_j f(\vec{a}) h_j \right]$$

Thus we need to prove that

$$\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{\left| f(\vec{a} + [\vec{h}]^j) - f(\vec{a} + [\vec{h}]^{j-1}) - D_j f(\vec{a}) h_j \right|}{|\vec{h}|} = 0$$

for all j . Fix some j . Then define $g_j : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by

$$g_j(x) = f(a_1 + h_1, \dots, a_{j-1} + h_{j-1}, a_j + x, a_{j+1}, \dots, a_n)$$

Since f is continuously differentiable, we can pick \vec{h} small enough that $D_j f$ exists at $\vec{a} + [\vec{h}]^{j-1}$. Then $D_j f(\vec{a} + [\vec{h}]^{j-1}) = g'_j(0)$, so we have

$$\begin{aligned} \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{\left| f(\vec{a} + [\vec{h}]^j) - f(\vec{a} + [\vec{h}]^{j-1}) - D_j f(\vec{a}) h_j \right|}{|\vec{h}|} &= \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|g_j(h_j) - g_j(0) - D_j f(\vec{a}) h_j|}{|\vec{h}|} \\ &= \lim_{h_j \rightarrow 0} \frac{|g_j(h_j) - g_j(0) - g'_j(0) h_j + g'_j(0) h_j - D_j f(\vec{a}) h_j|}{|h_j|} \\ &\leq \lim_{h_j \rightarrow 0} \frac{|g_j(h_j) - g_j(0) - g'_j(0) h_j|}{|h_j|} + \lim_{h_j \rightarrow 0} \frac{|g'_j(0) h_j - D_j f(\vec{a}) h_j|}{|h_j|} \\ &= \lim_{h_j \rightarrow 0} |g'_j(0) - D_j f(\vec{a})| \\ &= \lim_{h_j \rightarrow 0} |D_j f(\vec{a} + [\vec{h}]^{j-1}) - D_j f(\vec{a})| \\ &= 0 \end{aligned}$$

where the fourth line follows since g_j is differentiable at 0, and the last equality because $D_j f$ is continuous at a_j . Thus $Df(a) = \lambda$ exists. \square

The above theorem, in combination with the Chain Rule, allows us to derive a specific version of the Chain Rule that allows us to bypass checking for differentiability when the partial derivatives are known.

Corollary 2.9

Let $g_1, \dots, g_m : \mathbb{R}^n \rightarrow \mathbb{R}$ be continuously differentiable at a , and let $f : \mathbb{R}^m \rightarrow \mathbb{R}$ be differentiable at $(g_1(a), \dots, g_m(a))$. Let $F : \mathbb{R}^n \rightarrow \mathbb{R}$ be defined by

$$F(a) = f(g_1(a), \dots, g_m(a))$$

Then

$$D_i F(a) = \sum_{j=1}^m D_j f(g_1(a), \dots, g_m(a)) \cdot D_i g_j(a)$$

Proof. Define $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ by $g = (g_1, \dots, g_m)$. Then $F = f \circ g$. Since g_1, \dots, g_m are continuously differentiable, g is continuously differentiable, so it is differentiable. Thus the Chain Rule tells us that

$$F'(a) = (f \circ g)'(a) = f'(g(a))g'(a)$$

Matrix multiplication tells us that

$$[F'(a)]_{1i} = \sum_{j=1}^m [f'(g(a))]_{1j} [g'(a)]_{ji}$$

Moreover, Theorem 2.7 tells us that

$$\begin{aligned} [F'(a)]_{1i} &= D_i F(a) \\ [f'(g(a))]_{1j} &= D_j f(g(a)) \\ [g'(a)]_{ji} &= D_i g_j(a) = D_i g_j(a) \end{aligned}$$

Thus we conclude that

$$D_i F(a) = \sum_{j=1}^m D_j f(g_1(a), \dots, g_m(a)) \cdot D_i g_j(a) \quad \square$$

Example 2.4

Let $f(x, y, z) = xyz$, and let $g_1(a, b) = a \sin b$, $g_2(a, b) = b \cos a$, $g_3(a, b) = a^3 b$. Then

$$\begin{aligned} \frac{\partial}{\partial a}(f \circ g) \Big|_{(a,b)} &= D_1(f \circ g)(a, b) \\ &= D_1 f(g(a, b)) D_1 g_1(a, b) + D_2 f(g(a, b)) D_1 g_2(a, b) + D_3 f(g(a, b)) D_1 g_3(a, b) \\ &= a^3 b^2 \cos a \sin b - a^4 b^2 \sin b \sin a + 3a^2 b^2 \sin b \cos a \end{aligned}$$

In cases where one or more of the g_i do not explicitly depend on all of the variables, the derivatives with respect to those variables is zero.

Example 2.5

Let $f(x, y, z) = xyz$, and let $g_1(a, b) = ab$, $g_2(a) = a$, $g_3(b) = b$. Replacing D_1 with D_a for clarity, we consider

$$D_a g_3(b) = 0, D_b g_2(a) = 0$$

Thus

$$\begin{aligned} D_a(f \circ g)(a, b) &= D_1 f(g(a, b)) D_a g_1(a, b) + D_2 f(g(a, b)) D_a g_2(a) \\ &= ab^2 + ab^2 \\ &= 2ab^2 \end{aligned}$$

(This can be formally established by writing $\hat{g}_2(a, b) = a$, $\hat{g}_3(a, b) = b$, but this is generally unnecessary.)

2.5 Inverse Functions

In Exercise 2-16, we began our study of inverse functions, showing that in the case that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is differentiable with a differentiable inverse $f^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$,

$$(f^{-1})'(a) = [f'(f^{-1}(a))]^{-1}$$

However, the requirement that f has an inverse, and that both are differentiable is a relatively stringent condition. Thus, it is of interest to us to identify when the above equality may be obtained under weaker conditions. In particular, the requirement that f is invertible is a strong *global* condition. However, it can be weakened by instead requiring that f is invertible locally; that is, the restriction of f to a sufficiently small open set is invertible. Thus, it falls to us to determine the conditions where this occurs.

Consider the case of $f : \mathbb{R} \rightarrow \mathbb{R}$. We would like our conditions to be in terms of the differentiability of f , since that is what we have studied so far. One observation that we can make is that if f is strictly increasing or decreasing on a small interval, it is 1-1 on that interval. In other words, if $f'(x) > 0$ in an interval around a , then f is invertible in that interval, and similarly if $f'(x) < 0$. Moreover, if f is continuously differentiable, then $f'(a) > 0$ is sufficient to conclude that $f'(x) > 0$ in an interval around a . This leads to our multivariate generalization, but it will take some work to arrive there.

Lemma 2.10

Let $A \subseteq \mathbb{R}^n$ be a rectangle and let $f : A \rightarrow \mathbb{R}^n$ be continuously differentiable. If there is a number $M > 0$ such that $|D_j f^i(x)| \leq M$ for all $x \in \text{int } A$, then

$$|f(x) - f(y)| \leq n^2 M |x - y|$$

for all $x, y \in A$.

Proof. First, we have

$$|f(x) - f(y)| \leq \sum_{i=1}^n |f^i(x) - f^i(y)|$$

Now, let $z = y - x$ and define $h^i_z(t) = f^i(x + tz)$, so that $h^i_z(0) = f^i(x)$ and $h^i_z(1) = f^i(y)$. Since f^i is differentiable (this follows from Theorem 2.8), we know that the directional derivative $D_z f^i(x)$ exists, and moreover that $h^{i'}(t) = D_z f^i(x + tz)$ (see Exercise 2-35). Thus

$$|f^i(y) - f^i(x)| = |h^i(0) - h^i(1)| = \left| \int_0^1 h^{i'}(t) dt \right| = \left| \int_0^1 D_z f^i(x + tz) dt \right|$$

We also showed in Exercise 2-29 that D_* is linear with respect to direction, so we can expand this:

$$\begin{aligned}
\left| \int_0^1 D_z f^i(x + tz) dt \right| &= \left| \int_0^1 \sum_{j=1}^n z_j D_j f^i(x + tz) dt \right| \\
&\leq \sum_{j=1}^n \left| \int_0^1 z_j D_j f^i(x + tz) dt \right| \\
&\leq \sum_{j=1}^n |z_j| \left| \int_0^1 D_j f^i(x + tz) dt \right| \\
&\leq \sum_{j=1}^n |z_j| M \\
&\leq \sum_{j=1}^n |z_j| M \\
&= nM|y - x|
\end{aligned}$$

Thus we have

$$|f^i(y) - f^i(x)| \leq nM|y - x|$$

Combining this with our first inequality, we have

$$|f(x) - f(y)| \leq \sum_{i=1}^n |f^i(x) - f^i(y)| \leq \sum_{i=1}^n nM|y - x| = n^2M|y - x| \quad \square$$

Lemma 2.10 provides the necessary machinery to extend our result about locally invertible functions to the multivariate case:

Theorem 2.11: Inverse Function Theorem

Suppose that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuously differentiable in an open set containing a , and $\det f'(a) \neq 0$. Then there is an open set V containing a and an open set W containing $f(a)$ such that $f : V \rightarrow W$ has a continuous inverse $f^{-1} : W \rightarrow V$ which is differentiable and for all $y \in W$ satisfies

$$(f^{-1})'(y) = [f'(f^{-1}(y))]^{-1}$$

Briefly speaking, this theorem says that so long as $f'(a)$ is nonsingular, then we can find a restriction to a small open set where f is invertible and the derivative condition is met.

Proof. Let $\lambda = Df(a)$. Since $\det f'(a) \neq 0$, λ is invertible. Now suppose that the theorem is true for $\lambda^{-1} \circ f$. Then letting $\phi = (\lambda^{-1} \circ f)^{-1}$, I claim that $\phi \circ \lambda^{-1} = f^{-1}$. To see this, we check that $\phi \circ \lambda^{-1}$ is both a left and right identity:

$$\begin{aligned}
(\phi \circ \lambda^{-1}) \circ f &= (\lambda^{-1} \circ f)^{-1} \circ (\lambda^{-1} \circ f) = \text{id} \\
f \circ (\phi \circ \lambda^{-1}) &= f \circ f^{-1} \circ \lambda \circ \lambda^{-1} = \text{id}
\end{aligned}$$

Moreover, this composition is continuous and differentiable, so if the theorem holds for $\lambda^{-1} \circ f$, it holds for f . Thus it suffices to prove the case where λ is the identity.

Now we know that

$$\lim_{h \rightarrow 0} \frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = 0$$

so we can choose a small closed rectangle U containing a such that

$$\frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} < 1$$

Now suppose for contradiction that there exists $x \in U$ with $f(x) = f(a)$. Then we would have

$$\frac{|f(x) - f(a) - \lambda(x-a)|}{|x-a|} = \frac{|x-a|}{|x-a|} = 1$$

which contradicts the inequality we just established for U . So $f(x) \neq f(a)$ for all $x \neq a \in U$.

Now note that $x \mapsto \det f'(x)$ consists of sums and products of continuous functions (each $D_j f^i$ exists and is continuous since f is continuously differentiable), so it is continuous. Thus we can also choose U small enough such that $\det f'(x) \neq 0$ for $x \in U$.

Lastly, since f is continuously differentiable, we can pick U small enough such that for any i, j and $x \in U$ we have

$$|D_j f^i(x) - D_j f^i(a)| < \frac{1}{2n^2}$$

Next, let $g(x) = f(x) - x$. Then since $Df(a) = \text{id}$, for any $x \in \text{int } A$ we have

$$|D_j g^i(x)| = |D_j f^i(x) - D_j \text{id}^i(x)| = |D_j f^i(x) - D_j f^i(a)| < \frac{1}{2n^2}$$

so $|D_j g^i(x)| \leq M = 1/2n^2$ for all i, j and $x \in U$. Thus we may apply Lemma 2.10 to conclude that for any $x, y \in U$,

$$|f(x) - x - (f(y) - y)| = |g(x) - g(y)| \leq n^2 M |x - y| = \frac{|x - y|}{2}$$

Moreover, by the reverse triangle inequality,

$$|x - y| - |f(x) - f(y)| \leq |f(x) - x - (f(y) - y)|$$

so we know that for any $x, y \in U$,

$$|x - y| \leq 2|f(x) - f(y)|$$

Since U is a closed rectangle, $\partial U \subseteq U$, so for any $x \in \partial U$ we know $f(x) \neq f(a)$. Thus $f(a) \notin f(\partial U)$. Moreover, ∂U is compact, so $f(\partial U)$ is compact and there exists $d > 0$ such that $|f(a) - f(x)| \geq d$ for any $x \in \partial U$. Then define

$$W = \left\{ y : |y - f(a)| < \frac{d}{2} \right\}$$

If $y \in W$ and $x \in \partial U$, then

$$|y - f(a)| < |y - f(x)|$$

Then we show that for any $y \in W$, there exists a unique preimage $x \in \text{int } U$ with $f(x) = y$. To prove this, note that defining $g : U \rightarrow \mathbb{R}$ by

$$g(x) = |y - f(x)|^2 = \sum_{i=1}^n (y_i - f^i(x))^2$$

This function is continuous, so it achieves a minimum on U . But since $|y - f(a)| < |y - f(x)|$ for $x \in \partial U$, we know that $g(a) < g(x)$. So the minimum cannot be in ∂U . Thus there exists $x \in \text{int } U$ such that g is minimized, which allows us to conclude that $D_j g(x) = 0$ for all j . Thus

$$\sum_{i=1}^n 2(y_i - f^i(x)) D_j f^i(x) = 0$$

Since this holds for every j , we can rewrite this system of equations as

$$f'(x) \begin{bmatrix} y_1 - f^1(x) \\ \vdots \\ y_n - f^n(x) \end{bmatrix} = 0$$

But $\det f'(x) \neq 0$ so we conclude that $y_i - f^i(x) = 0$ for all i . Thus $y = f(x)$. So we know that a preimage x exists. If another preimage x_2 exists, then we have

$$|x - x_2| \leq 2|f(x) - f(x_2)| = 2|y - y| = 0$$

so $x = x_2$. Thus x is unique as well. Thus, we have shown that f is locally invertible. Letting $V = \text{int } U \cap f^{-1}(W)$, we may write that $f : V \rightarrow W$ has an inverse $f^{-1} : W \rightarrow V$. Moreover, for any $y_1, y_2 \in W$ with $x_1 = f^{-1}(y_1)$ and $x_2 = f^{-1}(y_2)$, we have

$$|f^{-1}(y_1) - f^{-1}(y_2)| = |x_1 - x_2| \leq 2|f(x_1) - f(x_2)| = 2|y_1 - y_2|$$

So f^{-1} is Lipschitz and is thus continuous.

Now we must show that f^{-1} is differentiable. Let $x \in V$, and write $\mu = Df(x)$. Let $y = f(x) \in W$. Then we show that f^{-1} is differentiable at y with $Df^{-1}(y) = \mu^{-1}$. Let $\varphi(x_1) = f(x_1) - f(x) - \mu(x_1 - x)$, such that

$$f(x_1) = f(x) + \mu(x_1 - x) + \varphi(x_1 - x)$$

Moreover, since f is differentiable at x we have

$$\lim_{x_1 \rightarrow x} \frac{|\varphi(x_1 - x)|}{|x_1 - x|} = 0$$

So

$$\mu^{-1}(f(x_1) - f(x)) = x_1 - x + \mu^{-1}(\varphi(x_1 - x))$$

or

$$x_1 = \mu^{-1}(f(x_1) - f(x)) + x - \mu^{-1}(\varphi(x_1 - x))$$

But any $y_1 \in W$ is of the form $f(x_1)$ for $x_1 \in V$, so without loss of generality we may write

$$f^{-1}(y_1) = f^{-1}(y) + \mu^{-1}(y_1 - y) - \mu^{-1}(\varphi(f^{-1}(y_1) - f^{-1}(y)))$$

and we only need to show that

$$\lim_{y_1 \rightarrow y} \frac{|\mu^{-1}(\varphi(f^{-1}(y_1) - f^{-1}(y)))|}{|y_1 - y|} = 0$$

By Exercise 1-10 the linear transformation μ^{-1} is irrelevant here and we only need to show that

$$\lim_{y_1 \rightarrow y} \frac{|\varphi(f^{-1}(y_1) - f^{-1}(y))|}{|y_1 - y|} = 0$$

We can apply a trick here, splitting the fraction:

$$\frac{|\varphi(f^{-1}(y_1) - f^{-1}(y))|}{|y_1 - y|} = \frac{|\varphi(f^{-1}(y_1) - f^{-1}(y))|}{|f^{-1}(y_1) - f^{-1}(y)|} \frac{|f^{-1}(y_1) - f^{-1}(y)|}{|y_1 - y|}$$

Since f^{-1} is continuous, $f^{-1}(y_1) \rightarrow f^{-1}(y)$ as $y_1 \rightarrow y$, so

$$\lim_{y_1 \rightarrow y} \frac{|\varphi(f^{-1}(y_1) - f^{-1}(y))|}{|f^{-1}(y_1) - f^{-1}(y)|} = \lim_{x_1 \rightarrow x} \frac{|\varphi(x_1 - x)|}{|x_1 - x|} = 0$$

and the second factor is bounded by 2, completing the proof. \square

2.6 Implicit Functions

Having now proved our major result concerning local invertibility of functions, we will apply it to the study of converting implicit function relations into explicit functions.

Example 2.6

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = x^2 + y^2 - 1$. Let C be the set of points (x, y) with $f(x, y) = 0$ (this defines a **level curve** of f). Then this curve is simply a circle of radius 1 centered at the origin.

To convert this curve into an explicit function, we attempt to answer the following question: given a point $(a, b) \in C$, do there exist intervals A around a and B around b such that for any $x \in A$ there exists exactly one $y \in B$ with $(x, y) \in C$. In the case that there is, we can then define a function $g : A \rightarrow B$ which maps each x to that unique y .

If we choose (x, y) such that $x \neq \pm 1$, then we can indeed do so. When $y > 0$, the graph of the function $g(x) = \sqrt{1 - x^2}$ traces out the upper semicircle. When $y < 0$, we instead pick $h(x) = -\sqrt{1 - x^2}$, tracing out the lower circle. In both cases, our choice of g or h is forced. However, when $x = \pm 1$, we cannot pick an interval around x where such a function can be defined.

It is also worth remarking that both g and h are differentiable.

To generalize the above discussion to multiple variables, we consider functions of the form $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$. If $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$, then we would like to find neighborhoods V around x and W around y such that any $\bar{x} \in V$ corresponds to exactly one $\bar{y} \in W$ with $f(\bar{x}, \bar{y}) = 0$, which allows us to implicitly define a function $g : V \rightarrow W$, which maps \bar{x} to \bar{y} .

Theorem 2.12: Implicit Function Theorem

Let $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be continuously differentiable in an open set around (a, b) , and suppose $f(a, b) = 0$. Let M be an $m \times m$ matrix defined by $M_{ij} = D_{n+j}f^i(a, b)$. If $\det M \neq 0$, then there is an open set $A \subseteq \mathbb{R}^n$ containing a and an open set $B \subseteq \mathbb{R}^m$ containing b , such that for any $x \in A$ there is a unique $y \in B$ such that $f(x, y) = 0$. Moreover, the function g defined by $x \mapsto y$ is differentiable.

Proof. Define $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n \times \mathbb{R}^m$ by $F(x, y) = (x, f(x, y))$. Then $F'(a, b)$ is given by a block matrix

$$F'(a, b) = \begin{bmatrix} I & O \\ O & M \end{bmatrix}$$

so $\det F'(a, b) = \det M \neq 0$. Apply the Inverse Function Theorem to produce open sets $V \subseteq \mathbb{R}^n \times \mathbb{R}^m$ containing (a, b) and $W \subseteq \mathbb{R}^n \times \mathbb{R}^m$ containing $F(a, b) = (a, 0)$. We can write $V = A \times B$ (Spivak asserts this but I'm not sure how), and thus the restriction $F : A \times B \rightarrow W$ has a differentiable inverse $h : W \rightarrow A \times B$. Moreover, since F preserves the first n coordinates, h must also, so that $h(x, y) = (x, k(x, y))$ for some differentiable function k . Then define the projection $\pi : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ by $\pi(x, y) = y$, such that $\pi \circ F = f$. Thus

$$f(x, k(x, y)) = f \circ h(x, y) = (\pi \circ F) \circ h(x, y) = \pi \circ (F \circ h)(x, y) = \pi(x, y) = y$$

Then $f(x, k(x, 0)) = 0$. So for any $x \in A$, we can pick $y = k(x, 0) \in B$, and we will have $f(x, y) = 0$. Moreover, if there exists another $y' \in B$ with $f(x, y') = 0$, then we would have

$$F(x, y') = (x, f(x, y')) = (x, 0) = (x, f(x, y)) = F(x, y)$$

But F is invertible so we cannot have $y \neq y'$. Thus our choice of y is unique, and the implicitly defined function k is differentiable. \square

Since we know that the implicitly defined g is differentiable, we can calculate its derivative. For any coordinate i , we have $f^i(x, g(x)) = 0$, so

$$D_j f^i(x, g(x)) + \sum_{\alpha=1}^m D_{n+\alpha} f^i(x, g(x)) D_j g^\alpha(x) = 0$$

which we can then solve for the various $D_j g^\alpha(x)$ by inverting M (which can be done since $\det M \neq 0$).

We can generalize the Implicit Function Theorem as follows:

Theorem 2.13

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^p$ be continuously differentiable in an open set containing a , where $p \leq n$. If $f(a) = 0$ and the $p \times n$ matrix P with $P_{ij} = D_j f^i(a)$ has rank p , then there is an open set $A \subseteq \mathbb{R}^n$ and a differentiable function $h : A \rightarrow \mathbb{R}^n$ with differentiable inverse such that $h(A)$ contains a and

$$f \circ h(x_1, \dots, x_n) = (x_{n-p+1}, \dots, x_n)$$

Note: Spivak states that A contains a . This is incorrect.

We can interpret the above theorem by saying that whenever the derivative of f has rank p , then we can find h such that $f \circ h$ acts to embed the last p coordinates of \vec{x} into \mathbb{R}^p .

Proof. Consider f as a function $\mathbb{R}^{n-p} \times \mathbb{R}^p \rightarrow \mathbb{R}^p$. Then if P has rank p , it has p linearly independent columns. Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ permute the coordinates such that those linearly independent columns are the last p columns. Taking $f \circ g$, the matrix M as defined in the Implicit Function Theorem, which is a $p \times p$ matrix with $M_{ij} = D_{n+j} (f \circ g)^i(a)$, has rank p , and thus has nonzero determinant.

Now, as in the proof of the Implicit Function Theorem, define $F : \mathbb{R}^{n-p} \times \mathbb{R}^p \rightarrow \mathbb{R}^{n-p} \times \mathbb{R}^p$ by $F(x, y) = (x, f \circ g(x, y))$. Again, $\det F'(a, b) = \det M \neq 0$, so we apply the Inverse Function Theorem to produce h which is locally an inverse of F . As in the previous proof, we have

$$(f \circ g) \circ h(x, y) = y$$

so taking $g \circ h$ produces the requested function. □

Chapter 3

Integration

3.1 Basic Definitions

The following treatment of the basic definitions of integrals over a closed rectangle $A \subseteq \mathbb{R}^n$ is rapid, as this case is similar to the single variable case of integration over an interval.

Definition 3.1

A **partition** of a closed interval $[a, b]$ is a finite sequence $\{t_0, \dots, t_k\}$, such that $a = t_0 \leq \dots \leq t_k = b$, such that $[a, b]$ is divided into k subintervals.

Definition 3.2

Let $A = [a_1, b_1] \times \dots \times [a_n, b_n] \subseteq \mathbb{R}^n$ be a closed rectangle. A partition of A is a collection of partitions $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_n)$, such that \mathcal{P}_i is a partition of $[a_i, b_i]$. If \mathcal{P}_i divides $[a_i, b_i]$ into N_i subintervals, then \mathcal{P} divides A into $N_1 \times \dots \times N_n$ **subrectangles** of \mathcal{P} . Using a slight abuse of notation, we will write $S \in \mathcal{P}$ to denote that S is a subrectangle of \mathcal{P} .

If $A \subseteq \mathbb{R}^n$ is a rectangle, $f : A \rightarrow \mathbb{R}$ is bounded, and \mathcal{P} is a partition, then we can define the maximum and minimum values for each subrectangle $S \in \mathcal{P}$:

$$m_S(f) = \inf\{f(x) : x \in S\}$$
$$M_S(f) = \sup\{f(x) : x \in S\}$$

Let $v(S)$ denote the volume of S , defined as the product of the side lengths (regardless of whether S is open or closed). Then the lower and upper sums of f with respect to \mathcal{P} are

$$L(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} m_S(f)v(S)$$
$$U(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} M_S(f)v(S)$$

Since $m_S(f) \leq M_S(f)$ for any s , we then have $L(f, \mathcal{P}) \leq U(f, \mathcal{P})$.

Definition 3.3

A partition \mathcal{P}' is called a **refinement** of a partition \mathcal{P} if each subrectangle of \mathcal{P}' is contained in a subrectangle of \mathcal{P} .

Lemma 3.1

Let \mathcal{P}' be a refinement of \mathcal{P} . Then

$$L(f, \mathcal{P}) \leq L(f, \mathcal{P}')$$

and

$$U(f, \mathcal{P}) \geq U(f, \mathcal{P}')$$

Proof. Let S be a subrectangle of \mathcal{P} . Then it contains subrectangles $S_1, \dots, S_k \in \mathcal{P}'$ which are disjoint and cover S , so that $\sum_{1 \leq i \leq k} v(S_i) = v(S)$. For each S_i , $m_{S_i}(f) \geq m_S(f)$. Thus

$$\sum_{1 \leq i \leq k} m_{S_i}(f)v(S_i) \geq m_S(f)v(S)$$

Since \mathcal{P}' refines \mathcal{P} , each subrectangle of \mathcal{P}' is contained in a subrectangle of \mathcal{P} . Thus we have

$$L(f, \mathcal{P}') = \sum_{S' \in \mathcal{P}'} m_{S'}(f)v(S') = \sum_{S \in \mathcal{P}} \sum_{1 \leq i \leq k} m_{S_i}(f)v(S_i) \geq \sum_{S \in \mathcal{P}} m_S(f)v(S) = L(f, \mathcal{P})$$

The proof for the other case is similar. □

In essence, as we refine a given partition, the upper and lower sums will grow closer to one another, and under the appropriate conditions, they will also converge to one another. This provides a candidate value for the integral of f over A ; however, it is dependent on our starting partition. Ideally, our integral may be defined independent of a particular choice of partition; to do so we must prove the following:

Corollary 3.2

If \mathcal{P} and \mathcal{P}' are partitions, then $L(f, \mathcal{P}') \leq U(f, \mathcal{P})$.

To prove this, we first introduce an auxiliary construction:

Definition 3.4

Let \mathcal{P} and \mathcal{P}' be partitions of an interval $[a, b]$. Then their **common refinement** \mathcal{Q} is the partition $\mathcal{P} \cup \mathcal{P}'$. If $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_n)$ and $\mathcal{P}' = (\mathcal{P}'_1, \dots, \mathcal{P}'_m)$ be partitions of a rectangle $A \subseteq \mathbb{R}^n$. Then the common refinement \mathcal{Q} is given by $(\mathcal{P}_1 \cup \mathcal{P}'_1, \dots, \mathcal{P}_n \cup \mathcal{P}'_m)$.

Proof. Let \mathcal{Q} be the common refinement of \mathcal{P} and \mathcal{P}' . Then by Lemma 3.1,

$$L(f, \mathcal{P}') \leq L(f, \mathcal{Q}) \leq U(f, \mathcal{Q}) \leq U(f, \mathcal{P}) \quad \square$$

Now let $U = \inf U(f, \mathcal{P})$, where the infimum is taken over all partitions \mathcal{P} of A , and let $L = \sup L(f, \mathcal{P})$. By Corollary 3.2, both U and L exist, and $L \leq U$. As mentioned above, if our continued refinements converge to a single value, then this provides a plausible definition of the integral. As Corollary 3.2 shows, this convergence is only possible if $U = L$, and it must converge to that common value. Moreover, the values of U and L are independent of our choice of partition, which allows us to define the integral:

Definition 3.5

Let $f : A \rightarrow \mathbb{R}$ be bounded, with $A \subseteq \mathbb{R}^n$ a rectangle. Then f is **integrable** if $U = L$. In this case, we denote the **integral** of f on A by $\int_A f = U = L$, which may alternatively be notated $\int_A f(x_1, \dots, x_n) dx_1 \dots dx_n$.

The following theorem gives us an equivalent criterion for integrability.

Theorem 3.3

A bounded function $f : A \rightarrow \mathbb{R}$ is integrable if and only if, for every $\varepsilon > 0$ there exists a partition \mathcal{P} of A such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

Proof. (\implies) If f is integrable then U exists, so there exists a partition \mathcal{P}_1 with $U(f, \mathcal{P}_1) \leq U + \frac{\varepsilon}{2}$. Similarly there exists \mathcal{P}_2 with $L(f, \mathcal{P}_2) \geq L - \frac{\varepsilon}{2}$. Let \mathcal{P} be the common refinement of $\mathcal{P}_1, \mathcal{P}_2$. Then

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) \leq U(f, \mathcal{P}_1) - L(f, \mathcal{P}_2) \leq U + \frac{\varepsilon}{2} - (L - \frac{\varepsilon}{2}) = \varepsilon$$

(\impliedby) By Corollary 3.2, both U and L exist. Let $\varepsilon > 0$ be arbitrary, and let \mathcal{P} be the partition produced by the condition. Then

$$U - L \leq U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

So $U - L < \varepsilon$ for all $\varepsilon > 0$ and thus $U = L$. So f is integrable over A . \square

Example 3.1

Let $f : A \rightarrow \mathbb{R}$ be constant with $f(x) = c$. Then if \mathcal{P} is a partition and $S \in \mathcal{P}$, $m_S(f) = M_S(f) = c$, so

$$\begin{aligned} U(f, \mathcal{P}) &= \sum_{S \in \mathcal{P}} m_S(f)v(S) = c \sum_{S \in \mathcal{P}} v(S) = cv(A) \\ L(f, \mathcal{P}) &= cv(A) \end{aligned}$$

so $U = L = cv(A)$ and f is integrable with $\int_A f = cv(A)$.

Example 3.2

Let $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} 0, & x \in \mathbb{Q} \\ 1, & x \notin \mathbb{Q} \end{cases}$$

If \mathcal{P} is a partition and $S \in \mathcal{P}$, by the density of \mathbb{Q} in \mathbb{R} we have $m_S(f) = 0$, and by the density of $\mathbb{I} \in \mathbb{R}$ we have $M_S(f) = 1$. So

$$U(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} M_S(f)v(S) = \sum_{S \in \mathcal{P}} v(S) = v(A)$$

$$L(f, \mathcal{P}) = 0$$

So f is not integrable over any rectangle A with $v(A) > 0$.

3.2 Measure Zero and Content Zero

In this section, we discuss the notions of measure and content zero. These quantify the concept of a set which is small enough to be insignificant in certain contexts. Moreover, in particular with the case of measure zero, this is a special case of a more general technique which serves as the formalization of volume in higher dimensions.

Definition 3.6

A subset $A \subseteq \mathbb{R}^n$ has **measure zero** if for any $\varepsilon > 0$ there exists a cover \mathcal{O} of A by closed rectangles such that $\sum_{O \in \mathcal{O}} v(O) < \varepsilon > 0$.

We may also use open rectangles rather than closed rectangles in the above.

Proposition 3.4

If a set $A \subseteq \mathbb{R}^n$ is countable, then it has measure zero.

Proof. Let $\varepsilon > 0$. Enumerate the points in A as a_1, a_2, \dots . Then for each a_i , pick a closed rectangle O_i containing a_i such that $v(O_i) < \frac{\varepsilon}{2^i}$. Then $\mathcal{O} = \{O_1, O_2, \dots\}$ covers A , and

$$\sum_{O \in \mathcal{O}} v(O) = \sum_{i=1}^{\infty} v(O_i) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon \sum_{i=1}^{\infty} \frac{1}{2^i} = \varepsilon$$

so A has measure zero. □

Importantly, \mathbb{Q} is countable, and thus has measure zero.

Theorem 3.5

Let $A = \bigcup_{i=1}^{\infty} A_i$ be a countable union of measure zero sets A_i . Then A has measure zero.

Proof. Let $\varepsilon > 0$. For each A_i , pick an open cover \mathcal{O}_i such that

$$\sum_{O \in \mathcal{O}_i} v(O) < \frac{\varepsilon}{2^i}$$

Now let $\mathcal{O} = \bigcup_{i=1}^{\infty} \mathcal{O}_i$. Then \mathcal{O} covers A , and

$$\sum_{O \in \mathcal{O}} v(O) = \sum_{i=1}^{\infty} \sum_{O \in \mathcal{O}_i} v(O) \leq \sum_{i=1}^{\infty} \frac{\varepsilon}{2^i} = \varepsilon$$

so A has measure zero. □

While sets of measure zero are important (and indeed, this notion hints at more important themes in measure theory), there are times when we would prefer to work with a finite cover rather than an open cover. This is analogous to our preference for compact sets. Thus, we have a corresponding notion of measure zero for finite covers:

Definition 3.7

A subset $A \subseteq \mathbb{R}^n$ has **content zero** if for any $\varepsilon > 0$ there exists a *finite* cover \mathcal{O} of A by closed rectangles such that

$$\sum_{O \in \mathcal{O}} v(O) < \varepsilon$$

By definition, a set having content zero is a special case of having measure zero.

Theorem 3.6

A nonsingleton interval $[a, b] \subseteq \mathbb{R}$ does not have content zero. For any finite cover $\{O_1, \dots, O_n\}$ of $[a, b]$, where each O_i is a closed interval,

$$\sum_{i=1}^n v(O_i) \geq b - a$$

Proof. Let \mathcal{O} be a finite cover. We can pick a cover $\mathcal{O}' = \{O_1 \cap [a, b], \dots, O_n \cap [a, b]\}$, which will be a cover if and only if \mathcal{O} is, and which has smaller total length, so without loss of generality we may consider \mathcal{O}' . Let t_0, \dots, t_k be the endpoints of the O'_i , with

$a = \mathcal{O}_0 \leq \dots \leq \mathcal{O}_k = b$. Then each \mathcal{O}'_i contains at least one interval $[t_{i-1}, t_i]$, and each interval is contained in at least one \mathcal{O}'_i . Then

$$\sum_{\mathcal{O}' \in \mathcal{O}'} v(\mathcal{O}') \geq \sum_{j=1}^k (t_j - t_{j-1}) = b - a$$

□

The reader should note that the above proof also shows that $[a, b]$ does not have measure zero (as long as $a < b$).

Theorem 3.7

If A is compact and has measure zero, then it has content zero.

Proof. Let $\varepsilon > 0$. There exists an open cover \mathcal{O} of A with

$$\sum_{\mathcal{O} \in \mathcal{O}} v(\mathcal{O}) < \varepsilon$$

Since A is compact, pick a finite subcover \mathcal{O}' . Then

$$\sum_{\mathcal{O}' \in \mathcal{O}'} v(\mathcal{O}') \leq \sum_{\mathcal{O} \in \mathcal{O}} v(\mathcal{O}) < \varepsilon$$

so A has content zero. □

Example 3.3

Although we pointed out earlier that \mathbb{Q} has measure zero, it does not have content zero. Let $\mathcal{O} = \{[a_i, b_i]\}$ be a finite cover of $\mathbb{Q} \cap [0, 1]$ by closed intervals. Then by the density of \mathbb{Q} , \mathcal{O} must cover $[0, 1]$. But then $\sum_{i=1}^n b_i - a_i \geq 1$, so $\mathbb{Q} \cap [0, 1]$ does not have content zero. It follows that \mathbb{Q} does not either.

3.3 Integrable Functions

In this section, we will expand on the theory of which functions may be (Riemann) integrated.

Recall that $o(f, x)$ denotes the oscillation of f at x , defined as

$$\lim_{\delta \rightarrow 0} M(x, f, \delta) - m(x, f, \delta)$$

where

$$\begin{aligned} M(x, f, \delta) &= \sup\{f(y) : |x - y| < \delta\} \\ m(x, f, \delta) &= \inf\{f(y) : |x - y| < \delta\} \end{aligned}$$

Lemma 3.8

Let A be a closed rectangle and let $f : A \rightarrow \mathbb{R}$ be a bounded function such that $o(f, x) < \varepsilon$ for all $x \in A$. Then there is a partition \mathcal{P} of A with

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon \cdot v(A)$$

Proof. For each x , because $o(f, x) < \varepsilon$ we may pick a closed rectangle U_x containing x such that $M_{U_x}(f) - m_{U_x}(f) < \varepsilon$. Then the collection of U_x covers A compact, so we can pick a finite subcover U_1, \dots, U_k . Then pick a partition \mathcal{P} such that each subrectangle of \mathcal{P} is entirely contained within one of the U_x . Then for any subrectangle $S \in \mathcal{P}$ we have

$$M_S(f) - m_S(f) \leq M_{U_x}(f) - m_{U_x}(f) < \varepsilon$$

Then

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} v(S)[M_S(f) - m_S(f)] < \varepsilon \sum_{S \in \mathcal{P}} v(S) = \varepsilon v(A) \quad \square$$

Lemma 3.9

Let \mathcal{R} be a finite collection of closed rectangles $R_1, \dots, R_k \subseteq \mathbb{R}^n$. Let $A \subseteq \mathbb{R}^n$ be a closed rectangle. Then there exists a partition \mathcal{P} of A such that for each $S \in \mathcal{P}$ and each R_i , exactly one of the following is true: $S \subseteq R_i$ or $S \cap \text{int } R_i = \emptyset$.

Proof. Let $a_{i,j}$ be the left endpoint of R_i in the j th direction and $b_{i,j}$ the right endpoint, such that

$$R_i = [a_{i,1}, b_{i,1}] \times \dots \times [a_{i,n}, b_{i,n}]$$

Let $\mathcal{P}_j = \{a_{1,j}, b_{1,j}, \dots, a_{k,j}, b_{k,j}\}$ (not necessarily in order). Suppose that when ordered, $\mathcal{P}_j = \{t_{j,1}, \dots, t_{j,2k}\}$ (note that the j has switched coordinates). Let $\mathcal{P} = (\mathcal{P}_1, \dots, \mathcal{P}_n)$. Then for each $S \in \mathcal{P}$,

$$S = [t_{1,i_1-1}, t_{1,i_1}] \times \dots \times [t_{n,i_n-1}, t_{n,i_n}]$$

for appropriately chosen i_1, \dots, i_n . Any R_j is of the form

$$R_j = [t_{1,i'_1-1}, t_{1,i'_1}] \times \dots \times [t_{n,i'_n-1}, t_{n,i'_n}]$$

for some other i'_1, \dots, i'_n . Now consider the first coordinate direction. Suppose $t_{1,i'_1} \leq t_{1,i_1-1}$. Then for any $x \in S$ and $y \in \text{int } R_j$, we have

$$y_1 < t_{1,i'_1} \leq t_{1,i_1-1} \leq x_1$$

so $x \neq y$ and thus $S \cap \text{int } R_i = \emptyset$. Similarly, if $t_{1,i_1} \leq t_{1,i'_1-1}$, then we have

$$x_1 \leq t_{1,i_1} \leq t_{1,i'_1-1} < y_1$$

so $x \neq y$ and $S \cap \text{int } R_i = \emptyset$. Thus we either immediately conclude that $S \cap \text{int } R_i = \emptyset$, or we know that

$$\begin{aligned} t_{1,i'_1} &> t_{1,i_1-1} \\ t_{1,i_1} &> t_{1,i'_1-1} \end{aligned}$$

This is equivalent to

$$\begin{aligned} t_{1,i'_1} &\geq t_{1,i_1} \\ t_{1,i_1-1} &\geq t_{1,i'_1-1} \end{aligned}$$

so we either have $S \cap \text{int } R_i = \emptyset$ or

$$t_{1,i'_1-1} \leq t_{1,i_1-1} \leq t_{1,i_1} \leq t_{1,i'_1}$$

We can apply this argument to each coordinate direction $1, \dots, n$, so that it is either the case that $S \cap \text{int } R_i = \emptyset$, or we have

$$\begin{aligned} t_{1,i'_1-1} &\leq t_{1,i_1-1} \leq t_{1,i_1} \leq t_{1,i'_1} \\ &\vdots \\ t_{n,i'_n-1} &\leq t_{n,i_n-1} \leq t_{n,i_n} \leq t_{n,i'_n} \end{aligned}$$

In this case, we have $S \subseteq R_i$. □

In particular, the above statement shows that if \mathcal{O} is a finite collection of rectangles such that their interiors cover some set $B \subseteq A \subseteq \mathbb{R}^n$, with A a closed rectangle, then there exists a partition of A such that each subrectangle is either contained in some $O \in \mathcal{O}$ or does not intersect B . Such a collection may be of interest, for instance, if B has content zero.

Theorem 3.10

Let A be a closed rectangle and let $f : A \rightarrow \mathbb{R}$ be a bounded function. Let $B = \{x : f \text{ is not continuous at } x\}$. Then f is integrable if and only if B is a set of measure zero.

Proof. (\implies) Suppose that f is integrable. Define $B_\varepsilon = \{x : o(f, x) \geq \varepsilon\}$. I claim that $B_{1/n}$ has measure zero for each n .

To see this, let \mathcal{P} be a partition of A such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \frac{\varepsilon}{n}$$

Then let \mathcal{S} be the collection of subrectangles $S \in \mathcal{P}$ such that $S \cap B_{1/n} \neq \emptyset$. Then \mathcal{S} covers $B_{1/n}$. Now, for each $S \in \mathcal{S}$ we know that $o(f, x) \geq \frac{1}{n}$ for some $x \in S$, so $M_S(f) - m_S(f) \geq \frac{1}{n}$. So

$$1 \leq n(M_S(f) - m_S(f))$$

Thus

$$\sum_{S \in \mathcal{S}} v(S) \leq \sum_{S \in \mathcal{S}} v(S)n(M_S(f) - m_S(f)) \leq n \sum_{S \in \mathcal{P}} v(S)[M_S(f) - m_S(f)] < \varepsilon$$

So $B_{\frac{1}{n}}$ has measure zero. Thus $B = \bigcup_{n=1}^{\infty} B_{1/n}$ has measure zero.

(\impliedby) Suppose that B has measure zero. Now let $\varepsilon > 0$. Suppose that $|f(x)| < M$ for all

x . Define $\varepsilon' = \varepsilon/2v(A)$. Define $B_\varepsilon := \{x : o(f, x) \geq \varepsilon'\}$. We have previously proved that a set of this form is compact. Then B_ε is compact and has measure zero, so it has content zero. Then there exists a finite cover \mathcal{O} of B_ε by the interior of closed rectangles such that

$$\sum_{O \in \mathcal{O}} v(O) < \frac{\varepsilon}{4M}$$

Apply Lemma 3.9 to produce a partition \mathcal{P}' such that the subrectangles which do not intersect B_ε may be enumerated as R_1, \dots, R_k , and $o(f, x) < \varepsilon' = \varepsilon/2v(A)$ for any x in any of those closed rectangles. Then apply Lemma 3.8 to each R_i to produce a refinement \mathcal{P}' such that for each R_i ,

$$\sum_{S \in \mathcal{P}' : S \subseteq R_i} v(S)[M_S(f) - m_S(f)] < \varepsilon'v(R_i) = \frac{\varepsilon}{2v(A)}v(R_i)$$

Now, for each subrectangle $S' \in \mathcal{P}'$, $S' \subseteq S$ for exactly one $S \in \mathcal{P}$. We either have $S \subseteq O$ for some $O \in \mathcal{O}$, or $S = R_i$ for some i . Thus either $S' \subseteq O$ for some $O \in \mathcal{O}$ or $S' \subseteq R_i$ for some i . Denote by \mathcal{L} the collection of S' such that $S' \subseteq O$ for $O \in \mathcal{O}$ and by \mathcal{R} the collection of S' such that $S' \subseteq R_i$ for some i . Then

$$\begin{aligned} U(f, \mathcal{P}') - L(f, \mathcal{P}') &= \sum_{S \in \mathcal{P}'} v(S)[M_S(f) - m_S(f)] \\ &= \sum_{S \in \mathcal{L}} v(S)[M_S(f) - m_S(f)] + \sum_{S \in \mathcal{R}} v(S)[M_S(f) - m_S(f)] \end{aligned}$$

We also have

$$\sum_{S \in \mathcal{L}} v(S)[M_S(f) - m_S(f)] \leq \sum_{O \in \mathcal{O}} v(O)[M_O(f) - m_O(f)]$$

and

$$\sum_{S \in \mathcal{R}} v(S)[M_S(f) - m_S(f)] = \sum_{i=1}^k \sum_{S' \in \mathcal{P}' : S' \subseteq R_i} v(S')[M_S(f) - m_S(f)]$$

so that

$$U(f, \mathcal{P}') - L(f, \mathcal{P}') \leq \sum_{O \in \mathcal{O}} v(O)[M_O(f) - m_O(f)] + \sum_{i=1}^k \sum_{S' \in \mathcal{P}' : S' \subseteq R_i} v(S')[M_S(f) - m_S(f)]$$

Since f is bounded by M , we must have $M_O(f) - m_O(f) \leq 2M$ for any O . Thus

$$\begin{aligned} \sum_{O \in \mathcal{O}} v(O)[M_O(f) - m_O(f)] + \sum_{i=1}^k U(f, \mathcal{P}_i) - L(f, \mathcal{P}_i) &< 2M \sum_{O \in \mathcal{O}} v(O) + \frac{\varepsilon}{2v(A)} \sum_{i=1}^k v(R_i) \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon v(A)}{2v(A)} \\ &= \varepsilon \end{aligned}$$

So f is integrable. □

We have thus presented an extremely useful criterion for determining when a function may be successfully integrated, without requiring the use of partitions to do so.

We will now progress to expanding our theory of integration from integration on rectangles to arbitrary bounded sets, which we define in terms of integrals on rectangles.

Definition 3.8

Let $C \subseteq \mathbb{R}^n$. The **characteristic function** of C is

$$\chi_C(x) = \begin{cases} 1, & x \in C \\ 0, & x \notin C \end{cases}$$

Definition 3.9

Suppose that $C \subseteq \mathbb{R}^n$ is bounded by a closed rectangle A , and $f : A \rightarrow \mathbb{R}$ is bounded. Then the integral of f on C is defined as

$$\int_C f = \int_A f\chi_C$$

provided this quantity is defined.

As we can see from the definition, $\int_C f$ is defined whenever $f\chi_C$ is integrable on A . As we prove in Exercise 3-14, the product of integrable functions is integrable, so if χ_C and f are both integrable, then $\int_C f$ is well defined. Since we are mainly concerned with integrating functions which are integrable to begin with, the main task for us is to determine when χ_C is integrable.

Theorem 3.11

If $C \subseteq A \subseteq \mathbb{R}^n$, where A is a closed rectangle, then $\chi_C : A \rightarrow \mathbb{R}$, is integrable if and only if ∂C has measure zero.

Proof. Note that whenever $x \in \partial C$, in any neighborhood of x there exists $y \in C$, such that $\chi_C(y) = 1$, and $z \notin C$, such that $\chi_C(z) = 0$. Thus χ_C is discontinuous on ∂C . On the other hand, if $x \notin \partial C$, then $x \in \text{int } A$ or $x \in \text{ext } A$. In either case, there exists a neighborhood around x such that χ_C is constant, so χ_C is continuous on $\text{int } A$ and $\text{ext } A$. Thus χ_C is discontinuous precisely on ∂C .

Since χ_C is integrable if and only if it is discontinuous on a set of measure zero, it is integrable if and only if ∂C has measure zero. □

We should note that since ∂C is closed and bounded, it also has content zero.

Definition 3.10

If C is bounded and ∂C has measure zero, then C is called **Jordan-measurable**.

Thus, for any integrable function f , $\int_C f$ is defined if C is Jordan-measurable. It is possible for $\int_C f$ to be defined in other cases (for instance, if f is identically zero), but this is of little interest to us. This also allows us to extend our definition of volume to non-rectangle sets.

Definition 3.11

The **volume** (or **content**) of a Jordan-measurable set C is defined as

$$v(C) = \int_C 1$$

Note that even if C is bounded and closed, it may not be Jordan-measurable, as we showed in Exercise 3-11. Thus, $\int_C f$ may not be defined even in the case of C open and f continuous.

3.4 Fubini's Theorem

As with our study of differentiation, we have so far been able to integrate on a case-by-case basis, and now need to produce a general method that will simplify the computation of integration in a broad class of cases. This section will develop Fubini's Theorem, which allows us to simplify computation of integrals into iterated integrals in single variables.

We will first proceed informally in order to develop intuition for the principle behind this theorem. Consider the case of a "sufficiently nice" function $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$. Then we can partition $[a, b]$ by t_0, \dots, t_k . For each t_i , the area under the graph of f above $\{t_i\} \times [c, d]$ is

$$\int_c^d f(t_i, y) dy$$

If f is nice, then we can approximate the volume under the graph of f above $[t_{i-1}, t_i] \times [c, d]$ by

$$\int_{[t_{i-1}, t_i] \times [c, d]} f \approx (t_i - t_{i-1}) \int_c^d f(x_i, y) dy$$

for any $x_i \in [t_{i-1}, t_i]$. Thus we can approximate the overall integral by

$$\int_{[a, b] \times [c, d]} f = \sum_{i=1}^k \int_{[t_{i-1}, t_i] \times [c, d]} f \approx \sum_{i=1}^k (t_i - t_{i-1}) \int_c^d f(x_i, y) dy$$

But if we consider the single variable integral $\int_a^b (\int_c^d f(x, y) dy) dx$, then this would be approximated by partitions of $[a, b]$ and sums of the form

$$\sum_{i=1}^k (t_i - t_{i-1}) \int_c^d f(x_i, y) dy$$

So it seems that for "sufficiently nice" functions, we should have

$$\int_{[a,b] \times [c,d]} f = \int_a^b \left(\int_c^d f(x,y) dy \right) dx$$

As it turns out, this indeed is the case, but the classification of which functions are "sufficiently nice" becomes a difficult problem. For instance, if $\int_c^d f(x,y) dy$ is not defined, then the above equation doesn't even make sense, although f may still be integrable.

Definition 3.12

Let $f : A \rightarrow \mathbb{R}$ be bounded with $A \subseteq \mathbb{R}^n$ a closed rectangle. Then the **lower integral** of f on A is

$$\mathbf{L} \int_A f = \sup_{\mathcal{P}} U(f, \mathcal{P})$$

and the **upper integral** is defined similarly as

$$\mathbf{U} \int_A f = \inf_{\mathcal{P}} L(f, \mathcal{P})$$

regardless of whether f is integrable on A .

Theorem 3.12: Fubini's Theorem

Let $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$ be closed rectangles, and let $f : A \times B \rightarrow \mathbb{R}$ be integrable. For any $x \in A$ define $g_x : B \rightarrow \mathbb{R}$ by $g_x(y) = f(x,y)$. Let

$$\mathcal{L}(x) = \mathbf{L} \int_B g_x = \mathbf{L} \int_B f(x,y) dy$$

$$\mathcal{U}(x) = \mathbf{U} \int_B g_x = \mathbf{U} \int_B f(x,y) dy$$

Then \mathcal{L} and \mathcal{U} are integrable on A and

$$\int_{A \times B} f = \int_A \mathcal{L} = \int_A \left(\mathbf{L} \int_B f(x,y) dy \right) dx$$

$$\int_{A \times B} f = \int_A \mathcal{U} = \int_A \left(\mathbf{U} \int_B f(x,y) dy \right) dx$$

We refer to integrals of the form $\int_A \left(\mathbf{L} \int_B f(x,y) dy \right) dx$ or $\int_A \left(\mathbf{U} \int_B f(x,y) dy \right) dx$ as **iterated integrals**.

Proof. Pick partitions \mathcal{P}_A of A and \mathcal{P}_B of B . Then $\mathcal{P} = (\mathcal{P}_A, \mathcal{P}_B)$ is a partition of $A \times B$.

Moreover, any subrectangle $S \in \mathcal{P}$ is of the form $S_A \times S_B$ for $S_A \in \mathcal{P}_A$, $S_B \in \mathcal{P}_B$. So

$$\begin{aligned} L(f, \mathcal{P}) &= \sum_{S \in \mathcal{P}} m_S(f)v(S) \\ &= \sum_{S_A \in \mathcal{P}_A, S_B \in \mathcal{P}_B} m_{S_A \times S_B}(f)v(S_A \times S_B) \\ &= \sum_{S_A \in \mathcal{P}_A} v(S_A) \left(\sum_{S_B \in \mathcal{P}_B} m_{S_A \times S_B}(f)v(S_B) \right) \end{aligned}$$

For any $x \in S_A$ we have $m_{S_A \times S_B}(f) \leq m_{S_B}(g_x)$. So for fixed $x \in S_A$,

$$\sum_{S_B \in \mathcal{P}_B} m_{S_A \times S_B}(f)v(S_B) \leq \sum_{S_B \in \mathcal{P}_B} m_{S_B}(g_x) \leq \mathbf{L} \int_B g_x = \mathcal{L}(x)$$

and thus

$$L(f, \mathcal{P}) = \sum_{S_A \in \mathcal{P}_A} v(S_A) \left(\sum_{S_B \in \mathcal{P}_B} m_{S_A \times S_B}(f)v(S_B) \right) \leq \sum_{S_A \in \mathcal{P}_A} m_{S_A}(\mathcal{L})v(S_A) = L(\mathcal{L}, \mathcal{P}_A)$$

so that

$$L(f, \mathcal{P}) \leq L(\mathcal{L}, \mathcal{P}_A) \leq U(\mathcal{L}, \mathcal{P}_A) \leq U(\mathcal{U}, \mathcal{P}_A) \leq U(f, \mathcal{P})$$

where the third inequality follows because $\mathcal{L} \leq \mathcal{U}$ and the fourth by a similar argument to what we just proved. Now, f is integrable, which means that

$$\sup L(f, \mathcal{P}) = \inf U(f, \mathcal{P}) = \int_{A \times B} f$$

So that

$$\sup L(\mathcal{L}, \mathcal{P}_A) = \inf U(\mathcal{L}, \mathcal{P}_A) = \int_{A \times B} f$$

Thus \mathcal{L} is integrable on A and

$$\int_A \mathcal{L} = \int_{A \times B} f$$

and similarly \mathcal{U} is integrable iwth

$$\int_A \mathcal{U} = \int_{A \times B} f \quad \square$$

Corollary

Under the same hypotheses,

$$\int_{A \times B} f = \int_B \left(\mathbf{L} \int_A f(x, y) dx \right) dy = \int_B \left(\mathbf{U} \int_A f(x, y) dx \right) dy$$

Proof. Analogous. □

The fact that this proof may be repeated in the other order may seem clear based on simply reading the proof. However, the important implication is that, for these sufficiently nice functions, not only may our integral be replaced with an iterated integral, but the order of the iterated integral may be changed.

Remark

If each g_x is integrable, then we may dispense with the functions \mathcal{L} and \mathcal{U} and simply write

$$\int_{A \times B} f = \int_A \left(\int_B f(x, y) \, dy \right) dx = \int_B \left(\int_A f(x, y) \, dx \right) dy$$

In particular, this is the case if f is continuous.

Alternatively, if all but a finite number of g_x are integrable, then we may still write the same, and arbitrarily define the quantity $\int_B f(x, y) \, dy$ if g_x is not integrable (since changing the value of \mathcal{L} at a finite number of points will not change its integral).

Example 3.4

Define $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ by

$$f(x, y) = \begin{cases} 1, & x \notin \mathbb{Q} \\ 1, & x \in \mathbb{Q}, y \notin \mathbb{Q} \\ 1 - \frac{1}{q}, & x = \frac{p}{q}, y \in \mathbb{Q} \end{cases}$$

where $x = p/q$ is assumed to be in lowest terms. Then f is integrable with $\int_{[0,1] \times [0,1]} f = 1$. But $\int_0^1 f(x, y) \, dy = 1$ when $x \in \mathbb{Q}$ and does not exist otherwise. So we cannot arbitrarily set the value of $\int_0^1 f(x, y) \, dy$ wherever the integral doesn't exist. For instance, defining this as zero gives Dirichlet's function, which is not integrable.

Remark

If $A = [a_1, b_1] \times \dots \times [a_n, b_n]$ and $f : A \rightarrow \mathbb{R}$ is "sufficiently nice," then repeated application of Fubini's theorem gives

$$\int_A f = \int_{a_n}^{b_n} \left(\dots \left(\int_{a_1}^{b_1} f(x_1, \dots, x_n) \, dx_1 \right) \dots \right) dx_n$$

An application of Fubini's theorem is to integrate over subsets $C \subseteq A \times B$ by appropriately setting bounds on the iterated integrals.

Example 3.5

Let

$$C = ([-1, 1] \times [-1, 1]) \setminus \{(x, y) : |(x, y)| < 1\}$$

Then

$$\int_C f = \int_{[-1, 1] \times [-1, 1]} \chi_C f$$

Assuming that f is integrable, $\chi_C f$ is integrable since C is Jordan-measurable. So we may write

$$\int_{[-1, 1] \times [-1, 1]} \chi_C f = \int_{-1}^1 \left(\int_{-1}^1 f(x, y) \chi_C(x, y) dy \right) dx$$

We have

$$\chi_C(x, y) = \begin{cases} 1, & |y| > \sqrt{1-x^2} \\ 0, & |y| \leq \sqrt{1-x^2} \end{cases}$$

so

$$\int_{-1}^1 f(x, y) \chi_C(x, y) dy = \int_{\sqrt{1-x^2}}^1 f(x, y) dy + \int_{-1}^{-\sqrt{1-x^2}} f(x, y) dy$$

and thus

$$\int_C f = \int_{-1}^1 \left(\int_{\sqrt{1-x^2}}^1 f(x, y) dy \right) dx + \int_{-1}^1 \left(\int_{-1}^{-\sqrt{1-x^2}} f(x, y) dy \right) dx$$

In general, the problem of determining bounds for arbitrary $C \subseteq A \times B$ is harder. However, one important result of Fubini's theorem is that these bounds may be set in either the $dy - dx$ order or the $dx - dy$ order, whichever is easier.

3.5 Partitions of Unity

In this section, we will discuss *partitions of unity*. These are an important tool that will help allow us to combine local results into global results, for instance when developing a theory of integration on manifolds.

Definition 3.13

Let $A \subseteq \mathbb{R}^n$. Then a **partition of unity** for A is a collection Φ of C^∞ functions φ which are defined on an open set containing A , such that

1. For all $x \in A$ and all $\varphi \in \Phi$, $0 \leq \varphi(x) \leq 1$.
2. For all $x \in A$ there exists an open set V containing x such that all but finitely many $\varphi \in \Phi$ are 0 on V .
3. For all $x \in A$ it is the case that $\sum_{\varphi \in \Phi} \varphi(x) = 1$, which is a finite sum by 2).

Definition 3.14

Let φ be a partition of unity for some $A \subseteq \mathbb{R}^n$, and let \mathcal{O} be an open cover of A . Then Φ is **subordinate** to \mathcal{O} if, for each $\varphi \in \Phi$ there exists an open set $O \in \mathcal{O}$ such that $\varphi = 0$ outside of some compact set contained in O .

Note: Spivak only requires that $\varphi = 0$ outside of a closed contained in O , but later he makes assumptions which require this set to be compact.

An important tool in proving the existence of partitions of unity will be the smooth bump functions that we proved the existence of in Exercise 2-26. Exercise 2-26 states that if $O \subseteq \mathbb{R}^n$ is open and $C \subseteq O$ is compact, then there exists a closed set $D \subseteq O$ and a C^∞ function which is positive on C and 0 outside of D .

Theorem 3.13

Let $A \subseteq \mathbb{R}^n$ and let \mathcal{O} be an open cover of A . Then there exists a partition of unity Φ for A which is subordinate to \mathcal{O} .

Proof. Case 1: A is compact.

Note that any partition of unity subordinate to a subcover of \mathcal{O} is also subordinate to \mathcal{O} . Since A is compact, we will simply assume $\mathcal{O} = \{U_1, \dots, U_k\}$ is finite. Now, we will construct a corresponding set of compact sets $D_i \subseteq U_i$ such that $\{\text{int } D_1, \dots, \text{int } D_k\}$ is also an open cover for A .

To do so, we apply an inductive argument. Let D_1, \dots, D_m be compact sets chosen so that $\{\text{int } D_1, \dots, \text{int } D_m, U_{m+1}, \dots, U_k\}$ covers A . Then let

$$C_{k+1} = A \setminus \left[\left(\bigcup_{i=1}^m \text{int } D_i \right) \cup \left(\bigcup_{j=m+2}^k U_j \right) \right]$$

Clearly U_{k+1} covers C_{k+1} , and C_{k+1} is the result of a closed set being finitely intersected with the complement of open sets, and is thus closed. So C_{k+1} is compact. Then by Exercise 1-22, there exists a compact set D_{k+1} that satisfies

$$C_{k+1} \subseteq \text{int } D_{k+1}, D_{k+1} \subseteq U_{k+1}$$

By construction, the collection of C_i will cover A , so the collection of $\text{int } D_i$ do as well, and $D_i \subseteq C_i \subseteq U_i$, so this is our desired set.

Now, by Exercise 2-26, we can construct a C^∞ "bump" function ψ_i : which is nonnegative everywhere, strictly positive on D_i , and 0 outside of a closed set contained in U_i . Now, let

$$U = \bigcup_{i=1}^k \text{int } D_i$$

$A \subseteq U$ since the $\text{int } D_i$ cover A . Moreover, for $x \in U$, x is in some D_i , and the rest are nonnegative, so

$$\sum_{i=1}^k \psi_i > 0$$

on U . So we may define $\varphi_i : U \rightarrow \mathbb{R}$ by

$$\varphi_i(x) = \frac{\psi_i(x)}{\sum_{j=1}^k \psi_j(x)}$$

which is also smooth on U . Then the collection $\{\varphi_1, \dots, \varphi_k\}$ is a partition of unity. However, it must be noted that this collection is not necessarily subordinate to \mathcal{O} . Indeed, we know that $\psi_i = 0$ outside of some closed set K contained in U_i . However, it may be the case that K is not completely contained within U . In this case, φ_1 is not even defined on K , let alone outside of it.

Moreover, it is not necessarily that case that φ_1 goes to zero at the boundary of its support. For instance, suppose $k = 1$, so that we have only a single bump function ψ_1 . Then ψ_1 goes to zero, but φ_1 is identically 1.

We can remedy this by applying Exercise 2-26 once more to construct a C^∞ function $f : U \rightarrow [0, 1]$ which is 1 on A and 0 outside of a closed set K' contained in U . Moreover, we can ensure that K' is bounded since A is, so K' is compact. Then the collection $\Phi = \{f\varphi_1, \dots, f\varphi_k\}$ is still a partition of unity for A (since $f\varphi_i = \varphi_i$ on A), but this time $f\varphi_i$ is zero outside of the compact set $K \cap K' \subseteq U_i$, so Φ is subordinate to \mathcal{O} .

Case 2: $A = \bigcup_{i=1}^\infty A_i$, where A_i is compact and $A_i \subseteq \text{int } A_{i+1}$.

Define $B_1 = A_1$ and $B_i = A_i \setminus \text{int } A_{i-1}$ for all $i \geq 2$.

Claim

Suppose $x \in A_i$. Then $x \in B_j$ for some $j \leq i$.

Proof. We prove this by induction. In the base case, $x \in A_1 \implies x \in B_1$ since $A_1 = B_1$.

For $i \geq 2$, if $x \in A_i$ then $x \in B_i$ or $x \in \text{int } A_{i-1}$. But $\text{int } A_{i-1} \subseteq A_{i-1}$, so $x \in A_{i-1}$. By the inductive hypothesis $x \in B_j$ for some $j \leq i-1 < i$. \square

By the claim, we have $A \subseteq \bigcup B_i$, and clearly $\bigcup B_i \subseteq A$, so $\bigcup B_i = A$.

Define the open cover \mathcal{O}_i by

$$\mathcal{O}_i = \begin{cases} \{O \cap \text{int } A_{i+1} : O \in \mathcal{O}\}, & i = 1, 2 \\ \{O \cap (\text{int } A_{i+1} \setminus A_{i-2}) : O \in \mathcal{O}\}, & i \geq 3 \end{cases}$$

We will construct a partition of unity for each B_i subordinate to \mathcal{O}_i .

Note that each B_i is compact, and that \mathcal{O}_i covers B_i . So by Case 1 there exists a partition of unity Φ_i for B_i subordinate to \mathcal{O}_i , where the functions are defined on some open set U_i containing B_i . Now let $x \in A$. Then $x \in B_i$ for some i . Thus $x \in A_i$. Moreover, for any $j \geq i + 2$, $x \in A_i \subseteq A_{j-2}$ so $x \notin O \cap (\text{int } A_{j+1} \setminus A_j)$ for any $O \in \mathcal{O}$, and thus $x \notin O'$ for any $O' \in \mathcal{O}_j$. Since Φ_j is subordinate to \mathcal{O}_j , $\varphi(x) = 0$ for any $\varphi \in \Phi_j$ with $j \geq i + 2$. As a result, the sum

$$\sigma(x) = \sum_{j=1}^{\infty} \sum_{\varphi \in \Phi_j} \varphi(x) = \sum_{j=1}^{i+2} \sum_{\varphi \in \Phi_j} \varphi(x) \geq 1$$

is a finite sum. Now for any $\varphi \in \Phi_j$ for any j , define $\varphi' : U_i \rightarrow \mathbb{R}$ by

$$\varphi'(x) = \frac{\varphi(x)}{\sigma(x)}$$

Moreover, the domain may be extended to $\bigcup U_i$ by simply setting $\varphi' = 0$ outside of U_i .¹ Then the collection $\Phi = \{\varphi' : \varphi \in \Phi_j, j \in \mathbb{N}\}$ satisfies conditions 1 and 3 for being a partition of unity. For condition 2, suppose $x \in A_i$. Then for each $j \leq i + 2$, there exists an open set V_j containing x such that all but finitely many $\varphi \in \Phi_j$ are zero on V_j . Let $V = V_1 \cup \dots \cup V_{i+2}$, which is open. By the argument above, $\varphi(x) = 0$ if $\varphi \in \Phi_j$ for $j > i + 2$, so there are only finitely many nonzero φ at x , and thus only finitely many φ' are nonzero at x .

So Φ is a partition of unity. Let $\varphi' \in \Phi$. Then $\varphi \in \Phi_j$ for some j . Φ_j is subordinate to \mathcal{O}_j , so there exists $O' = O \cap (\text{int } A_{j+1} \setminus A_{j-2}) \in \mathcal{O}_j$ such that φ is zero outside of a compact set contained in $O' \subseteq O$. Then φ' is also zero outside this set (assured since $U \subseteq \text{int } A_{j+1} \subseteq A$, and φ' is defined on $A \subseteq U$). So Φ' is subordinate to \mathcal{O} .

Case 3: A is open.

Let $d(x, \partial A)$ be the distance from x to ∂A as defined in Exercise 1-21 part a). Define

$$A_i : \{x : |x| \leq i, d(x, \partial A) \geq \frac{1}{i}\}$$

For any $x \in A$, $|x| < M$ for some $M \in \mathbb{N}$, and $d(x, \partial A) \geq \frac{1}{N}$ for some other $N \in \mathbb{N}$ since A is open. So $x \in A_i$ for some i and thus $A = \bigcup_{i=1}^{\infty} A_i$. So A is of the type considered in Case 2.

Case 4: A is arbitrary.

Let $B = \bigcup_{O \in \mathcal{O}} O$. Then apply Case 3 to get a partition of unity Φ for B subordinate to \mathcal{O} . Then this is also a partition of unity for A . \square

¹For details, see my answer here

Remark 3.1

For any $C \subseteq A$, if Φ is a partition of unity for A , then for $x \in C$, there exists V_x open containing x such that only finitely many φ are nonzero on V_x . Then these V_x are an open cover of C , so by compactness we only need finitely many and thus only finitely many φ are nonzero on C . In particular, if A is compact then we only need finitely many φ (this was already proved in Case 1).

Remark 3.2

Note that our proof shows that we may demand that our partition of unity is countable.

Similarly to compactness, partitions of unity will allow us to make local constructions and combine them into a global result. We will demonstrate this by extending our definition of the integral to general open sets.

Definition 3.15

Let $A \subseteq \mathbb{R}^n$ be open and let \mathcal{O} be an open cover of A . \mathcal{O} is said to be **admissible** if $O \subseteq A$ for each $O \in \mathcal{O}$ (equivalently, if $\bigcup_{O \in \mathcal{O}} O = A$).

Let Φ be a partition of unity for an open set $A \subseteq \mathbb{R}^n$ (not necessarily bounded) subordinate to an admissible open cover \mathcal{O} . Suppose also that $f : A \rightarrow \mathbb{R}$ is bounded in an open set around each point of A , and that its set of discontinuities has measure zero. Since φ has compact support, let $C_\varphi \subseteq A$ be a closed rectangle such that $\varphi = 0$ outside of C_φ ($C_\varphi \subseteq A$ is guaranteed since Φ is subordinate to \mathcal{O} , which is admissible). Since f is bounded in an open neighborhood around each point, we apply compactness to pick a finite number of them and conclude f is bounded on C_φ . $|f|$ is continuous whenever f is, so it is discontinuous on a set of measure zero and thus $|f|$ is integrable on C_φ . φ is also continuous, so $\int_{C_\varphi} \varphi |f|$ exists.

Now, by Remark 3.2, Φ is countable. So we may consider the series

$$\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi |f|$$

Suppose this series converges. Since $0 \leq \varphi \leq 1$, $\varphi |f| = |\varphi f|$, and thus by Exercise 3-6,

$$\left| \int_{C_\varphi} \varphi f \right| \leq \int_{C_\varphi} |\varphi f| = \int_{C_\varphi} \varphi |f|$$

so the series

$$\sum_{\varphi \in \Phi} \left| \int_{C_\varphi} \varphi f \right|$$

converges absolutely. This means it is independent of our ordering of Φ . Moreover, we will show that this value is also independent of our choices of Φ and \mathcal{O} , allowing us to define

this value without reference to any specific cover or partition of unity. Note that this is only the case if the series $\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi |f|$ converges; the convergence of $\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f$ is not a sufficient condition.

Definition 3.16

Let $A \subseteq \mathbb{R}^n$ be open. Suppose $f : A \rightarrow \mathbb{R}$ is bounded in an open set around each point of A , and its set of discontinuities has measure zero. Let Φ be a partition of unity for A subordinate to an admissible open cover \mathcal{O} of A . For each $\varphi \in \Phi$, let $C_\varphi \subseteq A$ be a closed rectangle such that $\varphi = 0$ outside of C_φ . Then if the series

$$\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi |f|$$

converges, then we say that f is **extended integrable** relative to Φ . Moreover, we define the extended integral of f on A relative to Φ to be

$$\text{ext}_{\Phi} \int_A f = \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f$$

Theorem 3.14

Let $A \subseteq \mathbb{R}^n$ be open. Suppose $f : A \rightarrow \mathbb{R}$ is bounded in an open set around each point of A , and its set of discontinuities has measure zero. Let Φ be a partition of unity for A subordinate to an admissible open cover \mathcal{O} of A . Let Ψ be another partition of unity for A subordinate to another admissible open cover \mathcal{O}' of A . If f is extended integrable relative to Φ , then it is extended integrable relative to Ψ , and

$$\text{ext}_{\Phi} \int_A f = \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f = \sum_{\psi \in \Psi} \int_{C_\psi} \psi f = \text{ext}_{\Psi} \int_A f$$

Proof. For each $\varphi \in \Phi$, C_φ is compact, so by Remark 3.1, only finitely many $\psi \in \Psi$ are nonzero on C_φ . Moreover, the finite sum $\sum_{\psi \in \Psi} \psi = 1$ on $C_\varphi \subseteq A$ (the subset follows since \mathcal{O} is admissible), so we have

$$\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi |f| = \sum_{\varphi \in \Phi} \left(\int_{C_\varphi} \varphi |f| \left(\sum_{\psi \in \Psi} \psi \right) \right) = \sum_{\varphi \in \Phi} \int_{C_\varphi} \sum_{\psi \in \Psi} \varphi \psi |f| = \sum_{\varphi \in \Phi} \sum_{\psi \in \Psi} \int_{C_\varphi} \varphi \psi |f|$$

Now, since the left side series converges by assumption, the right side series does as well. Since $|\varphi \psi |f|| = \varphi \psi |f|$,

$$\sum_{\varphi \in \Phi} \sum_{\psi \in \Psi} \int_{C_\varphi} \varphi \psi |f|$$

converges absolutely and thus we may switch the order of the sums:

$$\sum_{\varphi \in \Phi} \sum_{\psi \in \Psi} \int_{C_\varphi} \varphi \psi |f| = \sum_{\psi \in \Psi} \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi \psi |f|$$

Now, since ψ and φ are both zero outside of a compact set, if we let R be a rectangle containing both C_φ and C_ψ , we have

$$\int_{C_\varphi} \varphi \psi |f| = \int_R \varphi \psi |f| = \int_{C_\psi} \varphi \psi |f|$$

So

$$\sum_{\psi \in \Psi} \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi \psi |f| = \sum_{\psi \in \Psi} \sum_{\varphi \in \Phi} \int_{C_\psi} \varphi \psi |f|$$

By the argument we made at the beginning, the sum $\sum_{\varphi \in \Phi}$ is finite and equal to 1 on C_ψ , so we have

$$\sum_{\psi \in \Psi} \sum_{\varphi \in \Phi} \int_{C_\psi} \varphi \psi |f| = \sum_{\psi \in \Psi} \int_{C_\psi} \psi |f|$$

So we have shown that

$$\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi |f| = \sum_{\psi \in \Psi} \int_{C_\psi} \psi |f|$$

so the right side converges, and thus f is extended integrable relative to Ψ . Repeating this argument with f substituted for $|f|$ shows that

$$\text{ext}_{\Phi} \int_A f = \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f = \sum_{\psi \in \Psi} \int_{C_\psi} \psi f = \text{ext}_{\Psi} \int_A f \quad \square$$

By Theorem 3.14, our choice of partition is irrelevant when considering extended integrability and the value of the integral, so long as f is extended integrable with respect to some partition. Thus we may define this without reference to a particular partition.

Definition 3.17

Let $A \subseteq \mathbb{R}^n$ be open. Suppose $f : A \rightarrow \mathbb{R}$ is bounded in an open set around each point of A , and its set of discontinuities has measure zero. Then f is **extended integrable** if it is extended integrable relative to some partition of unity Φ , and the extended integral of f on A is

$$\text{ext} \int_A f = \text{ext}_{\Phi} \int_A f$$

Theorem 3.15

If $A \subseteq \mathbb{R}^n$ is open and bounded, $f : A \rightarrow \mathbb{R}$ is bounded, and its set of discontinuities is a set of measure zero, then f is extended integrable.

Proof. Let $\Phi = \{\varphi_1, \varphi_2, \dots\}$ be a countable (by Remark 3.2) partition of unity subordinate to some admissible cover \mathcal{O} . Suppose $|f| \leq M$ on A . Then let

$$S_k = \sum_{i=1}^k \int_{C_{\varphi_i}} \varphi_i |f|$$

be the k th partial sum of the corresponding infinite series. Since $\varphi_i |f| \geq 0$,

$$\int_{C_{\varphi_i}} \varphi_i |f| \geq 0$$

for each i , and thus (S_k) is increasing. Let B be some rectangle containing A . Since Φ is subordinate to an admissible cover, $C_{\varphi} \subseteq A \subseteq B$, and thus

$$\int_B \varphi = \int_{C_{\varphi}} \varphi$$

and thus

$$S_k = \sum_{i=1}^k \int_{C_{\varphi_i}} \varphi_i |f| \leq \sum_{i=1}^k M \int_{C_{\varphi_i}} \varphi = M \int_B \sum_{i=1}^k \varphi \leq M \int_B 1 = Mv(B)$$

which is constant. So (S_k) is increasing and bounded above, so it is convergent and thus f is extended integrable. \square

Theorem 3.16

Let $A \subseteq \mathbb{R}^n$ be open and Jordan-measurable. Let $f : A \rightarrow \mathbb{R}$ be bounded, and suppose its set of discontinuities has measure zero. Then

$$\int_A f = \text{ext} \int_A f$$

Proof. Note that since A is Jordan-measurable, it is bounded and thus f is extended integrable by Theorem 3.15.

Let $\varepsilon > 0$, and let Φ be an arbitrary partition of unity subordinate to an admissible open cover \mathcal{O} . Let M be such that $|f| \leq M$. Then by Exercise 3-22, there exists a compact Jordan-measurable set $C \subseteq A$ such that

$$\int_{A \setminus C} 1 < \frac{\varepsilon}{M}$$

By Remark 3.1, the subpartition Φ' of those $\varphi \in \Phi$ which are nonzero on C is finite. Then we have

$$\left| \int_A f - \text{ext} \int_A f \right| = \left| \int_A f - \sum_{\varphi \in \Phi'} \int_{C_{\varphi}} \varphi f \right|$$

Since \mathcal{O} is admissible, $C_\varphi \subseteq A$ for each $\varphi \in \Phi'$ and thus

$$\int_A \varphi f = \int_{C_\varphi} \varphi f$$

so that

$$\left| \int_A f - \sum_{\varphi \in \Phi'} \int_{C_\varphi} \varphi f \right| = \left| \int_A f - \int_A \sum_{\varphi \in \Phi'} \varphi f \right| \leq \int_A |f| \left(1 - \sum_{\varphi \in \Phi'} \varphi \right)$$

Now, we have

$$\sum_{\varphi \in \Phi} \varphi = 1$$

on A , so we may write

$$\int_A |f| \left(1 - \sum_{\varphi \in \Phi'} \varphi \right) = \int_A |f| \left(\sum_{\varphi \in \Phi} \varphi - \sum_{\varphi' \in \Phi'} \varphi' \right) \leq M \int_A \left(\sum_{\varphi \in \Phi} \varphi - \sum_{\varphi' \in \Phi'} \varphi' \right)$$

Let Ψ be the collection of $\varphi \in \Phi$ such that $\varphi \notin \Phi'$. In other words, Ψ is the collection of φ which are zero on C . Then

$$M \int_A \left(\sum_{\varphi \in \Phi} \varphi - \sum_{\varphi' \in \Phi'} \varphi' \right) = M \int_A \sum_{\psi \in \Psi} \psi$$

Since the $\psi \in \Psi$ are zero on C , they are only nonzero on $A \setminus C$. Thus

$$M \int_A \sum_{\psi \in \Psi} \psi \leq M \int_{A \setminus C} \sum_{\psi \in \Psi} \psi \leq M \int_{A \setminus C} 1 < \varepsilon$$

So

$$\int_A f = \text{ext} \int_A f \quad \square$$

3.6 Change of Variables

Consider the "u-substitution" strategy employed in single variable calculus. If $u : [a, b] \rightarrow \mathbb{R}$ is a continuously differentiable function and $f : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, then let F be such that $F' = f$. By the chain rule, $(F \circ u)' = (f \circ u)u'$. Thus

$$\int_{u(a)}^{u(b)} f = \int_{u(a)}^{u(b)} F' = F(u(b)) - F(u(a)) = \int_a^b (F \circ u)' = \int_a^b (f \circ u)u'$$

For instance, this strategy could be used computationally as follows:

$$\int_0^3 2x \sin(x^2) dx = \int_0^9 \sin u du = \cos 0 - \cos 9 = 1 - \cos 9$$

Claim

Let $u : [a, b] \rightarrow \mathbb{R}$ be continuously differentiable and injective. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be continuous. Then

$$\int_{u(a,b)} f = \int_{(a,b)} (f \circ u)|u'|$$

Proof. Since u is continuous and injective, it is strictly monotone. Suppose it is strictly increasing. Then $|u'| = u'$ and $u(b) > u(a)$, so $u(a, b) = (u(a), u(b))$ and the claim follows directly from the equality above.

If u is decreasing, then $|u'| = -u'$, and $u(b) < u(a)$, so that $u(a, b) = (u(b), u(a))$. Thus

$$\int_{u(a,b)} f = \int_{u(b)}^{u(a)} f = - \int_{u(a)}^{u(b)} f = \int_a^b -(f \circ u)u' = \int_a^b (f \circ u)|u'|$$

□

This method is invaluable for computational calculus, which motivates the development of an equivalent technique in multiple dimensions. We will do so by first proving it for linear transformations.

Lemma 3.17

Let $A \subseteq \mathbb{R}^n$ be open and let $u : A \rightarrow \mathbb{R}^n$ be injective and continuously differentiable with $\det u'(x) \neq 0$ on A . Suppose there exists an admissible cover \mathcal{O} for A such that for all $U \in \mathcal{O}$ and $f : U \rightarrow \mathbb{R}$ integrable it is the case that

$$\text{ext} \int_{u(U)} f = \text{ext} \int_U (f \circ u)|\det u'|$$

Then

$$\text{ext} \int_{u(A)} f = \text{ext} \int_A (f \circ u)|\det u'|$$

Proof. The collection $\mathcal{U} = \{u(O)\}_{O \in \mathcal{O}}$ is an open cover for $u(A)$, so we may pick a partition of unity Φ for $u(A)$ subordinate to \mathcal{U} . Suppose that $U_\varphi \in \mathcal{U}$ contains C_φ for each φ . Then we have

$$\begin{aligned} \text{ext} \int_{u(A)} f &= \sum_{\varphi \in \Phi} \int_{U_\varphi} \varphi f \\ &= \sum_{\varphi \in \Phi} \int_{u^{-1}(U_\varphi)} (\varphi \circ u)(f \circ u)|\det u'| \\ &= \sum_{\varphi \in \Phi} \int_A (\varphi \circ u)(f \circ u)|\det u'| \end{aligned}$$

Let Ψ be the partition of unity for A given by $\{\varphi \circ u\}_{\varphi \in \Phi}$. Then Ψ is thus subordinate to \mathcal{O} . Then

$$\sum_{\varphi \in \Phi} \int_A (\varphi \circ u)(f \circ u) |\det u| = \sum_{\psi \in \Psi} \int_A \psi(f \circ u) |\det u| = \text{ext} \int_A (f \circ u) |\det u| \quad \square$$

Lemma 3.18

Let $A \subseteq \mathbb{R}^n$ be open and let $u : A \rightarrow \mathbb{R}^n$ be linear, with $\det u(x) \neq 0$ on A . Then if $f : u(A) \rightarrow \mathbb{R}$ is integrable, we have

$$\text{ext} \int_{u(A)} f = \text{ext} \int_A (f \circ u) |\det u|$$

Proof. First note that by Exercise 3-35, if f is the constant function 1 and U is an open rectangle, then

$$\int_{u(U)} 1 = v(u(U)) = |\det u| v(U) = |\det u| \int_U 1 = \int_U |\det u|$$

We can make an analogous argument for u^{-1} , so for any open rectangle U we have

$$\int_{u^{-1}(U)} 1 = \int_U |\det u^{-1}| \implies \int_U 1 = \int_{u^{-1}(U)} |\det u|$$

Now suppose f is arbitrary. Let $V \subseteq u(A)$ be a rectangle, and let \mathcal{P} be a partition of V .

$$\begin{aligned} L(f, \mathcal{P}) &= \sum_{S \in \mathcal{P}} v(S) m_S(f) \\ &= \sum_{S \in \mathcal{P}} m_S(f) \int_S 1 \\ &= \sum_{S \in \mathcal{P}} m_S(f) \int_{u^{-1}(\text{int } S)} |\det u| \\ &= \sum_{S \in \mathcal{P}} \int_{u^{-1}(\text{int } S)} m_S(f) |\det u| \end{aligned}$$

For each $S \in \mathcal{P}$, define $f|_S : S \rightarrow \mathbb{R}$ to be the constant function $f|_S(x) = m_S(f)$. Then we have

$$\begin{aligned} \sum_{S \in \mathcal{P}} \int_{u^{-1}(\text{int } S)} m_S(f) |\det u| &= \sum_{S \in \mathcal{P}} \int_{u^{-1}(\text{int } V)} (f|_S \circ u) |\det u| \\ &\leq \sum_{S \in \mathcal{P}} \int_{u^{-1}(\text{int } S)} (f \circ u) |\det u| \\ &\leq \int_{u^{-1}(V)} (f \circ u) |\det u| \end{aligned}$$

So $\int_{u^{-1}(V)} (f \circ u) |\det u|$ is an upper bound for all $L(f, \mathcal{P})$, but $\int_V f$ is the least such upper bound, so we have

$$\int_V f \leq \int_{u^{-1}(V)} (f \circ u) |\det u|$$

An analogous argument shows the reverse inequality, so we conclude that

$$\int_V f = \int_{u^{-1}(V)} (f \circ u) |\det u|$$

for every $V \subseteq u(A)$ and any f .

Since A is open, and u is a continuous injection, $u(A)$ is open. So for each $\alpha \in u(A)$, we may pick $V_\alpha \subseteq u(A)$ containing α . Then the collection $\mathcal{V} = \{V_\alpha\}_{\alpha \in u(A)}$ is an admissible open cover for $u(A)$, and the hypothesis of Lemma 3.17 applies. So we conclude that

$$\text{ext} \int_{u(A)} f = \text{ext} \int_A (f \circ u) |\det u| \quad \square$$

We now progress to the general case. To do so, we will need to replace u with u' (which are equal in the linear case).

Theorem 3.19

Let $A \subseteq \mathbb{R}^n$ be open and let $u : A \rightarrow \mathbb{R}^n$ be one-to-one and continuously differentiable. Moreover, suppose that $\det u'(x) \neq 0$ on A . Then for any integrable $f : u(A) \rightarrow \mathbb{R}$, we have

$$\text{ext} \int_{u(A)} f = \text{ext} \int_A (f \circ u) |\det u'|$$

We first prove one simplifying lemma.

Lemma 3.20

Suppose that the conclusion of Theorem 3.19 holds for two change-of-variable functions $g : A \rightarrow \mathbb{R}^n$ and $h : B \rightarrow \mathbb{R}^n$. Moreover, assume that $g(A) \subseteq B$. Then the theorem holds for $h \circ g$.

Proof. We have

$$\begin{aligned} \text{ext} \int_{(h \circ g)(A)} f &= \text{ext} \int_{g(A)} (f \circ h) |\det h'| \\ &= \text{ext} \int_A (f \circ h \circ g) [|\det h'| \circ g] |\det g'| \\ &= \text{ext} \int_A (f \circ (h \circ g)) |\det (h \circ g)'| \end{aligned} \quad \square$$

Returning to the main proof,

Proof of Theorem 3.19. We induct on n . For the base case $n = 1$, we can form an admissible open cover of A by open intervals, and the result follows from the discussion beginning this section combined with Lemma 3.17.

Suppose the theorem is proved for $n - 1$. Then for n , we will attempt to find an open set $U_\alpha \subseteq A$ containing α for each $\alpha \in A$ such that

$$\int_{u(U_\alpha)} f = \int_{U_\alpha} (f \circ u) |\det u'|$$

Then the theorem follows from Lemma 3.17. Thus, fix some $\alpha \in A$. Then

$$(Du(\alpha)^{-1} \circ u)'(\alpha) = \frac{u(\alpha)^{-1}}{1} u'(\alpha) = I$$

Note that Lemma 3.18 implies that the theorem is true for $Du(\alpha)^{-1}$. If the theorem is true for $(Du(\alpha)^{-1} \circ u)$, then it follows from Lemma 3.20 that it is true for u . So we may assume that $Du(\alpha) = \text{id}$.

Define the function $h : A \rightarrow \mathbb{R}^n$ by

$$h(x) = (u_1(x), \dots, u_{n-1}(x), x_n)$$

Then $h'(\alpha) = I$. h is continuously differentiable, so there exists an open set $U' \subseteq A$ containing α where h is injective and invertible. Then define $k : h(U') \rightarrow \mathbb{R}^n$ by

$$k(x) = (x_1, \dots, x_{n-1}, u_n(h^{-1}(x)))$$

so that $u = k \circ h$. Both of these functions only change at most $n - 1$ variables, so we will be able to apply the inductive hypothesis. Afterward, we would now like to apply Lemma 3.20; however, we cannot be assured that k is injective with k' invertible.

To remedy this, note that

$$(g^n \circ h^{-1})'(\alpha) = (g^n)'(h^{-1}(h(\alpha))) \underbrace{[h'(h^{-1}(h(\alpha)))]^{-1}}_{\text{Inverse Function Thm}} = (g^n)'(\alpha)[h'(\alpha)]^{-1} = (g^n)'(\alpha)$$

So $D_n(g^n \circ h^{-1})(h(\alpha)) = D_n g^n(\alpha) = 1$, and thus $k'(h(\alpha)) = I$. So we can find an open set $V \subseteq h(U)$ containing $h(\alpha)$ where k is injective and k' is invertible. We can then restrict h to $U = k^{-1}(V)$, and then h, k satisfy the hypotheses of Lemma 3.20.

We now prove that the theorem applies to h . The proof for k is easier. Pick an open rectangle $W \subseteq U$, and suppose $W = D \times [a_n, b_n]$, with $D \subseteq \mathbb{R}^{n-1}$. Because h does not change the n -th coordinate, Fubini's Theorem gives

$$\int_{h(W)} 1 = \int_{[a_n, b_n]} \left(\int_{h(D \times \{x_n\})} 1 \, dx_1 \dots dx_{n-1} \right) dx_n$$

For each x_n , define $h_{x_n} : D \rightarrow \mathbb{R}^{n-1}$ by

$$h_{x_n}(x_1, \dots, x_{n-1}) = h(u_1(x_1, \dots, x_n), \dots, u_{n-1}(x_1, \dots, x_n))$$

so that

$$\det h'_{x_n}(x_1, \dots, x_{n-1}) = \det h'(x_1, \dots, x_n) \neq 0$$

Moreover, h_{x_n} is injective, so the inductive hypothesis applies. We also have

$$\int_{h(D \times \{x_n\})} 1 \, dx_1 \dots dx_{n-1} = \int_{h_{x_n}(D)} 1$$

Then using the inductive hypothesis, we have

$$\begin{aligned} \int_{h(W)} 1 &= \int_{[a_n, b_n]} \left(\int_{h(D \times \{x_n\})} 1 \, dx_1 \dots dx_{n-1} \right) dx_n \\ &= \int_{[a_n, b_n]} \left(\int_{h_{x_n}(D)} 1 \right) dx_n \\ &= \int_{[a_n, b_n]} \left(\int_D |\det h'_{x_n}| \right) dx_n \\ &= \int_{[a_n, b_n]} \left(\int_D |\det h'(x_1, \dots, x_n)| \right) dx_n \\ &= \int_W |\det h'| \end{aligned}$$

From the proof for Lemma 3.18, it is sufficient to prove the theorem for the constant function 1. So we conclude that the theorem holds for h . A similar argument holds for k . By Lemma 3.20, it holds for u . \square

We will now prove a simple version of an important theorem.

Theorem 3.21: Sard's Theorem

Suppose $g : A \rightarrow \mathbb{R}^n$ is continuously differentiable, with $A \subseteq \mathbb{R}^n$ open. Let $B = \{x \in A : \det g'(x) = 0\}$ be the set of critical values of g . Then $g(B)$ has measure zero.

Proof. Suppose that $U \subseteq A$ is a closed n -cube with side length ℓ . Since U is compact, each $D_j g^i$ is uniformly continuous on U . Thus there exists N large enough such that when U is divided into N^n subcubes, then for any x, y which are both in the same subcube and any i, j we have

$$|D_j g^i(y) - D_j g^i(x)| < \frac{\varepsilon}{n^2}$$

Fix some subcube S and some $x \in S$. Define $f(z) = Dg(x)(z) - g(z)$, so that its partial derivatives are bounded:

$$|D_j f^i(z)| = |D_j g^i(x) - D_j g^i(z)| < \frac{\varepsilon}{n^2}$$

Then by Lemma 2.10, for any $y \in S$ we have

$$|Dg(x)(y - x) - g(y) + g(x)| = |f(y) - f(x)| < \varepsilon |y - x| \leq \varepsilon \sqrt{n} \frac{\ell}{N}$$

We can repeat this for each x , so this holds whenever x, y are in the same subcube. If $S \cap B \neq \emptyset$, then fix $x \in S \cap B$. Then we have $\det g'(x) = 0$, so $Dg(x)(S)$ is a subset of an $n-1$ dimensional subspace V of \mathbb{R}^n . Then every point $\{g(y) - g(x) : y \in S\}$ is contained within $\varepsilon\sqrt{n}(\ell/N)$ of V , meaning that $g(S)$ is contained within $\varepsilon\sqrt{n}\ell/N$ of $V + g(x)$. Moreover, each $D_j g^i$ is uniformly continuous on S , so they are bounded by some M . Then by Lemma 2.10, we have

$$|g(x) - g(y)| < n^2 M |x - y| \leq n^2 M \sqrt{n} \frac{\ell}{N}$$

Thus $g(S)$ lies within a cylinder with height $2\varepsilon\sqrt{n}\ell/N$ and base given by a $n-1$ -sphere with radius $n^2 M \sqrt{n} \ell / N$, which has volume bounded by $C(\ell/N)^n \varepsilon$ for an appropriate constant C . Then the total volume of these cylinders (which covers $g(U \cap B)$ for each S is $C\ell^n \varepsilon$. So $g(U \cap B)$ has measure zero. Now, we can produce a cover of A (countable by Exercise 3-13) and repeat this process, so $g(B)$ is the countable union of measure zero sets, and thus measure zero. \square

Sard's Theorem, among many other applications, allows us prove Theorem 3.19 without the assumption $\det u'(x) \neq 0$. This is the content of Exercise 3-39.

Chapter 4

Integration on Chains

4.1 Algebraic Preliminaries

In this chapter, we will begin to develop our theory of integration over objects with richer structure than pure subsets of \mathbb{R}^n . This will allow us to define integrals over parameterized objects, such as line integrals and surface integrals, and we will prove a version of Stokes' Theorem for this setting. We will also set the groundwork for the development of a similar theory for manifolds in Chapter 5.

Definition 4.1

Let V be a real vector space, and let $V^k = V \times \dots \times V$ k times. A **multilinear** function $T : V^k \rightarrow \mathbb{R}$ is a function such that, for each $1 \leq i \leq k$ and each $v = (v_1, \dots, v_k) \in V^k$ the function $T_v^i : V \rightarrow \mathbb{R}$ defined by

$$T_v^i(y) = T(v_1, \dots, \underbrace{y}_i, \dots, v_k)$$

is linear. Such a function is also called a **k -tensor** on V .

Definition 4.2

The set of all k -tensors on a fixed vector space V is denoted $\mathfrak{T}^k(V)$. $\mathfrak{T}^k(V)$ is a real vector space if the operations are defined as

$$\begin{aligned}(S + T)(v_1, \dots, v_k) &= S(v_1, \dots, v_k) + T(v_1, \dots, v_k) \\ (aS)(v_1, \dots, v_k) &= a(S(v_1, \dots, v_k))\end{aligned}$$

Definition 4.3

Suppose $S \in \mathfrak{J}^k(V)$ and $T \in \mathfrak{J}^l(V)$. Then the **tensor product** of S and T is a $k + l$ -tensor $S \otimes T$ defined by

$$(S \otimes T)(v_1, \dots, v_k, v_{k+1}, \dots, v_{k+l}) = S(v_1, \dots, v_k) \cdot T(v_{k+1}, \dots, v_{k+l})$$

Note that the tensor product is clearly not commutative. Because tensors are maps into \mathbb{R} , we may use properties of \mathbb{R} to derive similar properties for tensors.

Proposition 4.1

The following are properties of the tensor product:

1. $(S_1 + S_2) \otimes T = S_1 \otimes T + S_2 \otimes T$
2. $S \otimes (T_1 + T_2) = S \otimes T_1 + S \otimes T_2$
3. $(aS) \otimes T = S \otimes (aT) = a(S \otimes T)$
4. $S \otimes (T \otimes U) = (S \otimes T) \otimes U$

Proof. 1. Let $S_1, S_2 \in \mathfrak{J}^k(V)$ and $T \in \mathfrak{J}^l(V)$. Let $v = (v_1, \dots, v_{k+l}) \in V^{k+l}$. Let $v^k = (v_1, \dots, v_k)$ and $v^l = (v_{k+1}, \dots, v_{k+l})$. Then

$$\begin{aligned} ((S_1 + S_2) \otimes T)(v) &= (S_1 + S_2)(v^k) \cdot T(v^l) \\ &= (S_1(v^k) + S_2(v^k)) \cdot T(v^l) \\ &= S_1(v^k) \cdot T(v^l) + S_2(v^k) \cdot T(v^l) \\ &= (S_1 \otimes T)(v) + (S_2 \otimes T)(v) \\ &= (S_1 \otimes T + S_2 \otimes T)(v) \end{aligned}$$

2. Let $S \in \mathfrak{J}^k(V)$ and $T_1, T_2 \in \mathfrak{J}^l(V)$. Using the same notation,

$$\begin{aligned} (S \otimes (T_1 + T_2))(v) &= S(v^k) \cdot (T_1 + T_2)(v^l) \\ &= S(v^k) \cdot (T_1(v^l) + T_2(v^l)) \\ &= S(v^k) \cdot T_1(v^l) + S(v^k) \cdot T_2(v^l) \\ &= (S \otimes T_1)(v) + (S \otimes T_2)(v) \\ &= (S \otimes T_1 + S \otimes T_2)(v) \end{aligned}$$

3. Let $a \in \mathbb{R}$, $S \in \mathfrak{J}^k(V)$, and $T \in \mathfrak{J}^l(V)$. Then

$$\begin{aligned} ((aS) \otimes T)(v) &= (aS)(v^k) \cdot T(v^l) \\ &= a(S(v^k) \cdot T(v^l)) \\ &= a(S \otimes T)(v) \\ &= S(v^k) \cdot a(T(v^l)) \\ &= (S \otimes (aT))(v) \end{aligned}$$

4. Let $S \in \mathfrak{J}^k(V)$, $T \in \mathfrak{J}^l(V)$, and $U \in \mathfrak{J}^m(V)$. Let $v = (v_1, \dots, v_{k+l+m})$, and let $v^k = (v_1, \dots, v_k)$, $v^l = (v_{k+1}, \dots, v_{k+l})$, and $v^m = (v_{k+l+1}, \dots, v_{k+l+m})$. Then

$$\begin{aligned}
(S \otimes (T \otimes U))(v) &= S(v^k) \cdot (T \otimes U)(v^l, v^m) \\
&= S(v^k) \cdot (T(v^l) \cdot U(v^m)) \\
&= (S(v^k) \cdot T(v^l)) \cdot U(v^m) \\
&= (S \otimes T)(v^k, v^l) \cdot U(v^m) \\
&= ((S \otimes T) \otimes U)(v)
\end{aligned}$$

□

Since the tensor product is associative, we will drop the parentheses in general. Note that we already know how to describe $\mathfrak{J}^1(V)$: since it is the set of all linear maps from $V \rightarrow \mathbb{R}$, it is precisely the dual space V^* . We can use this to help us understand higher order $\mathfrak{J}^k(V)$.

Theorem 4.2

Let v_1, \dots, v_n be a basis for V . Let $\varphi_1, \dots, \varphi_n$ be the natural dual basis given by $\varphi_i(v_j) = \delta_{ij}$. Then the set of k -tensors of the form

$$\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}$$

where $1 \leq i_j \leq n$ for each index, is a basis of $\mathfrak{J}^k(V)$.

Proof. We first show that this collection spans $\mathfrak{J}^k(V)$.

Let $T \in \mathfrak{J}^k(V)$. Suppose that $w_1, \dots, w_k \in V$, and $w_i = \sum_{j=1}^n a_{i,j} v_j$. Then

$$\begin{aligned}
T(w_1, \dots, w_k) &= T\left(\sum_{j_1=1}^n a_{1,j_1} v_{j_1}, w_2, \dots, w_k\right) \\
&= \sum_{j_1=1}^n a_{1,j_1} T(v_{j_1}, w_2, \dots, w_k) \\
&\vdots \\
&= \sum_{j_1=1}^n \dots \sum_{j_k=1}^n a_{1,j_1} \dots a_{k,j_k} T(v_{j_1}, \dots, v_{j_k}) \tag{*}
\end{aligned}$$

Now, we have

$$\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}(v_{j_1}, \dots, v_{j_k}) = \delta_{i_1, j_1} \dots \delta_{i_k, j_k} = \begin{cases} 1, & i_1 = j_1, \dots, i_k = j_k \\ 0 & \end{cases} \tag{**}$$

Since $\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k} \in \mathfrak{J}^k(V)$, we can use (*) and (**) to conclude that

$$\begin{aligned} \varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}(w_1, \dots, w_k) &= \sum_{j_1=1}^n \dots \sum_{j_k=1}^n a_{1,j_1} \dots a_{k,j_k} (\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k})(v_{j_1}, \dots, v_{j_k}) \\ &= \sum_{j_1=1}^n \dots \sum_{j_{k-1}=1}^n a_{1,j_1} \dots a_{k-1,j_{k-1}} a_{k,i_k} (\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k})(v_{j_1}, \dots, v_{i_k}) \\ &\quad \vdots \\ &= a_{1,i_1} \dots a_{k,i_k} \end{aligned}$$

Substituting into (*), we have

$$T(w_1, \dots, w_k) = \sum_{j_1=1}^n \dots \sum_{j_k=1}^n T(v_{j_1}, \dots, v_{j_k}) (\varphi_{j_1} \otimes \dots \otimes \varphi_{j_k})(w_1, \dots, w_k)$$

So

$$T = \sum_{j_1=1}^n \dots \sum_{j_k=1}^n T(v_{j_1}, \dots, v_{j_k}) (\varphi_{j_1} \otimes \dots \otimes \varphi_{j_k})$$

so T is a linear combination of the $\varphi_{j_1} \otimes \dots \otimes \varphi_{j_k}$.

To show that the $\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}$ are linearly independent, suppose that

$$\sum_{i_1=1}^n \dots \sum_{i_k=1}^n a_{i_1, \dots, i_k} (\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}) = 0$$

Then plugging in some combination of basis vectors $(v_{j_1}, \dots, v_{j_k})$, by (**) we have

$$0 = \sum_{i_1=1}^n \dots \sum_{i_k=1}^n a_{i_1, \dots, i_k} (\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k})(v_{j_1}, \dots, v_{j_k}) = a_{j_1, \dots, j_k}$$

Repeating this with each combination of basis vectors shows that the linear combination is trivial. So the $\varphi_{i_1} \otimes \dots \otimes \varphi_{i_k}$ are linearly independent and thus a basis. \square

Recall that if $T : V \rightarrow W$ is a linear transformation, then its adjoint $T^* : W^* \rightarrow V^*$ is the linear operator defined such that for any $\Phi \in W^*$ it is the case that $T^*(\Phi) = \Phi \circ T$. Then we can extend this notion to arbitrary $\mathfrak{J}^k(V)$.

Definition 4.4

Let $f : V \rightarrow W$ be linear. Then define the (k -tensor) **pullback** of f to be the linear transformation $f^* : \mathfrak{J}^k(W) \rightarrow \mathfrak{J}^k(V)$ by

$$(f^*(T))(v_1, \dots, v_k) = T(f(v_1), \dots, f(v_k))$$

where $T \in \mathfrak{J}^k(W)$ and $v_1, \dots, v_k \in V$.

Proposition 4.3

If $S \in \mathfrak{J}^k(V)$ and $T \in \mathfrak{J}^l(V)$, and $f : V \rightarrow W$, then

$$f^*(S \otimes T) = f^*S \otimes f^*T$$

Proof. Let $v^k = (v_1, \dots, v_k) \in V^k$ and $v^l = (v_{k+1}, \dots, v_{k+l}) \in V^l$. Then

$$\begin{aligned}
f^*(S \otimes T)(v^k, v^l) &= (S \otimes T)(f(v_1), \dots, f(v_k), f(v_{k+1}), \dots, f(v_{k+l})) \\
&= S(f(v_1), \dots, f(v_k)) \cdot T(f(v_{k+1}), \dots, f(v_{k+l})) \\
&= f^*S(v^k) \cdot f^*T(v^l) \\
&= (f^*S \otimes f^*T)(v^k, v^l)
\end{aligned}$$

□

An example of a k -tensor which is not a linear functional is the dot product on \mathbb{R}^n , which is a 2-tensor. We can use this language to make an equivalent definition for arbitrary real inner products.

Definition 4.5

An **inner product** on a real vector space V is a 2-tensor $T \in \mathfrak{J}^2(V)$ which satisfies the following:

- $T(v, w) = T(w, v)$ (symmetric)
- $T(v, v) > 0$ if $v \neq 0$ (positive definite)

We can similarly reproduce some theorems from linear algebra.

Definition 4.6

A basis v_1, \dots, v_n for a real vector space V is **orthonormal** with respect to an inner product $T \in \mathfrak{J}^2(V)$ if $T(v_i, v_j) = \delta_{ij}$.

Theorem 4.4

For any inner product T on V , there is an orthonormal basis with respect to T .

Proof. Pick a basis and apply Gram-Schmidt. □

Corollary 4.5

If T is an inner product on V , then there exists an isomorphism $f : \mathbb{R}^n \rightarrow V$ such that $T(f(x), f(y)) = x \cdot y$, or equivalently so that f^*T is the dot product on \mathbb{R}^n .

Proof. Let v_1, \dots, v_k be an orthonormal basis for T . Define f by $f(e_i) = v_i$. Then if $x = (a_1, \dots, a_n)$ and $y = (b_1, \dots, b_n)$, we have

$$\begin{aligned} T(f(x), f(y)) &= T\left(\sum_{i=1}^n a_i v_i, \sum_{j=1}^n b_j v_j\right) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i b_j T(v_i, v_j) \\ &= \sum_{i=1}^n \sum_{j=1}^n a_i b_j \delta_{ij} \\ &= \sum_{i=1}^n a_i b_i \\ &= x \cdot y \end{aligned}$$

□

Suppose we consider a square $n \times n$ matrix as a vector whose entries are column vectors. That is, we will associate $M_{n \times n}(\mathbb{R})$ with $(\mathbb{R}^n)^n$. Then $\det : M_{n \times n}(\mathbb{R}) \rightarrow \mathbb{R}$ may be considered as a k -tensor for \mathbb{R}^n . Recall that one definition of the determinant defines it as the unique alternating multilinear map with $\det I = 1$. Let us attempt to generalize this notion.

Definition 4.7

A k -tensor $T \in \mathfrak{J}^k(V)$ is **alternating** if, for every pair $i < j$, we have

$$T(v_1, \dots, v_k) = -T(v_1, \dots, v_{i-1}, v_j, v_{i+1}, \dots, v_{j-1}, v_i, v_{j+1}, \dots, v_k)$$

In other words, switching the role of two entries also switches the sign of T .

Definition 4.8

The set of all alternating k -tensors on V is denoted $A^k(V)$.^a

^aSpivak uses the notation $\Lambda^k(V)$, but writes in the Addenda that $\Omega^k(V)$ should be used instead. This definition, if V is finite dimensional, is naturally isomorphic to $\wedge(V^*)$. See here for why neither of these are quite accurate.

One can quickly verify that $A^k(V)$ is a subspace of $\mathfrak{J}^k(V)$. The close relationship of alternating tensors with signed quantities will help us to define oriented objects. Due to this, it is of interest to us to investigate how to consistently represent elements of $A^k(V)$.

Recall that the **sign** of a permutation σ , denoted $\text{sgn } \sigma$, is $+1$ if σ is even (that is, it is composed of an even number of transpositions), and -1 if it is odd.

Definition 4.9

Let $T \in \mathfrak{J}^k(V)$. Then $\text{Alt}(T) \in \mathfrak{J}^k(V)$ is defined by

$$\text{Alt}(T)(v_1, \dots, v_k) = \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot T(v_{\sigma(1)}, \dots, v_{\sigma(k)})$$

We can see Alt as a kind of projection from $\mathfrak{J}^k(V)$ into $A^k(V)$:

Theorem 4.6

Let V be a real vector space.

1. If $T \in \mathfrak{J}^k(V)$, then $\text{Alt}(T) \in A^k(V)$.
2. If $\omega \in A^k(V)$, then $\text{Alt}(\omega) = \omega$.
3. If $T \in \mathfrak{J}^k(V)$, then $\text{Alt}(\text{Alt}(T)) = \text{Alt}(T)$.

Proof. 1. Fix i, j , and let (i, j) be the transposition of i and j . For each $\sigma \in S_k$, write $\sigma' = \sigma \cdot (i, j)$. We have $S_k(i, j) = S_k$. So

$$\begin{aligned} \text{Alt}(T)(v_1, \dots, v_j, \dots, v_i, \dots, v_k) &= \text{Alt}(T)(v_{(i,j)(1)}, \dots, v_{(i,j)(k)}) \\ &= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot T(v_{\sigma((i,j)(1))}, \dots, v_{\sigma((i,j)(k))}) \\ &= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot T(v_{\sigma'(1)}, \dots, v_{\sigma'(k)}) \\ &= \frac{1}{k!} \sum_{\sigma' \in S_k(i,j)} -\text{sgn } \sigma' \cdot T(v_{\sigma'(1)}, \dots, v_{\sigma'(k)}) \\ &= -\frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot T(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= -\text{Alt}(T)(v_1, \dots, v_k) \end{aligned}$$

2. Let ω be alternating. For a transposition $\sigma = (i, j)$, we have

$$\omega(v_{\sigma(1)}, \dots, v_{\sigma(k)}) = -\omega(v_1, \dots, v_k) = \text{sgn } \sigma \cdot \omega(v_1, \dots, v_k) \quad (*)$$

For arbitrary permutations σ , σ can be decomposed into a product of transpositions $\sigma_1, \dots, \sigma_m$. Since $\text{sgn}(\sigma_1 \circ \dots \circ \sigma_m) = \text{sgn } \sigma_1 \cdots \text{sgn } \sigma_m$, we simply apply $(*)$ m times

to see that (*) holds when σ is arbitrary. Now,

$$\begin{aligned}
 \text{Alt}(\omega)(v_1, \dots, v_k) &= \frac{1}{k!} \sum_{\sigma \in S_k} \omega(v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\
 &= \frac{1}{k!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot \text{sgn } \sigma \cdot \omega(v_1, \dots, v_k) \\
 &= \frac{1}{k!} \sum_{\sigma \in S_k} \omega(v_1, \dots, v_k) \\
 &= \omega(v_1, \dots, v_k)
 \end{aligned}$$

3. Follows from 1) and 2). □

One way of describing alternating tensors would be to produce a basis of $A^k(V)$. Note that we cannot necessarily apply Theorem 4.2. This is because if $\omega \in A^k(V)$ and $\eta \in A^l(V)$, it is not necessarily the case that $\omega \otimes \eta$ is alternating (consider a transposition which swaps entries in the ω and η domains). Thus, we will need to define an analogous product which takes alternating tensors to alternating tensors.

Definition 4.10

Let $\omega \in A^k(V)$ and $\eta \in A^l(V)$. Then the **wedge product** of ω and η , denoted $\omega \wedge \eta \in A^{k+l}(V)$, is defined by

$$\omega \wedge \eta = \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta)$$

We can prove properties of \wedge using similar methods as we did for \otimes .

Proposition 4.7

Let $\omega, \omega_1, \omega_2 \in A^k(V)$, $\eta, \eta_1, \eta_2 \in A^l(V)$, and $a \in \mathbb{R}$. Then

1. $(\omega_1 + \omega_2) \wedge \eta = \omega_1 \wedge \eta + \omega_2 \wedge \eta$
2. $\omega \wedge (\eta_1 + \eta_2) = \omega \wedge \eta_1 + \omega \wedge \eta_2$
3. $(a\omega) \wedge \eta = \omega \wedge (a\eta) = a(\omega \wedge \eta)$

Proof. 1. Let $v^k = (v_1, \dots, v_k) \in V^k$ and $v^l = (v_{k+1}, \dots, v_{k+l}) \in V^l$. Write $\sigma(v^k, v^l) =$

$(v_{\sigma(1)}, \dots, v_{\sigma(k+l)})$. Then

$$\begin{aligned}
[(\omega_1 + \omega_2) \wedge \eta](v^k, v^l) &= \frac{(k+l)!}{k!l!} \text{Alt}((\omega_1 + \omega_2) \otimes \eta)(v^k, v^l) \\
&= \frac{(k+l)!}{k!l!} \text{Alt}(\omega_1 \otimes \eta + \omega_2 \otimes \eta)(v^k, v^l) \\
&= \frac{(k+l)!}{k!l!} \cdot \frac{1}{(k+l)!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot (\omega_1 \otimes \eta + \omega_2 \otimes \eta)(\sigma(v^k, v^l)) \\
&= \frac{(k+l)!}{k!l!} \cdot \frac{1}{(k+l)!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot (\omega_1 \otimes \eta(\sigma(v^k, v^l)) + \omega_2 \otimes \eta(\sigma(v^k, v^l))) \\
&= \frac{(k+l)!}{k!l!} (\text{Alt}(\omega_1 \otimes \eta) + \text{Alt}(\omega_2 \otimes \eta)) \\
&= \omega_1 \wedge \eta + \omega_2 \wedge \eta
\end{aligned}$$

2. Analogous.

3. We prove the first and third expressions are equal. The other equality is proved analogously. Then

$$\begin{aligned}
((a\omega) \wedge \eta)(v^k, v^l) &= \frac{(k+l)!}{k!l!} \text{Alt}((a\omega) \otimes \eta)(v^k, v^l) \\
&= \frac{(k+l)!}{k!l!} \text{Alt}(a(\omega \otimes \eta))(v^k, v^l) \\
&= \frac{(k+l)!}{k!l!} \cdot \frac{1}{(k+l)!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot a(\omega \otimes \eta)(\sigma(v^k, v^l)) \\
&= a \frac{(k+l)!}{k!l!} \cdot \frac{1}{(k+l)!} \sum_{\sigma \in S_k} \text{sgn } \sigma \cdot (\omega \otimes \eta)(\sigma(v^k, v^l)) \\
&= a \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta)(v^k, v^l) \\
&= a(\omega \wedge \eta)(v^k, v^l) \quad \square
\end{aligned}$$

We can also take advantage of the alternating nature of these tensors to prove additional properties.

Proposition 4.8

Let $\omega \in A^k(V)$ and $\eta \in A^l(V)$. Let $f : V \rightarrow V$ be linear. Then

1. $\omega \wedge \eta = (-1)^{kl}(\eta \wedge \omega)$
2. $f^*(\omega \wedge \eta) = f^*(\omega) \wedge f^*(\eta)$.

Proof. 1. Let $v = (v_1, \dots, v_{k+l}) \in V^{k+l}$. Let $\sigma^* \in S_{k+l}$ be a permutation which sends $\{1, 2, \dots, k+l\}$ to $\{k+l, 1, 2, \dots, k+l-1\}$. Note that this can be achieved using

$k+l-1$ permutations (as $(k+l, k+l-1) \cdot (k+l, k+l-2) \cdot \dots \cdot (k+l, 1)$). Then $(\sigma^*)^l$ is the permutation which takes $\{1, 2, \dots, k+l\}$ to $\{k+1, \dots, k+l, 1, \dots, k\}$, and

$$\operatorname{sgn}(\sigma^*)^l = (\operatorname{sgn}(\sigma^*))^l = ((-1)^{k+l-1})^l = (-1)^{kl+l^2-l} = (-1)^{kl}$$

Then we have

$$\begin{aligned} (\omega \wedge \eta)(v) &= \frac{(k+l)!}{k!l!} \operatorname{Alt}(\omega \otimes \eta)(v) \\ &= \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \operatorname{sgn} \sigma \cdot (\omega \otimes \eta)(v_{\sigma(1)}, \dots, v_{\sigma(k)}, v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)}) \\ &= (-1)^{kl} \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma \cdot (\sigma^*)^l} \operatorname{sgn}(\sigma) \cdot (\omega \otimes \eta)(v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)}, v_{\sigma(1)}, \dots, v_{\sigma(k)}) \\ &= (-1)^{kl} \frac{(k+l)!}{k!l!} \frac{1}{(k+l)!} \sum_{\sigma} \operatorname{sgn}(\sigma) \cdot (\eta \otimes \omega)(v_{\sigma(1)}, \dots, v_{\sigma(k)}, v_{\sigma(k+1)}, \dots, v_{\sigma(k+l)}) \\ &= (-1)^{kl} \frac{(k+l)!}{k!l!} \operatorname{Alt}(\eta \otimes \omega)(v) \\ &= (-1)^{kl} (\eta \wedge \omega)(v) \end{aligned}$$

2. This follows from Proposition 4.3:

$$\begin{aligned} f^*(\omega \wedge \eta) &= \frac{(k+l)!}{k!l!} \operatorname{Alt}(f^*(\omega \otimes \eta)) \\ &= \frac{(k+l)!}{k!l!} \operatorname{Alt}(f^*\omega \wedge f^*\eta) \\ &= f^*\omega \wedge f^*\eta \end{aligned} \quad \square$$

We can also prove associativity of the wedge product:

Theorem 4.9

1. Let $S \in \mathfrak{J}^k(V)$, $T \in \mathfrak{J}^l(V)$, and suppose $\operatorname{Alt}(S) = 0$. Then

$$\operatorname{Alt}(S \otimes T) = \operatorname{Alt}(T \otimes S) = 0$$

2. $\operatorname{Alt}(\operatorname{Alt}(\omega \otimes \eta) \otimes \theta) = \operatorname{Alt}(\omega \otimes \eta \otimes \theta) = \operatorname{Alt}(\omega \otimes \operatorname{Alt}(\eta \otimes \theta))$

3. If $\omega \in A^k(V)$, $\eta \in A^l(V)$, and $\theta \in A^m(V)$, then

$$(\omega \wedge \eta) \wedge \theta = \omega \wedge (\eta \wedge \theta) = \frac{(k+l+m)!}{k!l!m!} \operatorname{Alt}(\omega \otimes \eta \otimes \theta)$$

Proof. 1. Let $G < S_{k+l}$ be the set of all permutations which fix the k -th through $k+l$ -th elements. This is a subgroup of S_{k+l} , so we may consider the set of right cosets $G\sigma'$

for $\sigma' \in S_{k+l}$. Let $v^k = (v_1, \dots, v_k) \in V^k$, $v^l = (v_{k+1}, \dots, v_{k+l})$, and let $\sigma(v) = (v_{\sigma(1)}, \dots, v_{\sigma(k+l)})$. Then

$$\begin{aligned} \text{Alt}(S \otimes T)(v^k, v^l) &= \frac{1}{(k+l)!} \sum_{\sigma \in S_{k+l}} \text{sgn } \sigma \cdot (S \otimes T)(\sigma(v^k, v^l)) \\ &= \frac{1}{(k+l)!} \sum_{G\sigma'} \sum_{\sigma \in G} \text{sgn}(\sigma\sigma') \cdot S(\sigma\sigma'(v^k)) \cdot T(\sigma\sigma'(v^l)) \\ &= \frac{1}{(k+l)!} \sum_{G\sigma'} \text{sgn}(\sigma') T(\sigma'(v^l)) \sum_{\sigma \in G} \text{sgn}(\sigma) \cdot S(\sigma\sigma'(v^k)) \end{aligned}$$

Write $\sigma'(v^k) = w^k$. Noting that $G \cong S_k$, we have

$$\sum_{\sigma \in G} \text{sgn}(\sigma) \cdot S(\sigma\sigma'(v^k)) = \sum_{\sigma \in S_k} \text{sgn}(\sigma) \cdot S(\sigma(w^k)) = k! \text{Alt}(S)(w^k) = 0$$

So $\text{Alt}(S \otimes T) = 0$ and similarly $\text{Alt}(T \otimes S) = 0$.

2. Noting that Alt is linear, we know that

$$\text{Alt}(\text{Alt}(\omega \otimes \eta) - \omega \otimes \eta) = \text{Alt}(\text{Alt}(\omega \otimes \eta)) - \text{Alt}(\omega \otimes \eta) = \text{Alt}(\omega \otimes \eta) - \text{Alt}(\omega \otimes \eta) = 0$$

Applying part 1),

$$0 = \text{Alt}((\text{Alt}(\omega \otimes \eta) - \omega \otimes \eta) \otimes \theta) = \text{Alt}(\text{Alt}(\omega \otimes \eta) \otimes \theta) - \text{Alt}(\omega \otimes \eta \otimes \theta)$$

so

$$(\text{Alt}(\omega \otimes \eta) \otimes \theta) = \text{Alt}(\omega \otimes \theta \otimes \eta)$$

and the other equality is similar.

3. We have

$$\begin{aligned} (\omega \wedge \eta) \wedge \theta &= \frac{(k+l+m)!}{(k+l)!m!} \text{Alt}((\omega \wedge \eta) \otimes \theta) \\ &= \frac{(k+l+m)!}{(k+l)!m!} \text{Alt}\left(\frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \eta) \otimes \theta\right) \\ &= \frac{(k+l+m)!}{k!l!m!} \text{Alt}(\text{Alt}(\omega \otimes \eta) \otimes \theta) \\ &= \frac{(k+l+m)!}{k!l!m!} \text{Alt}(\omega \otimes \eta \otimes \theta) \quad \square \end{aligned}$$

As a result, we will also drop the parentheses when discussing wedge products.

Theorem 4.10

Let V be a real vector space with basis v_1, \dots, v_n , and let $\varphi_1, \dots, \varphi_n$ be the induced dual basis. Then the collection of all k -tensors of the form

$$\varphi_{i_1} \wedge \dots \wedge \varphi_{i_k}$$

where $1 \leq i_1 < \dots < i_k \leq n$, is a basis for $A^k(V)$.

Proof. Let $\omega \in A^k(V)$. Then $\omega \in \mathfrak{J}^k(V)$. By Theorem 4.2, the collection of $\varphi_{j_1} \otimes \dots \otimes \varphi_{j_k}$ is a basis for $\mathfrak{J}^k(V)$ and we have

$$\omega = \sum a_{j_1, \dots, j_k} \varphi_{j_1} \otimes \dots \otimes \varphi_{j_k}$$

Since ω is alternating, we have

$$\omega = \text{Alt}(\omega) = \sum a_{j_1, \dots, j_k} \text{Alt}(\varphi_{j_1} \otimes \dots \otimes \varphi_{j_k}) = \sum a_{j_1, \dots, j_k} \frac{(nk)!}{(k!)^n} \varphi_{j_1} \wedge \dots \wedge \varphi_{j_k}$$

Let j'_1, \dots, j'_k be a reordering of j_1, \dots, j_k such that $j'_1 \leq \dots \leq j'_k$. This may be accomplished by a series of transpositions, each of which changes only the sign of the wedge product by Proposition 4.8. Moreover, if any two of the $\varphi_{j'_i}$ are equal, then the entire wedge product is zero. So we may assume $j'_1 < \dots < j'_k$, and we have

$$\sum a_{j_1, \dots, j_k} \frac{(nk)!}{(k!)^n} (-1)^{M_{j_1, \dots, j_k}} \varphi_{j'_1} \wedge \dots \wedge \varphi_{j'_k}$$

So the $\varphi_{j'_1} \wedge \dots \wedge \varphi_{j'_k}$ span $A^k(V)$.

To show linear independence, suppose we have some linear combination

$$\omega = \sum a_{j_1, \dots, j_k} \varphi_{j_1} \wedge \dots \wedge \varphi_{j_k}$$

□

Appendix A

Solutions to Selected Exercises

A.1 Chapter 1 Exercises

Exercise 1-1 Prove that $|x| \leq \sum_{i=1}^n |x_i|$ for any $x \in \mathbb{R}^n$.

Proof. Let $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ be arbitrary. For each $1 \leq i \leq n$, let us denote by (x_i) the vector $[0 \ \dots \ x_i \ \dots \ 0]^T$, with the x_i term in the i th coordinate. Then for each i , we have the following:

$$|(x_i)| = \sqrt{(x_i)^2} = |x_i|$$

Moreover, by construction we have $x = (x_1) + \dots + (x_n)$. By repeated application of the triangle inequality, we have

$$|x| = \left| \sum_{i=1}^n (x_i) \right| \leq \sum_{i=1}^n |(x_i)| = \sum_{i=1}^n |x_i| \quad \square$$

Exercise 1-2 When does equality hold for the triangle inequality?

I claim that $|x + y| = |x| + |y|$ if and only if $y = \lambda x$ for some $\lambda \geq 0$, or $x = \mathbf{0}$.

Proof. $x = \mathbf{0}$ clearly satisfies the triangle inequality, so assume $x \neq \mathbf{0}$. Following the proof of the triangle inequality given by Spivak, we already see that x, y being linearly dependent

is certainly a necessary condition. Thus, assume that $y = \lambda x$ for some $\lambda \in \mathbb{R}$. Then

$$\begin{aligned} |x + y|^2 &= \sum_{i=1}^n (x_i + y_i)^2 \\ &= \sum_{i=1}^n (x_i + \lambda x_i)^2 \\ &= \sum_{i=1}^n x_i^2 + \lambda^2 \sum_{i=1}^n x_i^2 + 2\lambda \sum_{i=1}^n x_i^2 \\ &= |x|^2 + |\lambda x|^2 + 2\lambda |x|^2 \end{aligned}$$

When $\lambda \geq 0$ we have

$$|x|^2 + |\lambda x|^2 + 2\lambda |x|^2 = |x|^2 + 2|x||\lambda x| + |\lambda x|^2 = (|x| + |\lambda x|)^2 = (|x| + |y|)^2$$

where equality follows by taking the square root on both sides.

When $\lambda < 0$ this becomes

$$|x|^2 + |\lambda x|^2 + 2\lambda |x|^2 = |x|^2 - 2|x||\lambda x| + |\lambda x|^2 = (|x| - |\lambda x|)^2 = (|x| - |y|)^2$$

By taking square roots on both sides, we have $|x + y| = |x| - |y| \neq |x| + |y|$ where the inequality holds since $x \neq \mathbf{0}$, $\lambda \neq 0$ means that $|y| \neq 0$. Thus $y = \lambda x$ for $\lambda \geq 0$, or $x = \mathbf{0}$ is a necessary and sufficient condition. \square

Exercise 1-3 Prove that $|x - y| \leq |x| + |y|$ for any $x, y \in \mathbb{R}^n$.

Proof. Let $x, y \in \mathbb{R}^n$ be arbitrary. Then

$$|x - y| = |x + (-1 * y)| \leq |x| + |(-1) * y| = |x| + |-1||y| = |x| + |y| \quad \square$$

Exercise 1-4 Prove that $||x| - |y|| \leq |x - y|$.

Proof. We expand:

$$\begin{aligned} |x - y|^2 &= \sum_{i=1}^n (x_i - y_i)^2 \\ &= \sum_{i=1}^n x_i^2 + \sum_{i=1}^n y_i^2 - 2 \sum_{i=1}^n x_i y_i \\ &\geq |x|^2 + |y|^2 - 2|x||y| \text{ (by Cauchy-Schwarz)} \\ &= (|x| - |y|)^2 \end{aligned}$$

Taking square roots on both sides (using the fact that it is order-preserving), we get

$$|x - y| \geq ||x| - |y|| \quad \square$$

Exercise 1-5 The quantity $|y - x|$ is called the **distance** between x and y . Prove and interpret geometrically the inequality $|z - x| \leq |z - y| + |y - x|$.

Proof. Noting that $|z - x| = |(z - y) + (y - x)|$, this is a simple application of the triangle inequality. This says that the sum of the lengths of any two sides of a triangle must be greater than the length of the third. \square

Exercise 1-6 Let f, g be integrable on $[a, b]$.

- (a) Prove that $|\int_a^b fg| \leq (\int_a^b f^2)^{\frac{1}{2}} (\int_a^b g^2)^{\frac{1}{2}}$.
- (b) If equality holds, must it be true that $f = \lambda g$ for some $\lambda \in \mathbb{R}$? What if f, g are required to be continuous?
- (c) Show that the Cauchy-Schwarz inequality is a special case of (a).

- (a) *Proof.* We consider the cases $0 = \int_a^b (f - \lambda g)^2$ for some $\lambda \in \mathbb{R}$, and $0 < \int_a^b (f - \lambda g)^2$ for all λ .

Case 1: Here, we have

$$0 = \int_a^b (f - \lambda g)^2 = \int_a^b f^2 - 2\lambda fg + \lambda^2 g^2 = \int_a^b f^2 - 2\lambda \int_a^b fg + \lambda^2 \int_a^b g^2$$

if $\lambda = 0$, then f (and thus fg) is zero on a set of measure 1, immediately making both sides of the inequality 0. Thus assume that $\lambda \neq 0$, which implies

$$\int_a^b fg = \frac{1}{2\lambda} \int_a^b f^2 + \frac{\lambda}{2} \int_a^b g^2$$

so

$$\begin{aligned} \left(\int_a^b fg\right)^2 &= \left(\frac{1}{2\lambda} \int_a^b f^2\right)^2 + \left(\frac{\lambda}{2} \int_a^b g^2\right)^2 + \frac{1}{2} \left(\int_a^b f^2\right) \left(\int_a^b g^2\right) \\ &\leq \frac{1}{2} \left(\int_a^b f^2\right) \left(\int_a^b g^2\right) \\ &\leq \left(\int_a^b f^2\right) \left(\int_a^b g^2\right) \end{aligned}$$

Taking the square root on both sides gives $|\int_a^b fg| \leq (\int_a^b f^2)^{\frac{1}{2}} (\int_a^b g^2)^{\frac{1}{2}}$, as desired.

Case 2: Here, we have

$$\int_a^b (f - g)^2 > 0 \implies \int_a^b f^2 + \int_a^b g^2 > 2 \int_a^b fg$$

Squaring both sides,

$$\left(\int_a^b fg\right)^2 < \left(\frac{1}{2}\int_a^b f^2\right) + \left(\frac{1}{2}\int_a^b g^2\right) + \frac{1}{2}\left(\int_a^b f^2\right)\left(\int_a^b g^2\right)$$

and the rest of the proof is identical to the first case. \square

- (b) *Proof.* Examining the proof of part (a), we must have $0 = \int_a^b (f - \lambda g)^2$ for equality to hold. This implies $f - \lambda g$ is 0 almost everywhere, so $f = \lambda g$ almost everywhere. However, it may not be the case that $f = \lambda g$ everywhere (consider $f = 0$ and $g = 0$ except at countably many points). When f, g are required to be continuous, then they cannot differ on a set of measure zero, so equality implies $f = \lambda g$ for some $\lambda \in \mathbb{R}$. \square
- (c) *Proof.* Let $x, y \in \mathbb{R}^n$ be arbitrary. Define $f : [0, n) \rightarrow \mathbb{R}$ such that $f = x_i$ on the interval $[i - 1, i)$ and define g similarly for y . Then

$$\int_0^n f^2 = \sum_{i=1}^n x_i^2 = |x|^2, \int_0^n g^2 = \sum_{i=1}^n y_i^2 = |y|^2, \int_0^n fg = \sum_{i=1}^n x_i y_i$$

Then by part a,

$$\left|\sum_{i=1}^n x_i y_i\right| = \left|\int_0^n fg\right| \leq \left(\int_a^b f^2\right)^{\frac{1}{2}} \left(\int_a^b g^2\right)^{\frac{1}{2}} = |x||y| \quad \square$$

Exercise 1-7 A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is **norm preserving** if $|T(x)| = |x|$ for all $x \in \mathbb{R}^n$, and **inner product preserving** if $\langle Tx, Ty \rangle = \langle x, y \rangle$ for all $x, y \in \mathbb{R}^n$.

- (a) Prove that T is norm preserving if and only if T is inner product preserving.
- (b) Prove that such a linear transformation T is one-to-one and T^{-1} is of the same sort.

- (a) *Proof.* (\implies) Suppose T is norm preserving. Then for any $x, y \in \mathbb{R}^n$, we use bilinearity of the inner product:

$$\begin{aligned} \langle Tx, Ty \rangle &= \langle Tx - Ty + Ty, Ty \rangle \\ &= \langle Tx - Ty, Ty \rangle + \langle Ty, Ty \rangle \\ &= \langle Tx - Ty, Ty - Tx + Tx \rangle + |Ty| \\ &= \langle Tx - Ty, Ty - Tx \rangle + \langle Tx - Ty, Tx \rangle + |Ty| \\ &= |Tx| - \langle Ty, Tx \rangle + |Ty| - |Tx - Ty| \end{aligned}$$

which gives

$$\begin{aligned} \langle Tx, Ty \rangle &= \frac{1}{2}(|Tx| + |Ty| - |T(x - y)|) && \text{(by linearity of } T) \\ &= \frac{1}{2}(|x| + |y| - |x - y|) && \text{(by norm preserving)} \\ &= \langle x, y \rangle \end{aligned}$$

where the last line follows through a similar calculation as the first part.

(\Leftarrow) Suppose T is inner product preserving. Then for any $x \in \mathbb{R}^n$,

$$|Tx| = \langle Tx, Tx \rangle = \langle x, x \rangle = |x|$$

where the second equality follows since T preserves inner products. \square

- (b) *Proof.* Suppose T is inner product/norm preserving. Suppose $Tx = Ty$. Since T is linear, we have $T(x - y) = 0$. So $|T(x - y)| = 0$. But T is norm preserving, so $|x - y| = 0$, which occurs only when $x - y = 0$, showing that $x = y$. So T is one-to-one.

Let T^{-1} denote the inverse of T (which exists since T is an injective endomorphism on finite dimensional vector spaces). Then let $x \in \mathbb{R}^n$ be arbitrary. Since T is norm preserving, we have

$$|T^{-1}x| = |TT^{-1}x| = |x|$$

so T^{-1} is norm preserving as well. \square

Exercise 1-8 If $x, y \in \mathbb{R}^n$ are nonzero, then the angle between x and y is denoted $\angle(x, y)$, which is defined as $\arccos\left(\frac{\langle x, y \rangle}{|x| \cdot |y|}\right)$. This is well-defined since $\left|\frac{\langle x, y \rangle}{|x| \cdot |y|}\right| \leq 1$ by Cauchy-Schwarz. The linear transformation T is angle preserving if T is one-to-one and for any $x, y \neq 0$ we have $\angle(Tx, Ty) = \angle(x, y)$.

- (a) Prove that if T is norm preserving, then T is angle preserving.
- (b) If there is a basis x_1, \dots, x_n of \mathbb{R}^n and numbers $\lambda_1, \dots, \lambda_n$ such that $Tx_i = \lambda_i x_i$, prove that T is angle preserving only if all $|\lambda_i|$ are equal. (**Note:** Spivak's original exercise has an if and only if here, but this is false.)
- (c) What are all angle preserving $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$?

- (a) *Proof.* Since T is both norm preserving and inner product preserving by Exercise 1-7, we have

$$\frac{\langle Tx, Ty \rangle}{|Tx| \cdot |Ty|} = \frac{\langle x, y \rangle}{|x| \cdot |y|}$$

so

$$\angle(Tx, Ty) = \arccos\left(\frac{\langle Tx, Ty \rangle}{|Tx| \cdot |Ty|}\right) = \arccos\left(\frac{\langle x, y \rangle}{|x| \cdot |y|}\right) = \angle(x, y) \quad \square$$

- (b) *Proof.* Proof by contrapositive. Suppose $|\lambda_i| \neq |\lambda_j|$ for some $i \neq j$. Then consider the vectors

$$v_1 = x_i + x_j, v_2 = x_i - \frac{|x_i|}{|x_j|}x_j$$

Since x_i, x_j are linearly independent, neither v_1 or v_2 is the zero vector. Then we have

$$\begin{aligned}\cos \angle(v_1, v_2) &= \cos \arccos \left(\frac{\left\langle x_i + x_j, x_i - \frac{|x_i|}{|x_j|} x_j \right\rangle}{|x_i + x_j| |x_i - \frac{|x_i|}{|x_j|} x_j|} \right) \\ &= \frac{|x_i|^2 - \frac{|x_i|^2}{|x_j|^2} |x_j|^2}{|x_i + x_j| |x_i - \frac{|x_i|}{|x_j|} x_j|} \\ &= 0\end{aligned}$$

On the other hand,

$$\begin{aligned}\cos \angle(T(v_1), T(v_2)) &= \cos \angle(\lambda_i x_i + \lambda_j x_j, \lambda_i x_i - \lambda_j \frac{|x_i|}{|x_j|} x_j) \\ &= \frac{\lambda_i^2 |x_i|^2 - \lambda_j^2 |x_i|^2}{|\lambda_i x_i + \lambda_j x_j| |\lambda_i x_i - \lambda_j \frac{|x_i|}{|x_j|} x_j|} \neq 0\end{aligned}$$

where the last inequality holds since $|\lambda_i| \neq |\lambda_j| \implies \lambda_i^2 \neq \lambda_j^2$. So if $|\lambda_i| \neq |\lambda_j|$, then T is not angle preserving. So T is angle preserving only if $|\lambda_i| = |\lambda_j|$ for all i, j . \square

- (c) Intuitively, the answer is that T must consist of only rotation and scaling by a constant factor. More rigorously, the singular values of T must all be $\sigma_1 = \dots = \sigma_n = k$ for some $k > 0$. We do not provide a full proof here.

Exercise 1-9 If $0 \leq \theta < \pi$, then let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ have the matrix in the standard basis given by

$$\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

Show that T is angle preserving, and that for any $x \neq 0$, $\angle(x, Tx) = \theta$.

Proof. To show that T is one-to-one, we instead prove that T is invertible. Consider the matrix

$$T' = \begin{bmatrix} \cos(-\theta) & \sin(-\theta) \\ -\sin(-\theta) & \cos(-\theta) \end{bmatrix}$$

Then

$$\begin{aligned}TT' &= \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix} \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \\ &= \begin{bmatrix} \cos^2 \theta + \sin^2 \theta & -\cos \theta \sin \theta + \cos \theta \sin \theta \\ -\cos \theta \sin \theta + \cos \theta \sin \theta & \sin^2 \theta + \cos^2 \theta \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}\end{aligned}$$

Since T is square and $TT' = I$, we have $T'T = I$ so T is invertible and thus must be one-to-one.

Let $x, y \neq 0 \in \mathbb{R}^2$ be arbitrary. Suppose $x = (x_1, x_2)$, $y = (y_1, y_2)$. Then

$$\cos \angle(x, y) = \frac{x_1 y_1 + x_2 y_2}{\sqrt{x_1^2 + x_2^2} \sqrt{y_1^2 + y_2^2}}$$

Moreover, $Tx = (x_1 \cos \theta + x_2 \sin \theta, x_2 \cos \theta - x_1 \sin \theta)$ and $Ty = (y_1 \cos \theta + y_2 \sin \theta, y_2 \cos \theta - y_1 \sin \theta)$. Then we have

$$\begin{aligned} \langle Tx, Ty \rangle &= (x_1 \cos \theta + x_2 \sin \theta)(y_1 \cos \theta + y_2 \sin \theta) + (x_2 \cos \theta - x_1 \sin \theta)(y_2 \cos \theta - y_1 \sin \theta) \\ &= x_1 y_1 \underbrace{(\cos^2 \theta + \sin^2 \theta)}_{=1} + x_1 y_2 \underbrace{(\cos \theta \sin \theta - \sin \theta \cos \theta)}_{=0} \\ &\quad + x_2 y_1 \underbrace{(\sin \theta \cos \theta - \sin \theta \cos \theta)}_{=0} + x_2 y_2 \underbrace{(\sin^2 \theta + \cos^2 \theta)}_{=1} \\ &= x_1 y_1 + x_2 y_2 = \langle x, y \rangle \end{aligned}$$

and

$$\begin{aligned} |Tx| &= \sqrt{(x_1 \cos \theta + x_2 \sin \theta)^2 + (x_2 \cos \theta - x_1 \sin \theta)^2} \\ &= \sqrt{x_1^2 (\cos^2 \theta + \sin^2 \theta) + x_2^2 (\sin^2 \theta + \cos^2 \theta)} \\ &= \sqrt{x_1^2 + x_2^2} \\ &= |x| \end{aligned}$$

Similarly,

$$|Ty| = |y|$$

Then

$$\begin{aligned} \angle(Tx, Ty) &= \arccos \left(\frac{\langle Tx, Ty \rangle}{|Tx| |Ty|} \right) \\ &= \arccos \left(\frac{\langle x, y \rangle}{|x| |y|} \right) \\ &= \angle(x, y) \end{aligned}$$

Lastly, using the fact that $|x| = |Tx|$,

$$\begin{aligned} \angle(x, Tx) &= \arccos \left(\frac{\langle x, Tx \rangle}{|x| |Tx|} \right) \\ &= \arccos \left(\frac{x_1^2 \cos \theta + x_1 x_2 \sin \theta + x_2^2 \cos \theta - x_1 x_2 \sin \theta}{|x|^2} \right) \\ &= \arccos \left(\cos \theta \frac{x_1^2 + x_2^2}{|x|^2} \right) \\ &= \arccos \cos \theta \\ &= \theta \end{aligned} \quad \square$$

Exercise 1-10 If $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is a linear transformation, show that there is a number M such that $|T(h)| \leq M|h|$ for $h \in \mathbb{R}^m$.

Proof. By singular value decomposition, there are orthonormal bases $\mathcal{B} = \{u_1, \dots, u_m\} \subseteq \mathbb{R}^m$ and $\mathcal{C} = \{v_1, \dots, v_n\} \subseteq \mathbb{R}^n$ as well as scalars $\sigma_1 \geq \dots \geq \sigma_m \geq 0$ such that $Tu_i = \sigma_i v_i$ for all i (with $Tu_j = 0$ for any $j \geq n$). Then for any $h \in \mathbb{R}^m$, if we suppose that $h = a_1 u_1 + \dots + a_m u_m$, then we have

$$\begin{aligned} |Th| &= |T(a_1 u_1 + \dots + a_m u_m)| \\ &= |a_1 T u_1 + \dots + a_m T u_m| \\ &= |a_1 \sigma_1 v_1 + \dots + a_m \sigma_m v_m| \end{aligned}$$

(where the indices only run to n if $n < m$). Now since \mathcal{C} is orthonormal, the Pythagorean identity gives

$$|a_1 \sigma_1 v_1 + \dots + a_m \sigma_m v_m|^2 = a_1^2 \sigma_1^2 + \dots + a_m^2 \sigma_m^2 \leq (\sigma_m)^2 (a_1^2 + \dots + a_m^2)$$

But since \mathcal{B} is also orthonormal, we have $(a_1^2 + \dots + a_m^2) = |h|^2$. So

$$|Th|^2 \leq \sigma_m^2 |h|^2 \implies |Th| \leq \sigma_m |h|$$

so our choice of $M = \sigma_m$ works. □

Exercise 1-11 If $x, y \in \mathbb{R}^n$ and $z, w \in \mathbb{R}^m$, show that $\langle (x, z), (y, w) \rangle = \langle x, y \rangle + \langle z, w \rangle$, and that $|(x, z)| = \sqrt{|x|^2 + |z|^2}$. Recall that $(x, z) \in \mathbb{R}^{n+m}$ is the concatenation of x and z .

Proof. For the first statement,

$$\begin{aligned} \langle (x, z), (y, w) \rangle &= \sum_{i=1}^{n+m} (x, z)_i (y, w)_i \\ &= \sum_{i=1}^n (x, z)_i (y, w)_i + \sum_{j=1}^m (x, z)_{n+j} (y, w)_{n+j} \\ &= \sum_{i=1}^n x_i y_i + \sum_{j=1}^m z_j w_j \\ &= \langle x, y \rangle + \langle z, w \rangle \end{aligned}$$

For the second statement,

$$|(x, z)|^2 = \langle (x, z), (x, z) \rangle = \langle x, x \rangle + \langle z, z \rangle = |x|^2 + |z|^2$$

where the second equality is by the first statement. Taking square roots on both sides recovers $|(x, z)| = \sqrt{|x|^2 + |z|^2}$. □

Exercise 1-12 Let $(\mathbb{R}^n)^*$ denote the dual space of \mathbb{R}^n , which is the space of all linear functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$. If $x \in \mathbb{R}^n$, then define $\phi_x \in (\mathbb{R}^n)^*$ such that $\phi_x(y) := \langle x, y \rangle$. Define $T : \mathbb{R}^n \rightarrow (\mathbb{R}^n)^*$ such that $T(x) = \phi_x$. Show that T is one-to-one and conclude that each $\phi \in (\mathbb{R}^n)^*$ is ϕ_x for a unique $x \in \mathbb{R}^n$.

Proof. Suppose $\phi_x = \phi_y$. Then $\langle x, z \rangle = \langle y, z \rangle$ for all $z \in \mathbb{R}^n$. Choosing $z = x - y$, this gives

$$0 = \langle x, z \rangle - \langle y, z \rangle = \langle x - y, z \rangle = \langle x - y, x - y \rangle = |x - y|^2$$

which implies that $|x - y|^2 = 0$. So $x = y$. The rest of the proof follows since $\dim \mathbb{R}^n = \dim(\mathbb{R}^n)^*$, so T is injective between vector spaces of the same dimension and is thus surjective and bijective. \square

Exercise 1-13 (Pythagorean Identity) If $x, y \in \mathbb{R}^n$, then x and y are called orthogonal if $\langle x, y \rangle = 0$. If x and y are orthogonal, prove that $|x + y|^2 = |x|^2 + |y|^2$.

Proof. By the definition of the norm and bilinearity of the inner product,

$$\begin{aligned} |x + y|^2 &= \langle x + y, x + y \rangle \\ &= \langle x, x \rangle + \langle y, y \rangle + 2 \underbrace{\langle x, y \rangle}_{=0} \\ &= |x|^2 + |y|^2 \end{aligned} \quad \square$$

Exercise 1-14 Prove that the arbitrary union of open sets is open. Prove that the finite intersection of open sets is open. Show that an infinite union of open sets need not be open.

Proof. Let $U = \bigcup_{i \in I} U_i$ be the union of some open sets over an arbitrary indexing set I . Then for any $x \in U$, $x \in U_i$ for some i . Then $x \in B \subseteq U_i$ for some open rectangle B . Since $B \subseteq U_i$, $B \subseteq U$, so $x \in B \subseteq U$. So U is open.

Let $U = U_1 \cap U_2$ for some open sets U_1, U_2 . Let $x \in U$ be arbitrary. Then $x \in B_{r_1}(x) \subseteq U_1$ and $x \in B_{r_2}(x) \subseteq U_2$ for some radii r_1, r_2 . Taking $r = \min\{r_1, r_2\} > 0$, we have $x \in B_r(x) \subseteq B_{r_1} \subseteq U_1$ and $B_r(x) \subseteq B_{r_2} \subseteq U_2$, so $x \in B_r(x) \subseteq U$. By induction, this extends to any finite intersection.

The intersection of the sets $(-1/n, 1/n)$ for $n \in \mathbb{N}$ is the singleton $\{0\}$, which is not open. \square

Exercise 1-15 Prove that the open ball $B_r(a) := \{x \in \mathbb{R}^n : |x - a| < r\}$ is indeed open.

Proof. When $r = 0$, $B_r(a) = \emptyset$ which is vacuously open. If $r > 0$, then pick some $x \in B_r(a)$. Let $r' = r - |x - a|$. Then if $x = (x_1, \dots, x_n)$, consider the box B with sides

$(x_1 - r'/n, x_1 + r'/n) \times \dots \times (x_n - r'/n, x_n + r'/n)$. For any other $y \in B$, we have $|x_i - y_i| \leq r'/n$ by construction, so

$$|x - y| \leq |x_1 - y_1| + \dots + |x_n - y_n| \leq r'$$

By the triangle inequality,

$$|y - a| = |y - x - (a - x)| \leq |y - x| + |a - x| \leq r' + |a - x| = r - |x - a| + |x - a| = r$$

So $y \in B_r(a)$, and thus $B \subseteq B_r(a)$. So $B_r(a)$ is open. \square

Exercise 1-16 Find the interior, exterior, and boundary of the following sets:

1. $A := \{x \in \mathbb{R}^n : |x| \leq 1\}$
2. $B := \{x \in \mathbb{R}^n : |x| = 1\}$
3. $C := \{x \in \mathbb{R}^n : \text{each coordinate } x_i \in \mathbb{Q}\}$

1. We proved in Exercise 1-15 that $B_1(\mathbf{0}) \subseteq A$ is open. So $B_1(\mathbf{0}) \subseteq \text{int } A$.

I claim that $\mathbb{R}^n \setminus A = \text{ext } A$. Let $x \in \mathbb{R}^n \setminus A$. Then take the open ball $B_{|x|-1}(x)$. For any $y \in B_{|x|-1}(x)$, the reverse triangle inequality tells us

$$|y| \geq ||y - x| - |x||$$

Since $y \in B_{|x|-1}(x)$, $|y - x| \leq |x| - 1$. So $|y - x| - |x| \leq -1$, and thus

$$|y| \geq ||y - x| - |x|| \geq 1$$

so $y \in \mathbb{R}^n \setminus A$. Thus $\mathbb{R}^n \setminus A \subseteq \text{ext } A$, but $\text{ext } A \subseteq \mathbb{R}^n \setminus A$ (this is easy to see based on the definition of $\text{ext } A$), so $\mathbb{R}^n \setminus A = \text{ext } A$.

Lastly, for any x with $|x| = 1$, pick any open ball $B_r(x)$. Then the point $y = x + \frac{r}{2}x$ has

$$|y - x| = \left| \frac{r}{2}x \right| = \frac{r}{2} \underbrace{|x|}_{=1} < r$$

So $y \in B_r(x)$. Moreover,

$$|y| = \left(1 + \frac{r}{2}\right) \underbrace{|x|}_{=1} > 1$$

so $y \in \mathbb{R}^n \setminus A$. On the other hand, a similar calculation shows that $z = x - \frac{r}{2}x \in B_r(x)$ is in A . So the set of points with $|x| = 1$ is a subset of ∂A . But $\text{int } A \sqcup \partial A \sqcup \text{ext } A = \mathbb{R}^n$, and we have already partitioned \mathbb{R}^n , so our subsets must be equalities and we must have $\text{int } A = \{x : |x| < 1\}$, $\partial A = \{x : |x| = 1\}$, $\text{ext } A = \{x : |x| > 1\}$.

2. By the same argument as before, the set of $|x| > 1$ is a subset of $\text{ext } B$. By a similar argument, the set of $|x| < 1$ is also a subset of $\text{ext } B$. Lastly, the same argument shows that B itself is not a subset of $\text{int } B$. But B cannot be in $\text{ext } B$, so we must have $\text{int } B = \emptyset$, $\partial B = \{x : |x| = 1\}$, $\text{ext } B = \{x : |x| \neq 1\}$.

3. Let $x \in \mathbb{R}^n$ be arbitrary. Then let $D = (y_1, z_1) \times \dots \times (y_n, z_n)$ be an arbitrary open rectangle containing x . By the density of \mathbb{Q} in \mathbb{R} , we can pick rational numbers $q_i \in (y_i, z_i)$. Then the point $q = (q_1, \dots, q_n) \in C$ and $q \in D$, so D contains points of C . Similarly, we can construct a point with all irrational coordinates $p = (p_1, \dots, p_n) \notin C$ and $p \in D$, so D contains points of $\mathbb{R}^n \setminus C$. Thus $x \in \partial C$. x was arbitrary, so $\partial C = \mathbb{R}^n$ and $\text{int } C = \text{ext } C = \emptyset$.

Exercise 1-17 Construct a set $A \subseteq [0, 1] \times [0, 1]$ such that A contains at most one point on each horizontal and each vertical line but has $\text{ext } A = [0, 1] \times [0, 1]$.

We construct sets recursively as follows: for A_1 , pick a point in each quadrant of $[0, 1] \times [0, 1]$, such that none lie on the same horizontal or vertical line. For A_2 , pick a point in each sixteenth of $[0, 1] \times [0, 1]$ such that none lie on the same horizontal or vertical line, and none lie on the same horizontal or vertical line as the points in A_1 . Continue doing this, picking 4^i points for A_i such that no point $x \in A_i$ shares a vertical or horizontal line with a point $y \in \bigcup_{k=1}^i A_k$. This is possible because each choice of point removes only a single vertical line and horizontal line from our possible choices, which is a set of measure zero, so we always have a set of measure one to choose from. Then take our set to be $A = \bigcup_{i=1}^{\infty} A_i$. By construction, this set satisfies the vertical/horizontal line property. This set has no interior, since a nontrivial open rectangle being a subset of A would violate the vertical/horizontal line condition. Moreover, for any point $x \in [0, 1] \times [0, 1]$ and any radius r , we simply look in a (4^i) -ant of length $r/2$ or less in order to find a point y that is close to x . So $\partial A = [0, 1] \times [0, 1]$.

Exercise 1-18 If $A \subseteq [0, 1]$ is the union of open intervals (a_i, b_i) such that any rational number in $(0, 1)$ is in (a_i, b_i) for some i , prove that $\partial A = [0, 1] \setminus A$.

Proof. Since A is the union of open intervals, A is open and thus $\text{int } A = A$. I claim that $\text{ext } A = \mathbb{R}^n \setminus [0, 1]$. Clearly $\mathbb{R}^n \setminus [0, 1] \subseteq \text{ext } A$. Then take some point $x \in [0, 1]$. For any open interval (a, b) containing x , the density of \mathbb{Q} tells us that there is a rational number in $(a, b) \cap [0, 1]$, so $x \notin \text{ext } A$. So $\text{ext } A = \mathbb{R}^n \setminus [0, 1]$, $\text{int } A = A$, and this forces $\partial A = [0, 1] \setminus A$. \square

Exercise 1-19 If A is a closed set that contains every rational $r \in [0, 1]$, show that $[0, 1] \subseteq A$.

Proof. Suppose not. Then there is some $x \in [0, 1]$ with $x \in \mathbb{R}^n \setminus A$. x must be in $(0, 1)$, which is open. Moreover, $x \in \mathbb{R}^n \setminus A$, which is open since A is closed, so $x \in (\mathbb{R}^n \setminus A) \cap (0, 1)$ which is open (since the finite intersection of open sets is open). Take some open interval $(a, b) \subseteq (\mathbb{R}^n \setminus A) \cap (0, 1)$. By the density of \mathbb{Q} , there is a rational r in (a, b) . But $r \in A$ by definition, so $(a, b) \not\subseteq \mathbb{R}^n \setminus A$, so $\mathbb{R}^n \setminus A$ isn't open, which contradicts the assumption that A is closed. So we must have $[0, 1] \subseteq A$. \square

Exercise 1-20 Prove that a compact subset of \mathbb{R}^n is closed and bounded.

Proof. Suppose $K \subseteq \mathbb{R}^n$ is compact. The collection of open rectangles $(i-1, i+1) \times (j-1, j+1) \dots \times (k-1, k+1)$ for $i, j, \dots, k \in \mathbb{Z}$ covers \mathbb{R}^n , so it covers K . Then a finite number of these boxes covers K , so it is bounded.

We wish to show that $\mathbb{R}^n \setminus K$ is open. Suppose it is not. Then there is some $x \in \mathbb{R}^n \setminus K$ such that for all open balls $B_r(x)$, $B_r(x) \cap K \neq \emptyset$. We can construct a sequence of points $y_1, y_2, \dots \in K$ as follows: Pick some r_1 , say $r_1 = 1$. Then $B_{r_1}(x)$ contains some point $y_1 \in K$. Let $r_2 = |y_1 - x|$ (note this is strictly less than r_1 since $y_1 \in B_{r_1}(x) \implies |y_1 - x| < r_1$). Next, $B_{r_2}(x)$ contains some other point $y_2 \in K$, and $|y_2 - x| < r_2 = |y_1 - x|$. Continue this to construct a sequence of points $y_1, y_2, \dots \in K$ such that $|y_1 - x| > |y_2 - x| > \dots$

We use this sequence to create an open cover of K . Let $r_i = |y_i - x|$. Let C be the closed ball with radius r_2 and center x . The set $\mathbb{R}^n \setminus C$ is open, since its complement C is closed. Now let $R_i := \{y : r_{i+2} < |y-x| < r_i\}$ be the open ring with outer radius r_i and inner radius r_{i+2} . Then $\bigcup R_i = \{y : |y-x| < r_1\} = B_{r_1}(x)$ contains all points with distance $|y-x| < r_1$. $\mathbb{R}^n \setminus C$ contains all points with distance $|y-x| > r_2$. But $r_2 < r_1$, so $\mathbb{R}^n \setminus C \cup \bigcup R_i = \mathbb{R}^n$.

Thus the collection $\mathcal{O} = \{\mathbb{R}^n \setminus C, R_1, R_2, \dots\}$ covers \mathbb{R}^n and thus K . But if we pick only a finite number of these, then there is some R_i in the finite subcover such that i is maximal in the subcover, so the points y_{i+2}, y_{i+3}, \dots are not contained in the subcover, and thus K is not compact. So if K is compact, then it is closed. \square

Exercise 1-21

1. If A is closed and $x \notin A$ prove that there is a number $d > 0$ such that $|y-x| \geq d$ for all $y \in A$.
2. If A is closed, B is compact, and $A \cap B = \emptyset$, prove that there is $d > 0$ such that $|y-x| \geq d$ for all $y \in A$ and $x \in B$.
3. Give a counterexample in \mathbb{R}^2 if A and B are closed but neither is compact.

1. *Proof.* Since A is closed, $\mathbb{R}^n \setminus A$ is open. Let $x \notin A$. Then $x \in \mathbb{R}^n \setminus A$, so there is an open ball $B_r(x) \subseteq \mathbb{R}^n \setminus A$. Then we have $|x-y| < r \implies y \in B_r(x) \subseteq \mathbb{R}^n \setminus A \implies y \notin A$, and thus for any $y \in A$ we must have $|x-y| \geq r$. \square

2. *Proof.* For each point $b \in B$, part (a) tells us there is a distance r_b such that $|b-y| \geq r_b$ for any $y \in A$. Consider the collection of open balls $(B_{r_b/2}(b))_{b \in B}$. This collection covers B , so we pick a finite subcover $\{B_{r_{b_1}/2}(b_1), B_{r_{b_2}/2}(b_2), \dots, B_{r_{b_n}/2}(b_n)\}$. For any $x \in B$, $x \in B_{r_{b_i}/2}(b_i)$ for some i . Then by the reverse triangle inequality, for any $y \in A$, we have

$$|y-x| = |y-b_i - (x-b_i)| \geq ||y-b_i| - |x-b_i||$$

Since $y \in A$, $|y-b_i| \geq r_{b_i}$. Since $x \in B_{r_{b_i}/2}(b_i)$, $|x-b_i| \leq r_{b_i}/2 \leq r_{b_i} \leq |y-b_i|$. So the quantity $|y-b_i| - |x-b_i|$ is positive, so

$$|y-x| \geq |y-b_i| - |x-b_i| \geq r_{b_i} - \frac{r_{b_i}}{2} = \frac{r_{b_i}}{2} \geq \frac{\min_{1 \leq i \leq n} r_{b_i}}{2}$$

Since $r_{b_i} \geq 0$ for all i and there are finite i , $\min r_{b_i}$ is well defined and positive. Thus for arbitrary $y \in A$, $x \in B$, we have $|y - x| \geq \min r_{b_i}/2 = d > 0$. \square

3. We define two sets as follows: first, pick $A = \mathbb{N}$. Next, pick $B = \{x_1, x_2, \dots\}$, where $x_i = i + \frac{1}{i+1}$. Since x_i is never an integer, $A \cap B = \emptyset$. However, let $r > 0$ be arbitrary. Then pick i large enough that $\frac{1}{i+1} < r$. Choosing $x = x_i$, $y = i$, we have

$$|x - y| = |x_i - i| = \left| \frac{1}{i+1} \right| = \frac{1}{i+1} < r$$

Exercise 1-22 If U is open and $C \subseteq U$ is compact, show that there is a compact set D such that $C \subseteq \text{int } D$ and $D \subseteq U$.

Proof. Since U is open, $\mathbb{R}^n \setminus U$ is closed. Thus by Exercise 1-21 part (b), there is a distance d such that $|y - x| < d$ for any $x \in C$ and $y \in \mathbb{R}^n \setminus U$. Let $B_x = B_d(x)$ be the open ball of radius d and center x . Let $\overline{B}_x = \overline{B}_d(x) = \{y : |y - x| \leq d\}$ be the closed ball of radius d and center x .

The collection $(B_x)_{x \in C}$ is an open cover of C compact, so we pick a finite subcollection B_{x_1}, \dots, B_{x_n} . Then let $D = \overline{B}_{x_1} \cup \dots \cup \overline{B}_{x_n}$. We have $\overline{B}_{x_i} \supseteq B_{x_i}$ for all i , so

$$D = \bigcup_{i=1}^n \overline{B}_{x_i} \supseteq \bigcup_{i=1}^n B_{x_i} \supseteq C$$

so $C \subseteq D$. Moreover, for any point $y \in \mathbb{R}^n \setminus U$ and $x \in D$, $x \in B_{x_i}$ for some i . Then $|x - x_i| \leq d/2$, and $|y - x_i| \geq d$, so

$$|y - x| \geq |y - x_i| - |x - x_i| \geq d - \frac{d}{2} = \frac{d}{2} > 0$$

so $D \cap \mathbb{R}^n \setminus U = \emptyset$ and thus $D \subseteq U$. \square

Exercise 1-23 If $f : A \rightarrow \mathbb{R}^m$ and $a \in A$, show that $\lim_{x \rightarrow a} f(x) = b = (b_1, \dots, b_m)$ if and only if $\lim_{x \rightarrow a} f^i(x) = b_i$ for each i (recall f^i is the i th component function).

Proof. (\implies) Suppose $\lim_{x \rightarrow a} f(x) = b$. Then for any $\varepsilon > 0$, there is $\delta > 0$ such that $|x - a| < \delta$ and $x \in A$ implies $|f(x) - b| < \varepsilon$. Then for any such x , we have $|f^i(x) - b_i|^2 \leq \sum_{j=1}^m |f^j(x) - b_j|^2 = |f(x) - b|^2 < \varepsilon^2$ so $|f^i(x) - b_i| < \varepsilon$. So $\lim_{x \rightarrow a} f^i(x) = b_i$.

Suppose $\lim_{x \rightarrow a} f^i(x) = b_i$ for each i . Then for any $\varepsilon > 0$, pick $\delta_i > 0$ for each i such that $|x - a| < \delta_i \implies |f^i(x) - b_i| < \varepsilon/\sqrt{m}$. Let $\delta = \min \delta_i$. Then for any x with $|x - a| < \delta$,

$$|f(x) - b|^2 = \sum_{i=1}^m |f^i(x) - b_i|^2 < \varepsilon^2/m = \varepsilon^2$$

so $|f(x) - b| < \varepsilon$ and thus $\lim_{x \rightarrow a} f(x) = b$. \square

Exercise 1-24 Prove that $f : A \rightarrow \mathbb{R}^m$ is continuous at a if and only if each f^i is.

Proof. Immediate from Exercise 1-23. \square

Exercise 1-25 Prove that a linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous.

Proof. From Exercise 1-10, we know that there exists $M > 0$ such that $|T(h)| \leq M|h|$ for all h . Then at any point $a \in \mathbb{R}^n$, let $\varepsilon > 0$ be arbitrary. Set $\delta = \varepsilon/M$. Then for any $x \in \mathbb{R}^n$ with $|x - a| < \delta$, we have

$$|T(x) - T(a)| = |T(x - a)| \leq M|x - a| < M\frac{\varepsilon}{M} = \varepsilon \quad \square$$

Exercise 1-26 Let $A = \{(x, y) \in \mathbb{R}^2 : x > 0 \text{ and } 0 < y < x^2\}$.

- (a) Show that every straight line through $(0, 0)$ contains an interval around $(0, 0)$ which is in $\mathbb{R}^2 \setminus A$.
- (b) Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by $f(x) = 0$ if $x \notin A$ and $f(x) = 1$ if $x \in A$. For $h \in \mathbb{R}^2$ define $g_h : \mathbb{R} \rightarrow \mathbb{R}$ by $g_h(t) = f(th)$. Show that each g_h is continuous at 0, but f is not continuous at $(0, 0)$. (This problem shows that f is continuous in any direction, but not continuous as a two-variable function).

- (a) *Proof.* Suppose $y = mx$ defines a straight line through $(0, 0)$. When $m = 0$ one can verify that the entire line is in $\mathbb{R}^2 \setminus A$ since $y = 0$. (For a vertical line we similarly have $x = 0$ so the line is in $\mathbb{R}^2 \setminus A$). Then consider the interval $[-|m|, |m|]$. The portion of the line with $x \leq 0$ is automatically in $\mathbb{R}^2 \setminus A$, but for any $x \in (0, |m|]$,

$$x^2 \leq |m|x = y$$

so the entire interval $[-|m|, |m|]$ is in $\mathbb{R}^2 \setminus A$. \square

- (b) *Proof.* Pick some g_h . By part (a), there is an interval about 0 such that $th \in \mathbb{R}^2 \setminus A$, so $g_h(t) = 0$. So $g_h(t) = 0$ on an interval about 0, so $\lim_{t \rightarrow 0} g_h(t) = 0 = g_h(0)$. Thus each g_h is continuous at 0.

To show f is not continuous at 0, pick $\varepsilon = 1/2$. Let $\delta > 0$ be arbitrary. Assume $\delta < 1$ since this will automatically prove larger δ . Then the point $x = (\delta/2, \delta^2/5)$ is in A , so $f(x) = 1$. Moreover,

$$|x - 0| = |x| \leq \frac{\delta}{2} + \frac{\delta^2}{5} \leq \frac{\delta}{2} + \frac{\delta}{5} < \delta$$

But $|f(x) - f(0)| = |1| = 1 > \varepsilon$, so f is not continuous at 0. \square

Exercise 1-27 Prove that $\{x \in \mathbb{R}^n : |x - a| < r\}$ is open using the topological condition.

Proof. Consider the function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ with $f(x) = |x - a|$. To prove f is continuous, pick some point y . Then let $\varepsilon > 0$ and set $\delta = \varepsilon$. Then we have

$$|x - y| < \delta \implies |f(x) - f(y)| = ||x - a| - |y - a|| \leq |x - a - (y - a)| = |x - y| < \delta = \varepsilon$$

so f is continuous. Thus the preimage of the open ball $B_r(0)$ under f is open, but this is precisely the set $\{x \in \mathbb{R}^n : f(x) = |x - a| < r\}$. \square

Exercise 1-28 Suppose $A \subseteq \mathbb{R}^n$ is not closed. Show that there exists an unbounded continuous function $f : A \rightarrow \mathbb{R}$.

Proof. Let $A \subseteq \mathbb{R}^n$ be not closed. Then $\mathbb{R}^n \setminus A$ is not open, so there exists a point $x \in \mathbb{R}^n \setminus A$ such that every $B_r(x)$ contains a point in A . Then define $f : A \rightarrow \mathbb{R}$ by

$$f(y) = \frac{1}{|y - x|}$$

To verify that this function is continuous, first consider the function $|y - x|$. Letting $a \in A$ be arbitrary, for any $\varepsilon > 0$ set $\delta = \varepsilon$. Then for any $b \in A$ with $|b - a| < \delta$,

$$|f(b) - f(a)| = ||b - x| - |a - x|| \leq |b - a| < \delta = \varepsilon$$

So $y \mapsto |y - x|$ is continuous. Then since $|y - x| \neq 0$ for $y \in A$, and f is the quotient of nonzero continuous functions, f is continuous.

To show that f is unbounded, pick $M > 0$. Then by our choice of x , the ball $B_{1/M}(x)$ contains a point $y \in A$. Then

$$f(y) = \frac{1}{|y - x|} \geq \frac{1}{\frac{1}{M}} = M \quad \square$$

Exercise 1-29 Let $K \subseteq \mathbb{R}^n$ be compact, and let $f : K \rightarrow \mathbb{R}$ be continuous. Show that f attains a maximum and minimum value.

Proof. Since K is compact and f is continuous, $f(K)$ is compact. Specifically, it is bounded, so let $\alpha = \sup f(K)$. We want to show $\alpha \in f(K)$. By way of contradiction, suppose $\alpha \notin f(K)$. Then since $f(K)$ is closed, $\mathbb{R} \setminus f(K)$ is open, so there is an interval $(\alpha - \varepsilon, \alpha + \varepsilon)$ that doesn't intersect $f(K)$. But then $\alpha - \varepsilon$ is also an upper bound for $f(K)$, contradicting that fact that $\alpha = \sup f(K)$. So we must have $\sup f(K) = \max f(K) \in f(K)$, and thus there is a $y \in K$ such that $f(y) = \max f(K)$. The proof for the minimum is similar. \square

Exercise 1-30 Let $f : [a, b] \rightarrow \mathbb{R}$ be increasing. Let $x_1, \dots, x_n \in [a, b]$ be distinct. Show that

$$\sum_{i=1}^n o(f, x_i) \leq f(b) - f(a)$$

Proof. Note that since f is increasing, for any $[c, d] \subseteq [a, b]$, we have

$$\max_{[c,d]} f(x) = f(d), \min_{[c,d]} f(x) = f(c)$$

In particular, $M(f, x, \delta) = f(x+\delta)$ and $m(f, x, \delta) = f(x-\delta)$, so $f(x+\delta) - f(x-\delta) \geq o(f, x)$.

We may suppose that the x_i are ordered, so that $x_1 < \dots < x_n$. Pick δ small enough that $|x_{i+1} - x_i| < \delta$ for all i . This gives us disjoint intervals $[x_1 - \delta, x_1 + \delta], \dots, [x_n - \delta, x_n + \delta]$. Then we have

$$\begin{aligned} \sum_{i=1}^n o(f, x_i) &\leq \sum_{i=1}^n f(x_i + \delta) - f(x_i - \delta) \\ &= f(x_n + \delta) - f(x_n - \delta) + \dots + f(x_1 + \delta) - f(x_1 - \delta) \\ &\leq \underbrace{f(b) - f(x_n + \delta)}_{\geq 0} + f(x_n + \delta) - f(x_n - \delta) + \underbrace{f(x_n - \delta) - f(x_{n-1} + \delta)}_{\geq 0} \\ &\quad + f(x_{n-1} + \delta) - \dots - f(x_1 - \delta) + \underbrace{f(x_1 - \delta) - f(a)}_{\geq 0} \\ &= f(b) - f(a) \end{aligned}$$

The first and last intervals may be adjusted slightly for the case where $x_1 = a$ or $x_n = b$. \square

A.2 Chapter 2 Exercises

Exercise 2-1 Prove that if a function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$, then it is continuous at a .

Proof. Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable at $a \in \mathbb{R}^n$. Then $Df(a)$ is linear transformation. By Exercise 1-10, there exists a number $M > 0$ such that

$$\frac{|Df(a)(h)|}{|h|} \geq M, \quad \forall h \in \mathbb{R}^n$$

Then since f is differentiable at a , there exists $\delta > 0$ such that for any $|h| < \delta$,

$$\frac{|f(a+h) - f(a) - Df(a)(h)|}{|h|} < 1$$

Now let $\varepsilon > 0$ be arbitrary, and pick $\delta' = \min \left\{ \delta, \frac{\varepsilon}{M+1} \right\}$. Then for any x with $|x - a| < \delta'$ we have

$$\begin{aligned} |f(x) - f(a)| &= |f(x) - f(a) - Df(a)(x-a) + Df(a)(x-a)| \\ &\leq |f(x) - f(a) - Df(a)(x-a)| + |Df(a)(x-a)| \\ &< |x-a| + M|x-a| \\ &< (M+1) \frac{\varepsilon}{M+1} = \varepsilon \end{aligned} \quad \square$$

Exercise 2-2 A function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is **independent of the second variable** if for any $x \in \mathbb{R}$ and $y_1, y_2 \in \mathbb{R}$ we have $f(x, y_1) = f(x, y_2)$. Show that f is independent of the second variable if and only if there is a function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x, y) = g(x)$. What is $f'(a, b)$ in terms of g' ?

Proof. (\implies) Suppose f is independent of the second variable. Then define $g(x) = f(x, 0)$. For any x, y we have

$$f(x, y) = f(x, 0) = g(x)$$

(\impliedby) Suppose $g(x) = f(x, y)$. Then let $x, y_1, y_2 \in \mathbb{R}$ be arbitrary. We have

$$f(x, y_1) = g(x) = f(x, y_2) \quad \square$$

Claim: $f'(a, b) = [g'(a) \ 0]$.

Proof. Fix $(a, b) \in \mathbb{R}^2$. Then let $\varepsilon > 0$. Since g is differentiable at a , there exists $\delta > 0$ such that for any $|h| < \delta$,

$$\frac{|g(a+h) - g(a) - g'(a)(h)|}{|h|} < \varepsilon$$

Then if $h = (h_1, h_2)$ satisfies $|h| < \delta$, it must also be the case that $|h_1| \leq |(h_1, h_2)| < \delta$. Thus for any $|(h_1, h_2)| = |h| < \delta$, we have

$$\begin{aligned} \frac{\left| f(a+h_1, b+h_2) - f(a, b) - [g'(a) \ 0] \begin{bmatrix} h_1 \\ h_2 \end{bmatrix} \right|}{|h|} &= \frac{|g(a+h_1) - g(a) - g'(a)(h_1)|}{|h|} \\ &\leq \frac{|g(a+h_1) - g(a) - g'(a)(h_1)|}{|h_1|} < \varepsilon \end{aligned}$$

Thus we have $f'(a, b) = [g'(a) \ 0]$. \square

Exercise 2-3 Define when a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ is independent of the first variable, and find $f'(a, b)$ for such f . Which functions are independent of both the first and second variables?

A function $f; \mathbb{R}^2 \rightarrow \mathbb{R}$ is independent of the first variable if for any $x_1, x_2, y \in \mathbb{R}$ we have $f(x_1, y) = f(x_2, y)$, or equivalently if there exists $h : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x, y) = h(y)$. In this case, $f'(a, b) = [0 \ h'(b)]$. If a function is independent of both variables, then $f(a_1, b_1) = f(a_2, b_1) = f(a_2, b_2)$ for any $(a_1, b_1), (a_2, b_2) \in \mathbb{R}^2$ so f is constant.

Exercise 2-4 Let g be a continuous real-valued function on the unit circle such that $g(0, 1) = g(1, 0) = 0$ and $g(-x) = -g(x)$. Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} |x|g\left(\frac{x}{|x|}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$$

- (a) If $x \in \mathbb{R}^2$ and $h_x : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $h_x(t) = f(tx)$, show that h_x is differentiable.
- (b) Show that f is not differentiable at $(0, 0)$ unless $g = 0$ everywhere.

- (a) *Proof.* If $x = 0$ then h is identically 0 and is differentiable. If $x \neq 0$, then for $t \neq 0$ we have

$$h(t) = |tx|g\left(\frac{tx}{|tx|}\right) = |t||x|g\left(\underbrace{\text{sign}(t)\frac{x}{|x|}}_{g(-x)=-g(x)}\right) = |t|\text{sign}(t)|x|g\left(\frac{x}{|x|}\right) = t\left[|x|g\left(\frac{x}{|x|}\right)\right]$$

We also have $h(0) = f(0) = 0 = 0|x|g\left(\frac{x}{|x|}\right)$ so h is a linear function of t . Thus it is differentiable from single-variable analysis. \square

- (b) *Proof.* Suppose that f can be differentiated. Then since $Df(0, 0)$ is linear, it is uniquely determined by its behavior on the basis $\{e_1, e_2\}$. In particular, pick $\varepsilon > 0$. Then there exists a $\delta > 0$ such that whenever $0 < |h| < \delta$ we have

$$\frac{|f(h) - f(0, 0) - Df(0, 0)(h)|}{|h|} < \varepsilon$$

Then picking some $h_1 \in \mathbb{R}$ with $0 < |h_1| < \delta$,

$$|Df(0, 0)(e_1)| = \frac{|h_1 Df(0, 0)(e_1)|}{|h_1|} = \frac{|f(h_1 e_1) - f(0, 0) - \underbrace{Df(0, 0)(h_1 e_1)}_{=0}|}{|h_1 e_1|} < \varepsilon$$

This works for all epsilon, so $Df(0, 0)(e_1) = 0$. Similarly, $Df(0, 0)(e_2) = 0$, so $Df(0, 0)$ is the zero transformation. Now suppose $g(x) \neq 0$ for some x . Then for $\varepsilon = g(x)$ and arbitrary, δ ,

$$\frac{|f\left(\frac{\delta x}{2}\right) - f(0, 0) - \underbrace{Df(0, 0)\left(\frac{\delta x}{2}\right)}_{=0}|}{\left|\frac{\delta x}{2}\right|} = \frac{\frac{\delta}{2}g\left(\frac{\delta x/2}{\delta/2}\right)}{\frac{\delta}{2}} = g(x) \geq \varepsilon$$

so f is not differentiable. Thus f is only differentiable when $g(x) = 0$ everywhere. \square

Exercise 2-5 Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} \frac{x|y|}{\sqrt{x^2+y^2}}, & (x, y) \neq 0 \\ 0 & (x, y) = 0 \end{cases}$$

Show that f is a function of the kind considered in Exercise 2-4, so that f is not differentiable at $(0, 0)$.

Proof. Let

$$g(x, y) = \begin{cases} \frac{x|y|}{x^2+y^2}, & (x, y) \neq 0 \\ 0, & (x, y) = 0 \end{cases}$$

Then for $(x, y) \neq 0$ we have

$$\begin{aligned} |(x, y)|g\left(\frac{(x, y)}{|(x, y)|}\right) &= \sqrt{x^2+y^2}g\left(\frac{x}{\sqrt{x^2+y^2}}, \frac{y}{\sqrt{x^2+y^2}}\right) \\ &= \sqrt{x^2+y^2} \frac{x|y|}{x^2+y^2} \\ &= \frac{x|y|}{\sqrt{x^2+y^2}} \\ &= f(x, y) \end{aligned}$$

Moreover,

$$g(1, 0) = \frac{0}{\sqrt{1}} = 0 = \frac{|0|}{\sqrt{1}} = g(0, 1)$$

and

$$g(-x, -y) = \frac{-x|-y|}{(-x)^2+(-y)^2} = -\frac{x|y|}{x^2+y^2} = -g(x, y)$$

so f is of the form in Exercise 2-4. However,

$$g\left(\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}}\right) = \left(\frac{1}{\sqrt{2}}\right)^2 = \frac{1}{2} \neq 0$$

so g is not 0 everywhere and thus f is not differentiable at $(0, 0)$. □

Exercise 2-6 Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by $f(x, y) = \sqrt{|xy|}$. Show that f is not differentiable at $(0, 0)$.

Proof. Following the proof of Exercise 2-4 part (a), first suppose f is differentiable at $(0, 0)$. Then $Df(0, 0)$ exists, and it is determined by its behavior on the basis $\{e_1, e_2\}$. Letting $\varepsilon > 0$ be arbitrary, there must exist $\delta > 0$ such that for any $0 < |h| < \delta$,

$$\frac{|f(h) - f(0, 0) - Df(0, 0)(h)|}{|h|} < \varepsilon$$

Pick some $h_1 \in \mathbb{R}$ with $0 < |h_1| < \delta$. Then

$$|Df(0,0)(e_1)| = \frac{|h_1 Df(0,0)(e_1)|}{|h_1|} = \frac{|f(h_1 e_1) - f(0,0) - Df(0,0)(h_1 e_1)|}{|h_1 e_1|} < \varepsilon$$

So $|Df(0,0)(e_1)| < \varepsilon$ for all ε , and thus $Df(0,0)(e_1) = 0$. Similarly, $Df(0,0)(e_2) = 0$, so $Df(0,0)$ is the zero transformation. However, let $\varepsilon = \frac{1}{\sqrt{2}}$, and let $\delta > 0$ be arbitrary. Then the point $(x, y) = (\frac{\delta}{\sqrt{3}}, \frac{\delta}{\sqrt{3}})$ satisfies $0 < |(x, y)| < \delta$, but

$$\frac{|f(x, y) - f(0,0) - Df(0,0)(x, y)|}{|(x, y)|} = \frac{\sqrt{\frac{\delta^2}{3}}}{\sqrt{\frac{2\delta^2}{3}}} = \frac{1}{\sqrt{2}} \geq \varepsilon$$

so no δ works and f is not differentiable. □

Exercise 2-7 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be a function such that $|f(x)| \leq |x|^2$. Show that f is differentiable at $\mathbf{0}$.

Proof. Let $\lambda : \mathbb{R}^n \rightarrow \mathbb{R}$ be the zero transformation. Then let $\varepsilon > 0$ be arbitrary, and set $\delta = \varepsilon$. Whenever $0 < |x| < \delta$, by assumption we have

$$\frac{|f(x)|}{|x|} \leq |x|$$

In particular, $|f(0)| \leq |0|^2 = 0$ so $f(0) = 0$. Thus

$$\frac{|f(x) - f(0) - \lambda(x)|}{|x|} = \frac{|f(x)|}{|x|} \leq |x| < \delta = \varepsilon$$

so f is differentiable at 0 with derivative $Df(0) = \lambda$ the zero transformation. □

Exercise 2-8 Let $f : \mathbb{R} \rightarrow \mathbb{R}^2$. Prove that f is differentiable at $a \in \mathbb{R}$ if and only if f^1 and f^2 are, and that in this case

$$f'(a) = \begin{bmatrix} (f^1)'(a) \\ (f^2)'(a) \end{bmatrix}$$

Proof. (\implies) Suppose f is differentiable at $a \in \mathbb{R}$. Then let $\varepsilon > 0$ be arbitrary. Since f is differentiable, there exists $\delta > 0$ such that whenever $0 < |h| < \delta$ we have

$$\frac{|f(a+h) - f(a) - Df(a)(h)|}{|h|} < \varepsilon$$

If we suppose that $Df(a)(h)$ has matrix representation given by

$$f'(a) = \begin{bmatrix} b \\ c \end{bmatrix}$$

then it is the case that

$$Df(a)(h) = \begin{bmatrix} bh \\ ch \end{bmatrix}$$

Now if we write for convenience $(x, y) = f(a+h) - f(a) - Df(a)(h)$, then we know that $|x| \leq |(x, y)|$, so whenever $0 < |h| < \delta$

$$\frac{|f^1(a+h) - f(a) - bh|}{|h|} = \frac{|x|}{|h|} \leq \frac{|(x, y)|}{|h|} = \frac{|f(a+h) - f(a) - Df(a)(h)|}{|h|} < \varepsilon$$

so f^1 is differentiable at a . The proof for f^2 is similar. Moreover, this proves that in this case $bh = (f^1)'(a)$ and $ch = (f^2)'(a)$, so that

$$f'(a) = \begin{bmatrix} (f^1)'(a) \\ (f^2)'(a) \end{bmatrix}$$

(\Leftarrow) Now suppose that f^1 and f^2 are differentiable at a . Let $\varepsilon > 0$ be arbitrary. Then there exist $\delta_1, \delta_2 > 0$ such that whenever $0 < |h| < \delta_1$ we have

$$\frac{|f^1(a+h) - f^1(a) + (f^1)'(a)(h)|}{|h|} < \frac{\varepsilon}{2}$$

and whenever $0 < |h| < \delta_2$ we have

$$\frac{|f^2(a+h) - f^2(a) + (f^2)'(a)(h)|}{|h|} < \frac{\varepsilon}{2}$$

Let $\delta = \min\{\delta_1, \delta_2\}$. Let $\lambda : \mathbb{R} \rightarrow \mathbb{R}^2$ have the matrix

$$[\lambda] = \begin{bmatrix} (f^1)'(a) \\ (f^2)'(a) \end{bmatrix}$$

Then whenever $0 < |h| < \delta$,

$$\frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = \frac{\left\| \begin{bmatrix} x \\ y \end{bmatrix} \right\|}{|h|}$$

where

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} f^1(a+h) - f^1(a) - (f^1)'(a)(h) \\ f^2(a+h) - f^2(a) - (f^2)'(a)(h) \end{bmatrix}$$

Then

$$\frac{|f(a+h) - f(a) - \lambda(h)|}{|h|} = \frac{\left\| \begin{bmatrix} x \\ y \end{bmatrix} \right\|}{|h|} \leq \frac{|x|}{|h|} + \frac{|y|}{|h|} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

so f is differentiable at a , and once again we have

$$f'(a) = [\lambda] = \begin{bmatrix} (f^1)'(a) \\ (f^2)'(a) \end{bmatrix}$$

□

Exercise 2-9 Two functions $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are **equal up to n th order** at $a \in \mathbb{R}$ if

$$\lim_{h \rightarrow 0} \frac{f(a+h) - g(a+h)}{h^n} = 0$$

- (a) Show that a continuous function f is differentiable at a if and only if there is a function g of the form $g(x) = a_0 + a_1(x-a)$ such that f and g are equal up to first order at a . (**Note:** Spivak did not assume continuity in the original exercise, but it is required in the if direction, and continuity in the only if direction follows from differentiability).
- (b) If $f'(a), \dots, f^{(n)}(a)$ exist, show that f and the function g defined by

$$g(x) = \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x-a)^i$$

are equal up to n th order at a . (This is the n th degree Taylor polynomial of f expanded about a).

- (a) *Proof.* (\implies) Suppose f is differentiable at a . Then define

$$g(x) = f(a) + f'(a)(x-a)$$

We have

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a+h)}{h} = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - f'(a)h}{h} = 0$$

since f is differentiable, so f and g are equal up to first order.

(\impliedby) Now suppose $g(x) = a_0 + a_1(x-a)$ is equal to f up to first order. Since f (and g) are continuous,

$$\lim_{h \rightarrow 0} f(a+h) - g(a+h) = f(a) - g(a) = 0$$

so $f(a) = g(a) = a_0$. Thus we have

$$\lim_{h \rightarrow 0} \frac{f(a+h) - f(a) - a_1h}{h} = \lim_{h \rightarrow 0} \frac{f(a+h) - g(a+h)}{h} = 0$$

so f is differentiable at a . □

- (b) *Proof.* We induct on n . Suppose that for any function f , whenever $f'(a), \dots, f^{(n-1)}(a)$ exist, then

$$f(x) \stackrel{n-1}{\sim} \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (x-a)^i$$

where $\overset{n-1}{\sim}$ represents equality up to order $n - 1$. Now suppose that $f'(a), \dots, f^{(n)}(a)$ all exist. Then we have

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(a+h) - g(a+h)}{h^n} &= \lim_{h \rightarrow 0} \frac{f(a+h) - \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} (a+h-a)^i}{h^n} \\ &= \lim_{h \rightarrow 0} \frac{f(a+h) - \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} h^i}{h^n} \end{aligned}$$

Note that since f and g are continuous (where f is continuous since it is differentiable), we have

$$\lim_{h \rightarrow 0} f(a+h) - \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} h^i = f(a) - \frac{f^{(0)}(a)}{0!} - \sum_{i=1}^n \frac{f^{(i)}(a)}{i!} 0^i = f(a) - f(a) = 0$$

Clearly g is differentiable and so is f , so $f(a+h) - g(a+h)$ is differentiable, and thus L'Hopital's Rule applies. So

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(a+h) - \sum_{i=0}^n \frac{f^{(i)}(a)}{i!} h^i}{h^n} &\stackrel{\text{LH}}{=} \lim_{h \rightarrow 0} \frac{f'(a+h) - \sum_{i=1}^n \frac{f^{(i)}(a)}{(i-1)!} h^{i-1}}{nh^{n-1}} \\ &= \lim_{h \rightarrow 0} \frac{f'(a+h) - \sum_{i=0}^{n-1} \frac{f^{(i+1)}(a)}{i!} h^i}{h^{n-1}} \\ &= \lim_{h \rightarrow 0} \frac{f'(a+h) - \sum_{i=0}^{n-1} \frac{(f')^{(i)}(a)}{i!} h^i}{h^{n-1}} \end{aligned}$$

Since $f''(a), \dots, f^{(n)}(a)$ all exist, $(f')'(a), \dots, (f')^{(n-1)}(a)$ all exist, so the inductive hypothesis applies and

$$f'(x) \overset{n-1}{\sim} \sum_{i=0}^{n-1} \frac{(f')^{(i)}(a)}{i!} (x-a)^i$$

so

$$\lim_{h \rightarrow 0} \frac{f'(a+h) - \sum_{i=0}^{n-1} \frac{(f')^{(i)}(a)}{i!} h^i}{h^{n-1}} = 0$$

Thus

$$\lim_{h \rightarrow 0} \frac{f(a+h) - g(a+h)}{h^n} = \lim_{h \rightarrow 0} \frac{f'(a+h) - \sum_{i=0}^{n-1} \frac{(f')^{(i)}(a)}{i!} h^i}{h^{n-1}} = 0$$

so f and g are equal up to n th order. □

Exercise 2-10 Use the theorems of this section [Section 2.2] to find f' for the following:

- (a) $f(x, y, z) = x^y$.
- (b) $f(x, y, z) = (x^y, z)$.
- (c) $f(x, y) = \sin(x \sin y)$.
- (d) $f(x, y, z) = \sin(x \sin(y \sin z))$.
- (e) $f(x, y, z) = x^{y^z}$.
- (f) $f(x, y, z) = x^{y+z}$.
- (g) $f(x, y, z) = (x + y)^z$.
- (h) $f(x, y) = \sin(xy)$.
- (i) $f(x, y) = [\sin(xy)]^{\cos 3}$.
- (j) $f(x, y) = (\sin(xy), \sin(x \sin y), x^y)$.

(a) We write

$$f = [\pi^1]^{[\pi^2]} = (e^{\ln \circ [\pi^1]})^{[\pi^2]} = e^{\pi^2 \cdot \ln \circ \pi^1}$$

. Then

$$\begin{aligned} f'(a, b, c) &= (e^{\pi^2 \cdot \ln \circ \pi^1})'(a, b, c) \\ &= e^{b \ln a} (\pi^2 \cdot \ln \circ \pi^1)'(a, b, c) \\ &= a^b (\ln a (\pi^2)'(a, b, c) + b (\ln \circ \pi^1)'(a, b, c)) \\ &= a^b (\ln a \pi^2 + b \frac{1}{a} (\pi^1)')(a, b, c) \\ &= a^b (\ln a \pi^2 + \frac{b}{a} \pi^1) \\ &= (ba^{b-1}, a^b \ln a, 0) \end{aligned}$$

(b) Following easily from part (a) we have:

$$\begin{aligned} f'(a, b, c) &= \begin{bmatrix} - & (x^y)'(a, b, c) & - \\ - & (\pi^3)'(a, b, c) & - \end{bmatrix} \\ &= \begin{bmatrix} ba^{b-1} & a^b \ln a & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(c) Similarly to the example, we have $f = \sin \circ (\pi^1 \cdot \sin \circ \pi^2)$. Thus,

$$\begin{aligned}
f'(a, b) &= (\sin \circ (\pi^1 \cdot \sin \circ \pi^2))'(a, b) \\
&= \cos(a \sin b) (\pi^1 \cdot \sin \circ \pi^2)'(a, b) \\
&= \cos(a \sin b) (\sin b (\pi^1)'(a, b) + a (\sin \circ \pi^2)'(a, b)) \\
&= \cos(a \sin b) \sin b \pi^1 + a \cos(a \sin b) \cos b \pi^2 \\
&= (\cos(a \sin b) \sin b, a \cos(a \sin b) \cos b)
\end{aligned}$$

(d) As above, we have

$$f = \sin \circ (\pi^1 \cdot (\sin \circ (\pi^2 \cdot (\sin \circ \pi^3))))$$

so

$$\begin{aligned}
f'(a, b, c) &= (\sin \circ (\pi^1 \cdot (\sin \circ (\pi^2 \cdot (\sin \circ \pi^3)))))'(a, b, c) \\
&= \cos(a \sin(b \sin c)) (\pi^1 \cdot (\sin \circ (\pi^2 \cdot (\sin \circ \pi^3))))'(a, b, c) \\
&= \cos(a \sin(b \sin c)) (\sin(b \sin c) \pi^1 + a \cos(b \sin c) (\pi^2 \cdot (\sin \circ \pi^3))'(a, b, c)) \\
&= \cos(a \sin(b \sin c)) (\sin(b \sin c) \pi^1 + a \cos(b \sin c) (\sin c \pi^2 + b \cos c \pi^3)) \\
&= \cos(a \sin(b \sin c)) * (\sin(b \sin c), a \cos(b \sin c) \sin c, ab \cos(b \sin c) \cos c)
\end{aligned}$$

(e) Let $g(x, y) = x^y$. Then we have

$$f(x, y, z) = g(x, g(y, z))$$

so that

$$f = g \circ (\pi^1, g \circ (\pi^2, \pi^3))$$

Using our result from part (a),

$$\begin{aligned}
f'(a, b, c) &= g'(a, g(b, c)) \begin{bmatrix} - & (\pi^1)'(a, b, c) & - \\ - & (g \circ (\pi^2, \pi^3))'(a, b, c) & - \end{bmatrix} \\
&= [b^c a^{b^c-1} \quad a^{b^c} \ln a] \begin{bmatrix} 1 & 0 & 0 \\ 0 & cb^{c-1} & b^c \ln b \end{bmatrix} \\
&= [b^c a^{b^c-1} \quad a^{b^c} cb^{c-1} \ln a \quad a^{b^c} b^c \ln a \ln b]
\end{aligned}$$

(f) Letting g be as defined in part (e), we have

$$f = g \circ (\pi^1, \pi^2 + \pi^3)$$

Thus

$$\begin{aligned}
f'(a, b, c) &= (g \circ (\pi^1, \pi^2 + \pi^3))'(a, b, c) \\
&= g'(a, b + c) \begin{bmatrix} - & (\pi^1)'(a, b, c) & - \\ - & (\pi^2 + \pi^3)'(a, b, c) & - \end{bmatrix} \\
&= [(b + c) a^{b+c-1} \quad a^{b+c} \ln a] \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \\
&= [(b + c) a^{b+c-1} \quad a^{b+c} \ln a \quad a^{b+c} \ln a]
\end{aligned}$$

(g) Again letting g be as in part (e), we have

$$f = g \circ (\pi^1 + \pi^2, \pi^3)$$

so that

$$\begin{aligned} f'(a, b, c) &= (g \circ (\pi^1 + \pi^2, \pi^3))'(a, b, c) \\ &= g'(a + b, c) \begin{bmatrix} - & (\pi^1 + \pi^2)'(a, b, c) & - \\ - & (\pi^3)'(a, b, c) & - \end{bmatrix} \\ &= [c(a + b)^{c-1} \quad (a + b)^c \ln(a + b)] \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \\ &= [c(a + b)^{c-1} \quad c(a + b)^{c-1} \quad (a + b)^c \ln(a + b)] \end{aligned}$$

(h) We can straightforwardly write this as

$$f = \sin \circ (\pi^1 \cdot \pi^2)$$

Then

$$\begin{aligned} f'(a, b) &= (\sin \circ (\pi^1 \cdot \pi^2))'(a, b) \\ &= \cos(ab)(b\pi^1 + a\pi^2) \\ &= (b \cos(ab), a \cos(ab)) \end{aligned}$$

(i) Using the same definition of g ,

$$f = g \circ (\sin \circ (\pi^1 \cdot \pi^2), \cos 3)$$

Since $\cos 3$ is constant,

$$\begin{aligned} f'(a, b) &= (g \circ (\sin \circ (\pi^1 \cdot \pi^2), \cos 3))'(a, b) \\ &= g'(\sin(ab), \cos 3) \begin{bmatrix} - & (\pi^1 \cdot \pi^2)'(a, b) & - \\ - & (\cos 3)'(a, b) & - \end{bmatrix} \\ &= [\cos 3 [\sin(ab)]^{\cos 3} \quad [\sin(ab)]^{\cos 3} \ln \sin(ab)] \begin{bmatrix} b & a \\ 0 & 0 \end{bmatrix} \\ &= [b \cos 3 [\sin(ab)]^{\cos 3} \quad a \cos 3 [\sin(ab)]^{\cos 3}] \end{aligned}$$

(j) From parts (h), (c), and (a), respectively, we already know that

$$\begin{aligned} (\sin(xy))'(a, b) &= [b \cos(ab) \quad a \cos(ab)] \\ (\sin(x \sin y))'(a, b) &= [\cos(a \sin b) \sin b \quad a \cos(a \sin b) \cos b] \\ (x^y)'(a, b) &= [ba^{b-1} \quad a^b \ln a] \end{aligned}$$

Then f' is simply given by putting each of these matrices in as row vectors, such that

$$f'(a, b, c) = \begin{bmatrix} - & (\sin(xy))'(a, b) & - \\ - & (\sin(x \sin y))'(a, b) & - \\ - & (x^y)'(a, b) & - \end{bmatrix} = \begin{bmatrix} b \cos(ab) & a \cos(ab) \\ \cos(a \sin b) \sin b & a \cos(a \sin b) \cos b \\ ba^{b-1} & a^b \ln a \end{bmatrix}$$

Exercise 2-11 Find f' for the following (where $g : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, and $s \in \mathbb{R}$ is fixed):

(a) $f(x, y) = \int_s^{x+y} g.$

(b) $f(x, y) = \int_s^{xy} g.$

(c) $f(x, y, z) = \int_{xy}^{\sin(x \sin(y \sin z))} g.$

(a) Define $F : \mathbb{R} \rightarrow \mathbb{R}$ by

$$F(x) = \int_s^x g(t) dt$$

Since g is continuous, the fundamental theorem of calculus tells us that

$$F'(x) = g(x)$$

Then we can here write f as

$$f = F \circ (\pi^1 + \pi^2)$$

so that

$$\begin{aligned} f'(a, b) &= (F \circ (\pi^1 + \pi^2))'(a, b) \\ &= F'(a + b)(\pi^1 + \pi^2)'(a, b) \\ &= g(a + b)(\pi^1 + \pi^2) \\ &= (g(a + b), g(a + b)) \end{aligned}$$

(b) Similarly, write

$$f = F \circ (\pi^1 \cdot \pi^2)$$

Then

$$\begin{aligned} f'(a, b) &= (F \circ (\pi^1 \cdot \pi^2))'(a, b) \\ &= (bg(ab), ag(ab)) \end{aligned}$$

(c) First note that we can pick any $s \in \mathbb{R}$ and separate this integral:

$$f(x, y, z) = \int_{xy}^{\sin(x \sin(y \sin z))} g = \int_s^{\sin(x \sin(y \sin z))} g + \int_{xy}^s g = \int_s^{\sin(x \sin(y \sin z))} g - \int_s^{xy} g$$

Then using the same method as parts (a) and (b) of this problem, and using the results from parts (d) and (a) of Exercise 2-10, the Jacobian of the first term, evaluated at (a, b, c) , is given by

$$\begin{bmatrix} g(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \sin(b \sin c) \\ ag(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \cos(b \sin c) \sin c \\ abg(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \cos(b \sin c) \cos c \end{bmatrix}^T$$

and the Jacobian of the second by

$$\begin{bmatrix} g(a^b)ba^{b-1} \\ g(a^b)a^b \ln a \\ 0 \end{bmatrix}$$

Thus we have

$$f'(a, b, c) = \begin{bmatrix} g(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \sin(b \sin c) - g(a^b)ba^{b-1} \\ ag(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \cos(b \sin c) \sin c - g(a^b)a^b \ln a \\ abg(\sin(a \sin(b \sin c))) \cos(a \sin(b \sin c)) \cos(b \sin c) \cos c \end{bmatrix}^T$$

Exercise 2-12 A function $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^p$ is **bilinear** if for $x, x_1, x_2 \in \mathbb{R}^n$, $y, y_1, y_2 \in \mathbb{R}^m$ and $a \in \mathbb{R}$ we have

$$\begin{aligned} f(ax, y) &= af(x, y) = f(x, ay) \\ f(x_1 + x_2, y) &= f(x_1, y) + f(x_2, y) \\ f(x, y_1 + y_2) &= f(x, y_1) + f(x, y_2) \end{aligned}$$

(a) Prove that if f is bilinear, then

$$\lim_{(h,k) \rightarrow \mathbf{0}} \frac{|f(h, k)|}{|(h, k)|} = 0$$

(b) Prove that $Df(a, b)(x, y) = f(a, y) + f(x, b)$.

(c) Show that the formula for $Dp(a, b)$ in Section 2.2 is a special case of (b).

(a) *Proof.* Suppose f is bilinear, and suppose $h = (h_1, \dots, h_n)$, $k = (k_1, \dots, k_m)$. Then we can write

$$\begin{aligned} \lim_{(h,k) \rightarrow \mathbf{0}} \frac{|f(h, k)|}{|(h, k)|} &= \lim_{(h,k) \rightarrow \mathbf{0}} \frac{|f(\sum_{i=1}^n h_i e_i, \sum_{j=1}^m k_j e_j)|}{|(h, k)|} \\ &= \lim_{(h,k) \rightarrow \mathbf{0}} \frac{|\sum_{i=1}^n \sum_{j=1}^m h_i k_j f(e_i, e_j)|}{|(h, k)|} \\ &\leq \sum_{i=1}^n \sum_{j=1}^m |f(e_i, e_j)| \lim_{(h_i, k_j) \rightarrow \mathbf{0}} \frac{|h_i k_j|}{|(h, k)|} \\ &\leq \sum_{i=1}^n \sum_{j=1}^m |f(e_i, e_j)| \lim_{(h_i, k_j) \rightarrow \mathbf{0}} \frac{|h_i k_j|}{|(h_i, k_j)|} \end{aligned}$$

Now we proved in the proof of $Dp(a, b)$ that

$$\lim_{(h_i, k_j) \rightarrow \mathbf{0}} \frac{|h_i k_j|}{|(h_i, k_j)|} = 0$$

so we have

$$\lim_{(h,k) \rightarrow \mathbf{0}} \frac{|f(h,k)|}{|(h,k)|} = 0 \quad \square$$

(b) *Proof.* Note that

$$\begin{aligned} f(a+x, b+y) - f(a,b) - f(a,y) - f(x,b) &= f(a+x, b+y) - f(a, b+y) - f(x,b) \\ &= f(a+x, b+y) - f(a, b+y) - f(x,b) - f(x,y) + f(x,y) \\ &= f(a+x, b+y) - f(a, b+y) - f(x, b+y) + f(x,y) \\ &= f(a+x, b+y) - f(a+x, b+y) + f(x,y) \\ &= f(x,y) \end{aligned}$$

Then we have

$$\lim_{(x,y) \rightarrow \mathbf{0}} \frac{|f(a+x, b+y) - f(a,b) - f(a,y) - f(x,b)|}{|(x,y)|} = \lim_{(x,y) \rightarrow \mathbf{0}} \frac{|f(x,y)|}{|(x,y)|}$$

and by part (a) we know this limit is 0. □

(c) *Proof.* Note that our work in part (a) implies that f is completely determined by its values on the various pairs (e_i, e_j) . So $Dp(a, b)$ is simply the case where $n = m = 1$ and $f(1, 1) = 1$. □

Exercise 2-13 Define $IP : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ by $IP(x, y) = \langle x, y \rangle$.

- (a) Find $D(IP)(a, b)$ and $(IP)'(a, b)$.
- (b) If $f, g : \mathbb{R} \rightarrow \mathbb{R}^n$ are differentiable and $h : \mathbb{R} \rightarrow \mathbb{R}$ is defined by $h(t) = \langle f(t), g(t) \rangle$, show that

$$h'(a) = \langle f'(a)^T, g(a) \rangle + \langle f(a), g'(a)^T \rangle$$

(Note that $f'(a)$ is an $n \times 1$ matrix; its transpose $f'(a)^T$ is a $1 \times n$ matrix, which we consider as a member of \mathbb{R}^n .)

- (c) If $f : \mathbb{R} \rightarrow \mathbb{R}^n$ is differentiable and $|f(t)| = 1$ for all t , show that $\langle f'(t)^T, f(t) \rangle = 0$.
- (d) Exhibit a differentiable function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that the function $|f|$ defined by $|f|(t) = |f(t)|$ is not differentiable.

(a) Since the (real) inner product is bilinear by definition, we can apply Exercise 2-12 to conclude that

$$D(IP)(a, b)(x, y) = IP(a, y) + IP(x, b) = \langle a, y \rangle + \langle x, b \rangle = \langle a, y \rangle + \langle b, x \rangle = \langle (b, a), (x, y) \rangle$$

Moreover, we can rewrite this to be

$$D(IP)(a, b)(x, y) = (b, a)(x, y)^T$$

from which we can conclude that

$$(IP)'(a, b) = (b, a)$$

where (b, a) is the $1 \times 2n$ matrix given by concatenating the row vectors b and a .

(b) *Proof.* Directly from the definition of h , we have

$$h = IP \circ (f, g)$$

so the chain rule says that

$$\begin{aligned} h'(a) &= IP'(f(a), g(a)) \begin{bmatrix} | & | \\ f'(a) & \\ | & | \\ | & | \\ g'(a) & \\ | & | \end{bmatrix} \\ &= [-g(a) \quad -f(a)] \begin{bmatrix} | & | \\ f'(a) & \\ | & | \\ | & | \\ g'(a) & \\ | & | \end{bmatrix} \\ &= [-g(a)] \begin{bmatrix} | & | \\ f'(a) & \\ | & | \end{bmatrix} + [-f(a)] \begin{bmatrix} | & | \\ g'(a) & \\ | & | \end{bmatrix} \\ &= \langle g(a), f'(a)^T \rangle + \langle f(a), g'(a)^T \rangle \\ &= \langle f'(a)^T, g(a) \rangle + \langle f(a), g'(a)^T \rangle \quad \square \end{aligned}$$

(c) *Proof.* Define

$$h(t) := \langle f(t), f(t) \rangle = \sqrt{|f(t)|}$$

Then by part (b),

$$h'(t) = 2 \langle f'(t)^T, f(t) \rangle$$

But the assumption that $|f(t)|$ is identically 1 means that h is constant, and thus

$$\langle f'(t)^T, f(t) \rangle = \frac{h'(t)}{2} = 0 \quad \square$$

(d) The identity function satisfies this, since $x \mapsto |x|$ is not differentiable at $x = 0$.

Exercise 2-14 Let E_i , $i = 1, \dots, k$ be Euclidean spaces of various dimensions. A function $f : E_1 \times \dots \times E_k \rightarrow \mathbb{R}^p$ is called **multilinear** if for each choice of $x_j \in E_j$, $j \neq i$, the function $g : E_i \rightarrow \mathbb{R}^p$ defined by $g(x) = f(x_1, \dots, x_{i-1}, x, x_{i+1}, \dots, x_k)$ is a linear transformation.

- (a) If f is multilinear and $i \neq j$, show that for $h = (h_1, \dots, h_k)$ with $h_l \in E_l$, we have

$$\lim_{h \rightarrow \mathbf{0}} \frac{|f(a_1, \dots, h_i, \dots, h_j, \dots, a_k)|}{|h|} = 0$$

- (b) Prove that

$$Df(a_1, \dots, a_k)(x_1, \dots, x_k) = \sum_{i=1}^k f(a_1, \dots, a_{i-1}, x_i, a_{i+1}, \dots, a_k)$$

- (a) *Proof.* Suppose that $\dim E_i = k_1$ and $\dim E_j = k_2$. Then define the function $g : \mathbb{R}^{k_1} \times \mathbb{R}^{k_2} \rightarrow \mathbb{R}^p$ by

$$g(x, y) = f(a_1, \dots, x, \dots, y, \dots, a_k)$$

Then we need to prove that

$$\lim_{(h_i, h_j) \rightarrow \mathbf{0}} \frac{|g(h_i, h_j)|}{|(h_i, h_j)|} = 0$$

To do this, we first prove that g is bilinear. Using multilinearity, we have that

$$g(ax, y) = f(a_1, \dots, ax, \dots, y, \dots, a_k) = af(a_1, \dots, x, \dots, y, \dots, a_k) = ag(x, y)$$

and

$$\begin{aligned} g(x_1 + x_2, y) &= f(a_1, \dots, x_1 + x_2, \dots, y, \dots, a_k) \\ &= f(a_1, \dots, x_1, \dots, y, \dots, a_k) + f(a_1, \dots, x_2, \dots, y, \dots, a_k) \\ &= g(x_1, y) + g(x_2, y) \end{aligned}$$

The last property is similar. So g is bilinear, and Exercise 2-12 part (a) tells us that

$$\lim_{h \rightarrow \mathbf{0}} \frac{|f(a_1, \dots, h_i, \dots, h_j)|}{|h|} = \lim_{(h_i, h_j) \rightarrow \mathbf{0}} \frac{|g(h_i, h_j)|}{|(h_i, h_j)|} = 0 \quad \square$$

- (b) *Proof.* For notational convenience, we define the following. Given a set of distinct indices $i_1, \dots, i_n \in [1, k]$, and vectors $\vec{a} = (a_1, \dots, a_k)$, $\vec{h} = (h_1, \dots, h_k)$, we write

$$f_{\{i_1, \dots, i_n\}}(\vec{a}, \vec{h}) = f(a_1, \dots, h_{i_1}, \dots, h_{i_2}, \dots, h_{i_n}, \dots, a_k)$$

In other words, if $S \subseteq [1, k]$, then $f_S(\vec{a}, \vec{h})$ passes in h_i if $i \in S$ and a_i otherwise.

Now, we prove an extension of part (a), namely, that for any k -linear function f , if we pick $n \leq k$ indices i_1, \dots, i_n , then

$$\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f_{\{i_1, \dots, i_n\}}(\vec{a}, \vec{h})|}{|\vec{h}|} = 0$$

We skip the proof that $f_{\{i_1, \dots, i_n\}}$ is n -linear, so this reduces to simply showing that for any multilinear function ($n > 1$) we have

$$\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(\vec{h})|}{|\vec{h}|} = 0$$

Let $d_i = \dim E_i$ for each i . Suppose also that $h_i = (h_{i,1}, \dots, h_{i,d_i})$. Then

$$\begin{aligned} \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(h_1, \dots, h_k)|}{|\vec{h}|} &= \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(\sum_{j_1=1}^{d_1} h_{1,j_1}, \dots, \sum_{j_k=1}^{d_k} h_{k,j_k})|}{|\vec{h}|} \\ &= \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|\sum_{j_1=1}^{d_1} \dots \sum_{j_k=1}^{d_k} h_{1,j_1} \dots h_{k,j_k} f(e_{j_1}, \dots, e_{j_k})|}{|\vec{h}|} \\ &\leq \sum_{j_1=1}^{d_1} \dots \sum_{j_k=1}^{d_k} |f(e_{j_1}, \dots, e_{j_k})| \lim_{(h_{1,j_1}, \dots, h_{k,j_k}) \rightarrow \mathbf{0}} \frac{|h_{1,j_1} \dots h_{k,j_k}|}{|\vec{h}|} \\ &\leq \sum_{j_1=1}^{d_1} \dots \sum_{j_k=1}^{d_k} |f(e_{j_1}, \dots, e_{j_k})| \lim_{(h_{1,j_1}, \dots, h_{k,j_k}) \rightarrow \mathbf{0}} \frac{|h_{1,j_1} \dots h_{k,j_k}|}{|(h_{1,j_1}, \dots, h_{k,j_k})|} \\ &= \sum_{j_1=1}^{d_1} \dots \sum_{j_k=1}^{d_k} |f(e_{j_1}, \dots, e_{j_k})| \cdot 0 \\ &= 0 \end{aligned}$$

Thus we have shown that any multilinear function satisfies

$$\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(\vec{h})|}{|\vec{h}|} = 0$$

Now, I claim that

$$f(a_1 + x_1, \dots, a_k + x_k) = f(\vec{a} + \vec{x}) = \sum_{S \in \mathcal{P}([1,k])} f_S(\vec{a}, \vec{x})$$

where $\mathcal{P}([1,k])$ represents the set of all subsets of $[1,k]$. We prove this by induction. Supposing it is true for $k-1$, we can then partition $\mathcal{P}([1,k])$ into X , consisting of those subsets which contain k , and A , consisting of those subsets which do not. Then

$$\sum_{S \in \mathcal{P}([1,k])} f_S(\vec{a}, \vec{x}) = \sum_{S \in X} f_S(\vec{a}, \vec{x}) + \sum_{S \in A} f_S(\vec{a}, \vec{x})$$

Now, the inductive hypothesis applies, and we have

$$\sum_{S \in X} f_S(\vec{a}, \vec{x}) = f(a_1 + x_1, \dots, a_{k-1} + x_{k-1}, x_k)$$

and

$$\sum_{S \in A} f_S(\vec{a}, \vec{x}) = f(a_1 + x_1, \dots, a_{k-1} + x_{k-1}, a_k)$$

and by applying multilinearity, we conclude that

$$\begin{aligned} \sum_{S \in \mathcal{P}([1,k])} f_S(\vec{a}, \vec{x}) &= \sum_{S \in X} f_S(\vec{a}, \vec{x}) + \sum_{S \in A} f_S(\vec{a}, \vec{x}) \\ &= f(a_1 + x_1, \dots, a_{k-1} + x_{k-1}, x_k) + f(a_1 + x_1, \dots, a_{k-1} + x_{k-1}, a_k) \\ &= f(\vec{a} + \vec{x}) \end{aligned}$$

Lastly, we have

$$\begin{aligned} &\lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|f(\vec{a} + \vec{h}) - f(\vec{a}) - \sum_{i=1}^k f_{\{i\}}(\vec{a}, \vec{h})|}{|\vec{h}|} \\ &= \lim_{\vec{h} \rightarrow \mathbf{0}} \frac{|\sum_{S \in \mathcal{P}([1,k])} f_S(\vec{a}, \vec{h}) - f(\vec{a}) - \sum_{i=1}^k f_{\{i\}}(\vec{a}, \vec{h})|}{|\vec{h}|} \end{aligned}$$

Now, after cancelling, the numerator will be left only with terms of the form $f_S(\vec{a}, \vec{h})$ where S contains at least two elements, and f_S is therefore n -linear for $n > 1$. Thus the first part of this proof shows that the quotient goes to 0. \square

Exercise 2-15 Regard an $n \times n$ matrix as a point in the n -fold product $\mathbb{R}^n \times \dots \times \mathbb{R}^n$ by considering each column as a member of \mathbb{R}^n . (**Note:** Spivak considers the rows as elements of \mathbb{R}^n , but we use columns here for convention.)

(a) Prove that $\det : \mathbb{R}^n \times \dots \times \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable and

$$D(\det)(a_1, \dots, a_n)(x_1, \dots, x_n) = \sum_{i=1}^n \det \begin{bmatrix} | & & | & & | \\ a_1 & \dots & x_i & \dots & a_n \\ | & & | & & | \end{bmatrix}$$

(b) If $a_{ij} : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable, let $A(t)$ be the matrix such that $A(t)_{ij} = a_{ij}(t)$. If $f(t) = \det(A(t))$, show that

$$f'(t) = \sum_{j=1}^n \det \begin{bmatrix} a_{11}(t) & \dots & a'_{1j}(t) & \dots & a_{1n}(t) \\ \vdots & & \vdots & & \vdots \\ a_{n1}(t) & \dots & a'_{nj}(t) & \dots & a_{nn}(t) \end{bmatrix}$$

(c) If $\det(A(t)) \neq 0$ for all t and $b_1, \dots, b_n : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable, let $s_1, \dots, s_n : \mathbb{R} \rightarrow \mathbb{R}$ be the functions such that $s_1(t), \dots, s_n(t)$ are solutions of the equations

$$\sum_{j=1}^n a_{ij}(t)s_j(t) = b_i(t)$$

Show s_i is differentiable and find $s'_i(t)$.

(a) *Proof.* We take it for granted that \det is multilinear, as this is one possible definition of the determinant, and otherwise can easily be concluded from Laplace expansion along various columns. Then \det is differentiable by Exercise 2-14 part (b), and moreover the result from that problem shows that

$$D(\det)(a_1, \dots, a_n)(x_1, \dots, x_n) = \sum_{i=1}^n \det \begin{bmatrix} | & & | & & | \\ a_1 & \dots & x_i & \dots & a_n \\ | & & | & & | \end{bmatrix} \quad \square$$

(b) *Proof.* Note that $f'(t)$ is just a number, so we ignore the distinction between $Df(t)$

and $f'(t)$. By the chain rule, and using the result from part (a),

$$\begin{aligned}
 Df(t) &= D \left(\det \circ \left(\begin{bmatrix} a_{11} \\ \vdots \\ a_{n1} \end{bmatrix}, \dots, \begin{bmatrix} a_{1n} \\ \vdots \\ a_{nn} \end{bmatrix} \right) \right) (t) \\
 &= D(\det) \left(\begin{bmatrix} a_{11}(t) \\ \vdots \\ a_{n1}(t) \end{bmatrix}, \dots, \begin{bmatrix} a_{1n}(t) \\ \vdots \\ a_{nn}(t) \end{bmatrix} \right) \left(\begin{bmatrix} a'_{11}(t) \\ \vdots \\ a'_{n1}(t) \end{bmatrix}, \dots, \begin{bmatrix} a'_{1n}(t) \\ \vdots \\ a'_{nn}(t) \end{bmatrix} \right) \\
 &= \sum_{i=1}^n \det \begin{bmatrix} a_1 & \dots & a_i' & \dots & a_n \\ | & & | & & | \\ a_1(t) & \dots & a_i(t) & \dots & a_n(t) \\ | & & | & & | \end{bmatrix} \quad \square
 \end{aligned}$$

(c) For any fixed t , we essentially have the condition that

$$\begin{bmatrix} a_{11}(t) & \dots & a_{1n}(t) \\ \vdots & \ddots & \vdots \\ a_{n1}(t) & \dots & a_{nn}(t) \end{bmatrix} \begin{bmatrix} s_1(t) \\ \vdots \\ s_n(t) \end{bmatrix} = \begin{bmatrix} b_1(t) \\ \vdots \\ b_n(t) \end{bmatrix}$$

or more concisely, we can write

$$A(t)\vec{s}(t) = \vec{b}(t)$$

Since we are given that $\det A(t) \neq 0$, we know that $A(t)$ is invertible. Then by Cramer's Rule,

$$s_i(t) = \frac{\det(A_i(t))}{\det(A(t))}$$

where

$$A_i(t) = \begin{bmatrix} a_{11}(t) & \dots & b_1(t) & \dots & a_{1n}(t) \\ \vdots & & \vdots & & \vdots \\ a_{n1}(t) & \dots & b_n(t) & \dots & a_{nn}(t) \end{bmatrix}$$

Then $s_i(t)$ is differentiable as the quotient of differentiable functions. To calculate $s_i'(t)$, we have

$$s_i'(t) = \frac{\det(A(t))D(\det \circ A_i)(t) - \det(A_i(t))D(\det \circ A)(t)}{[\det(A(t))]^2}$$

Define the following matrices for convenience:

$$A^j(t) = \begin{bmatrix} | & & | & & | \\ a_1(t) & \dots & a'_j(t) & \dots & a_n(t) \\ | & & | & & | \end{bmatrix}$$

$$A_i^j(t) = \begin{cases} \begin{bmatrix} | & & | & & | & & | \\ a_1(t) & \dots & a'_j(t) & \dots & b_i(t) & \dots & a_n(t) \\ | & & | & & | & & | \end{bmatrix}, & i \neq j \\ \begin{bmatrix} | & & | & & | \\ a_1(t) & \dots & b'_i(t) & \dots & a_n(t) \\ | & & | & & | \end{bmatrix}, & i = j \end{cases}$$

Then the results from part (b), and the quotient rule,

$$s'_i(t) = \frac{\det(A(t)) \sum_{j=1}^n \det A_i^j(t) - \det(A_i(t)) \sum_{j=1}^n \det A^j(t)}{[\det(A(t))]^2}$$

Exercise 2-16 Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is differentiable and has a differentiable inverse $f^{-1} : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Show that

$$(f^{-1})'(a) = [f'(f^{-1}(a))]^{-1}$$

Proof. By definition,

$$f \circ f^{-1} = \text{Id}$$

Since both f and f^{-1} are differentiable, we can apply the chain rule in matrix form:

$$I_n = f'(f^{-1}(a)) \cdot (f^{-1})'(a)$$

Since both $f'(f^{-1}(a))$ and $(f^{-1})'(a)$ are $n \times n$ matrices, being single sided inverses is equivalent to being inverses, so we conclude that

$$(f^{-1})'(a) = [f'(f^{-1}(a))]^{-1} \quad \square$$

Exercise 2-17 Find the partial derivatives of the following functions:

(a) $f(x, y, z) = x^y$

(b) $f(x, y, z) = z$

(c) $f(x, y) = \sin(x \sin y)$

(d) $f(x, y, z) = \sin(x \sin(y \sin z))$

(e) $f(x, y, z) = x^{y^z}$

(f) $f(x, y, z) = x^{y+z}$

(g) $f(x, y, z) = (x + y)^z$

(h) $f(x, y) = \sin(xy)$

(i) $f(x, y) = [\sin(xy)]^{\cos 3}$

(a)

$$D_1 f(x, y, z) = yx^{y-1}$$

$$D_2 f(x, y, z) = x^y \ln x$$

$$D_3 f(x, y, z) = 0$$

(b)

$$D_1 f(x, y, z) = 0$$

$$D_2 f(x, y, z) = 0$$

$$D_3 f(x, y, z) = 1$$

(c)

$$D_1 f(x, y) = \sin y \cos(x \sin y)$$

$$D_2 f(x, y) = x \cos y \cos(x \sin y)$$

(d)

$$D_1 f(x, y, z) = \sin(y \sin z) \cos(x \sin(y \sin z))$$

$$D_2 f(x, y, z) = x \sin z \cos(y \sin z) \cos(x \sin(y \sin z))$$

$$D_3 f(x, y, z) = xy \cos z \cos(y \sin z) \cos(x \sin(y \sin z))$$

(e)

$$D_1 f(x, y, z) = y^z x^{y^z-1}$$

$$D_2 f(x, y, z) = zy^{z-1} x^{y^z} \ln x$$

$$D_3 f(x, y, z) = y^z x^{y^z} \ln x \ln y$$

(f)

$$\begin{aligned}D_1 f(x, y, z) &= (y + z)x^{y+z-1} \\D_2 f(x, y, z) &= x^z x^y \ln x^{y+z} \ln x \\D_3 f(x, y, z) &= x^{y+z} \ln x\end{aligned}$$

(g)

$$\begin{aligned}D_1 f(x, y, z) &= z(x + y)^{z-1} \\D_2 f(x, y, z) &= z(x + y)^{z-1} \\D_3 f(x, y, z) &= (x + y)^z \ln(x + y)\end{aligned}$$

(h)

$$\begin{aligned}D_1 f(x, y) &= y \cos(xy) \\D_2 f(x, y) &= y \cos(xy)\end{aligned}$$

(i)

$$\begin{aligned}D_1 f(x, y) &= y \cos 3[\sin(xy)]^{\cos 3-1} \cos(xy) \\D_2 f(x, y) &= x \cos 3[\sin(xy)]^{\cos 3-1} \cos(xy)\end{aligned}$$

Exercise 2-18 If $g : \mathbb{R} \rightarrow \mathbb{R}$ is continuous, find the partial derivatives of each of the following functions:

(a) $f(x, y) = \int_a^{x+y} g$

(b) $f(x, y) = \int_y^x g$

(c) $f(x, y) = \int_a^{xy} g$

(d) $f(x, y) = \int_a^{(\int_b^y g)} g$

(a) By the fundamental theorem of calculus,

$$\begin{aligned}D_1 f(x, y) &= g(x + y) \\D_2 f(x, y) &= g(x + y)\end{aligned}$$

(b) Let $a \in \mathbb{R}$. Then

$$\int_y^x g = \int_a^x g - \int_a^y g$$

so

$$\begin{aligned}D_1 f(x, y) &= g(x) \\D_2 f(x, y) &= -g(y)\end{aligned}$$

(c)

$$D_1 f(x, y) = yg(xy)$$

$$D_2 f(x, y) = xg(xy)$$

(d)

$$D_1 f(x, y) = 0$$

$$D_2 f(x, y) = g\left(\int_b^y g\right)g(y)$$

Exercise 2-19 If

$$f(x, y) = x^{x^{x^{x^y}}} + (\ln x)(\arctan(\arctan(\arctan(\sin(\cos xy) - \ln(x + y))))))$$

Find $D_2 f(1, y)$.

Since we are calculating D_2 , we treat x as constant, and in particular, we can substitute in $x = 1$. So we have

$$g_2(y) = f(1, y) = \underbrace{1^{1^{1^y}}}_{=1} + \underbrace{(\ln 1)}_{=0}(\arctan(\arctan(\arctan(\sin(\cos y) - \ln(1 + y))))))$$

So $g_2(y) = 1$ for all y , and thus $g_2'(y) = D_2 f(1, y) = 0$.

Exercise 2-20 Find the partial derivatives of f in terms of g, h, g', h' .

(a) $f(x, y) = g(x)h(y)$

(b) $f(x, y) = g(x)^{h(y)}$

(c) $f(x, y) = g(x)$

(d) $f(x, y) = g(y)$

(e) $f(x, y) = g(x + y)$

(a)

$$D_1 f(x, y) = h(y)g'(x)$$

$$D_2 f(x, y) = g(x)h'(y)$$

(b)

$$D_1 f(x, y) = h(y)g(x)^{h(y)-1}$$

$$D_2 f(x, y) = g(x)^{h(y)} \ln(g(x))$$

(c)

$$\begin{aligned}D_1f(x, y) &= g'(x) \\D_2f(x, y) &= 0\end{aligned}$$

(d)

$$\begin{aligned}D_1f(x, y) &= 0 \\D_2f(x, y) &= g'(y)\end{aligned}$$

(e)

$$\begin{aligned}D_1f(x, y) &= g'(x + y) \\D_2f(x, y) &= g'(x + y)\end{aligned}$$

Exercise 2-21 Let $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuous. Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$f(x, y) = \int_0^x g_1(t, 0)dt + \int_0^y g_2(x, t)dt$$

(a) Show that $D_2f(x, y) = g_2(x, y)$.

(b) How should f be defined such that $D_1f(x, y) = g_1(x, y)$?

(c) Find a function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ such that $D_1f(x, y) = x$ and $D_2f(x, y) = y$. Find one such that $D_1f(x, y) = y$ and $D_2f(x, y) = x$.

(a) *Proof.* Define

$$h_2(y) := f(x, y)$$

Then

$$D_2f(x, y) = h_2'(y) = \frac{d}{dy} \int_0^x g_1(t, 0)dt + \frac{d}{dy} \int_0^y g_2(x, t)dt$$

Since the first integral is constant with respect to y ,

$$\frac{d}{dy} \int_0^x g_1(t, 0)dt = 0$$

By the fundamental theorem of calculus,

$$\frac{d}{dy} \int_0^y g_2(x, t)dt = g_2(x, y)$$

Thus

$$D_2f(x, y) = h_2'(y) = g_2(x, y)$$

□

(b) Define

$$f(x, y) = \int_0^x g_1(t, y) dt + \int_0^y g_2(0, t) dt$$

Then by a similar argument as above, $D_1 f(x, y) = g_1(x, y)$.

(c) The function $f(x, y) = \frac{x^2}{2} + \frac{y^2}{2}$ satisfies

$$D_1 f(x, y) = x, D_2 f(x, y) = y$$

The function $f(x, y) = xy$ satisfies

$$D_1 f(x, y) = y, D_2 f(x, y) = x$$

Exercise 2-22 If $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $D_2 f = 0$, show that f is independent of the second variable. If $D_1 f = D_2 f = 0$, show that f is constant.

Proof. Fix some $x \in \mathbb{R}$, and define $h_x(y) = f(x, y)$. Since $D_2 f = 0$, $h'_x(y) = 0$ everywhere, so h_x is constant. Thus for any $y_1, y_2 \in \mathbb{R}$,

$$f(x, y_1) = h_x(y_1) = h_x(y_2) = f(x, y_2)$$

and thus f is independent of the second variable.

When $D_1 f = 0$, f is independent of the first variable as well. Moreover, we showed in Exercise 2-3 that functions which are independent of both variables are constant, so f is constant. \square

Exercise 2-23 Let $A = \{(x, y) \in \mathbb{R}^2 : x < 0, \text{ or } x \geq 0 \text{ and } y \neq 0\}$.

(a) If $f : A \rightarrow \mathbb{R}$ and $D_1 f = D_2 f = 0$, show that f is constant.

(b) Find a function $f : A \rightarrow \mathbb{R}$ such that $D_2 f = 0$ but f is not independent of the second variable.

Note: The set A as defined here is the plane excluding the nonnegative x -axis.

(a) *Proof.* Let $(x_1, y_1), (x_2, y_2) \in \mathbb{R}^2$ be arbitrary. Suppose $y_1 \neq 0$ and $y_2 \neq 0$. Define $g_x(y) = f(x, y)$ and $h_y(x) = f(x, y)$. Pick some $a < 0$. Then

$$\begin{aligned} f(x_1, y_1) - f(x_2, y_2) &= f(x_1, y_1) - f(a, y_1) + f(a, y_1) - f(a, y_2) + f(a, y_2) - f(x_2, y_2) \\ &= h_{y_1}(x_1) - h_{y_1}(a) + g_a(y_1) - g_a(y_2) + h_{y_2}(a) - h_{y_2}(x_2) \end{aligned}$$

Since $y_1 \neq 0$, h_{y_1} is defined on all of \mathbb{R} and h'_{y_1} is identically 0, h_{y_1} is constant. Similarly, h_{y_2} is constant, and g_a is also constant since $a < 0$. Thus

$$f(x_1, y_1) - f(x_2, y_2) = \underbrace{h_{y_1}(x_1) - h_{y_1}(a)}_{=0} + \underbrace{g_a(y_1) - g_a(y_2)}_{=0} + \underbrace{h_{y_2}(a) - h_{y_2}(x_2)}_{=0} = 0$$

The case where $y_1 = 0$ or $y_2 = 0$ is proved similarly. (Geometrically, we have connected the points (x_1, y_1) and (x_2, y_2) using three segments, but this can be adjusted to use only two or one if either y -coordinate is 0.) Thus $f(x_1, y_1) = f(x_2, y_2)$ for all points, and thus f is constant. \square

(b) Define $f : A \rightarrow \mathbb{R}$ by

$$f(x, y) = \begin{cases} 1, & x = 0, y > 0 \\ 0, & \text{otherwise} \end{cases}$$

Pick some point (x, y) . Then there exists an interval $(y - \varepsilon, y + \varepsilon) \subseteq A$. Moreover, f is constant on this interval. Thus $D_2 f(x, y) = 0$ everywhere, but f is not constant.

Exercise 2-24 Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$f(x, y) = \begin{cases} xy \frac{x^2 - y^2}{x^2 + y^2}, & (x, y) \neq 0 \\ 0, & (x, y) = 0 \end{cases}$$

- (a) Show that $D_2 f(x, 0) = x$ for all x and $D_1 f(0, y) = -y$ for all y .
 (b) Show that $D_{1,2} f(0, 0) \neq D_{2,1} f(0, 0)$.

(a) *Proof.* Define $g_x(y) = g(x, y)$ and $h_y(x) = f(x, y)$. Then

$$\begin{aligned} D_2 f(x, 0) &= g'_x(0) \\ &= \frac{d}{dy} \left(xy \frac{x^2 - y^2}{x^2 + y^2} \right) \Big|_{y=0} \\ &= \left(x \frac{x^2 - y^2}{x^2 + y^2} + xy \frac{-2x^2 y - 2y^3 - 2x^2 y + 2y^3}{(x^2 + y^2)^2} \right) \Big|_{y=0} \\ &= x \frac{x^2}{x^2} \end{aligned}$$

And

$$\begin{aligned} D_1 f(0, y) &= h'_y(0) \\ &= \frac{d}{dx} \left(xy \frac{x^2 - y^2}{x^2 + y^2} \right) \Big|_{x=0} \\ &= \left(y \frac{x^2 - y^2}{x^2 + y^2} + xy \frac{2x^3 + 2xy^2 - 2x^3 + 2xy^2}{(x^2 + y^2)^2} \right) \Big|_{x=0} \\ &= y \frac{-y^2}{y^2} \\ &= -y \end{aligned} \quad \square$$

(b) Taking the derivative of the functions we computed in part (a),

$$D_{1,2}f(0,0) = \frac{d}{dy}D_1f(0,y) = \frac{d}{dy}(-y) = -1$$

$$D_{2,1}f(0,0) = \frac{d}{dx}D_2f(x,0) = \frac{d}{dx}x = 1$$

so

$$D_{1,2}f(0,0) = -1 \neq 1 = D_{2,1}f(0,0)$$

Exercise 2-25 Define $f : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f(x) = \begin{cases} e^{-x^{-2}}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Show that f is C^∞ , and $f^{(i)}(0) = 0$ for all i .

Proof. For points $x \neq 0$, we have

$$f'(x) = \frac{2}{x^3}e^{-x^{-2}}$$

and

$$f''(x) = \frac{-6}{x^4}e^{-x^{-2}} + \frac{4}{x^6}e^{-x^{-2}}$$

Claim: In general, for any $i > 0$ and $x \neq 0$, $f^{(i)}(x)$ is composed of terms of the form

$$\frac{a}{x^b}e^{-x^{-2}}, \quad a \in \mathbb{Z}, b \in \mathbb{Z}_{\geq 0}$$

We prove this by induction. As shown, we already know this is true for $i = 1, 2$. Now suppose it is true for $i = k$. Then for $k + 1$, it is sufficient to show that each term of the above form differentiates into further terms of that form. Differentiating,

$$\frac{d}{dx} \frac{a}{x^b}e^{-x^{-2}} = \frac{-ab}{x^{b+1}}e^{-x^{-2}} + \frac{2a}{x^{b+3}}e^{-x^{-2}}$$

and the two terms are also of the form requested. Thus the claim is proved. This shows that $f^{(i)}(x)$ exists for all i when $x \neq 0$.

For $x = 0$, we use L'Hopital's rule:

$$\begin{aligned} f'(0) &= \lim_{h \rightarrow 0} \frac{e^{-h^{-2}}}{h} \\ &= \lim_{h \rightarrow 0} \frac{\frac{1}{h}}{e^{h^{-2}}} \\ (LH) &= \lim_{h \rightarrow 0} \frac{-\frac{1}{h^2}}{-2h^{-3}e^{h^{-2}}} \\ &= \lim_{h \rightarrow 0} \frac{h}{2e^{h^{-2}}} \\ &= 0 \end{aligned}$$

Similarly, for higher derivatives, we can apply the claim proved above to write

$$f^{(i)}(x) = \sum_{j=1}^n \frac{a_j}{x^{b_j}} e^{-x^{-2}}, \quad a_j \in \mathbb{Z}, b_j \in \mathbb{Z}_{\geq 0}$$

for some finite n . Then

$$\begin{aligned} f^{(i+1)}(0) &= \lim_{h \rightarrow 0} \frac{f^{(i)}(h)}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sum_{j=1}^n \frac{a_j}{h^{b_j}} e^{-h^{-2}}}{h} \\ &= \sum_{j=1}^n \left(\lim_{h \rightarrow 0} \frac{a_j}{h^{b_j+1}} e^{-h^{-2}} \right) \\ &= \sum_{j=1}^n \left(a_j \lim_{h \rightarrow 0} \frac{1}{h^{b_j+1} e^{h^{-2}}} \right) \\ (LH) &= \sum_{j=1}^n \left(a_j \lim_{h \rightarrow 0} \frac{-\frac{(b_j+1)}{h^{b_j+2}}}{\frac{-2}{h^3} e^{h^{-2}}} \right) \\ &= \sum_{j=1}^n \left(a_j \frac{b_j+1}{2} \lim_{h \rightarrow 0} \frac{e^{x^{-2}}}{h^{b_j-1}} \right) \\ &\quad \vdots \\ &= 0 \end{aligned}$$

Thus $f^{(i)}(x)$ exists for all i, x , so f is C^∞ , and $f^{(i)}(0) = 0$ for all i . □

Exercise 2-26 Let

$$f(x) = \begin{cases} e^{-(x-1)^{-2}} e^{-(x+1)^{-2}}, & x \in (-1, 1) \\ 0, & x \notin (-1, 1) \end{cases}$$

- (a) Show that $f : \mathbb{R} \rightarrow \mathbb{R}$ is a C^∞ function which is positive on $(-1, 1)$ and 0 elsewhere.
- (b) Show that there is a C^∞ function $s : \mathbb{R} \rightarrow [0, 1]$ such that $s(x) = 0$ for $x \leq 0$ and $s(x) = 1$ for $x \geq \varepsilon$.
- (c) If $a = (a_1, \dots, a_n) \in \mathbb{R}^n$, define $g_a : \mathbb{R}^n \rightarrow \mathbb{R}$ by

$$g_a(x) = g_a(x_1, \dots, x_n) = f\left(\frac{x_1 - a_1}{\varepsilon}\right) \cdots f\left(\frac{x_n - a_n}{\varepsilon}\right)$$

Show that g_a is a C^∞ function which is positive on

$$(a_1 - \varepsilon, a_1 + \varepsilon) \times \dots \times (a_n - \varepsilon, a_n + \varepsilon)$$

- (d) If $A \subseteq \mathbb{R}^n$ is open and $C \subseteq A$ is compact, show that there is a nonnegative C^∞ function $h : A \rightarrow \mathbb{R}$ such that $f(x) > 0$ for $x \in C$ and $f = 0$ outside of some closed set contained in A .
- (e) Show that we can choose such an h so that $h : A \rightarrow [0, 1]$ and $h(x) = 1$ for $x \in C$.

- (a) *Proof.* By definition, f is 0 outside of $(-1, 1)$, and it must be positive on $(-1, 1)$ since each of the exponential factors are positive.

To show that f is C^∞ , define $f_1, f_2 : (-1, 1) \rightarrow \mathbb{R}$ by

$$\begin{aligned} f_1(x) &= e^{-(x-1)^{-2}} \\ f_2(x) &= e^{-(x+1)^{-2}} \end{aligned}$$

We proved in Exercise 2-25 that both f_1, f_2 are C^∞ , so

$$f'(x) = f_1(x)f_2'(x) + f_1'(x)f_2(x)$$

and higher order derivatives will in general be sums of products of $f_1^{(i)}(x)$ and $f_2^{(j)}(x)$, which all exist and are continuous. Thus f is C^∞ . \square

- (b) *Proof.* Fix $\varepsilon > 0$. Then define

$$s(x) = \begin{cases} 1, & x \geq 2\varepsilon \\ \frac{f(1-\frac{x}{\varepsilon})}{f(\frac{x}{\varepsilon}) + f(1-\frac{x}{\varepsilon})}, & -\varepsilon < x < 2\varepsilon \\ 0, & x \leq -\varepsilon \end{cases}$$

By definition, $s(x) = 0$ for $x \leq -\varepsilon$ and $s(x) = 1$ for $x \geq 2\varepsilon$.

On the interval $(-\varepsilon, 0]$, $1 - \frac{x}{\varepsilon} \geq 1$, so $f(1 - \frac{x}{\varepsilon}) = 0$ and thus $s = 0$. So $s = 0$ for any $x \leq 0$.

Similarly, for the interval $[\varepsilon, 2\varepsilon)$, $\frac{x}{\varepsilon} \geq 1$, so $f(\frac{x}{\varepsilon}) = 0$ and

$$s(x) = \frac{f(1 - \frac{x}{\varepsilon})}{f(1 - \frac{x}{\varepsilon})} = 1$$

So $s = 1$ for any $x \geq \varepsilon$.

To prove that s is C^∞ , we can obviously ignore the constant regions.

On $(0, \varepsilon)$, at least one of $f(\frac{x}{\varepsilon})$, $f(1 - \frac{x}{\varepsilon})$ will be positive, so the quotient rule says that $s'(x)$ exists. In general, we can continue to apply the quotient rule, since the quotient will never be zero, and f is smooth. Thus $s^{(i)}(x)$ exists and is continuous for all i and $x \in (0, \varepsilon)$, and we conclude that s is C^∞ . \square

(c) *Proof.* The fact that g_a is positive follows from the fact that for each i ,

$$\left| \frac{x_i - a_i}{\varepsilon} \right| < 1$$

so $f(\frac{x_i - a_i}{\varepsilon}) > 0$. Thus their product g_a is positive.

To show that g_a is C^∞ , we need to prove that the mixed partials of all orders exist. Here, we can actually prove a more general result:

Lemma

If $f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R}$ are C^∞ , then $f : \mathbb{R}^n \rightarrow \mathbb{R}$ defined by

$$f(x_1, \dots, x_n) := f_1(x_1) \cdot \dots \cdot f_n(x_n)$$

is C^∞ .

Proof. To prove that derivatives of all orders exist and are continuous, pick any index i . Then define

$$g_i(x_i) = f_i(x_i) \left(\prod_{j \neq i} f_j(x_j) \right)$$

Then $g_i(x_i)$ is just a constant multiple of $f_i(x_i)$, so $g_i'(x_i)$ exists. Moreover, f_i' is also C^∞ , so the function

$$D_i f(x_1, \dots, x_n) = f_1(x_1) \dots f_i'(x_i) \dots f_n(x_n)$$

satisfies the hypotheses of this lemma and we can differentiate it again using the above method. So derivatives of all orders exist and are continuous. Thus f is C^∞ . \square

We can then apply the above lemma to conclude that g_a is C^∞ . \square

(d) *Proof.* For each $x = (x_1, \dots, x_n) \in C$, there exists ε_x such that the rectangle

$$R_x = (x_1 - \varepsilon_x, x_1 + \varepsilon_x) \times \dots \times (x_n - \varepsilon_x, x_n + \varepsilon_x) \subseteq A$$

In fact, we may choose ε_x small enough such that the closed rectangle is contained in A as well. Let \mathcal{O} be the collection of R_x for $x \in C$. Since C is compact, we pick a finite subcover $\mathcal{O}' = \{R_{x_i}\}_{i=1}^m$. Then define $h : A \rightarrow \mathbb{R}$ by

$$h(x_1, \dots, x_n) = \sum_{i=1}^m g_{x_i}(x_1, \dots, x_n)$$

h is C^∞ since it is the product of C^∞ functions (by the lemma in part (d)). For any $y \in C$, \mathcal{O}' covers C , so $y \in R_{x_i}$ for some x_i . Then $g_{x_i} > 0$, and each other g_{x_j} is at least nonnegative, so $h(y) > 0$.

Now let \bar{R}_{x_i} be the closed rectangle about x_i .

$$B = \bigcup_{i=1}^m \bar{R}_{x_i}$$

We showed that we can pick ε small enough that $\bar{R}_{x_i} \subseteq A$. Thus B is a closed set contained in A . Moreover, if $y \notin B$, then $y \notin R_{x_i}$ for any i , and hence $h(y) = 0$. So h is 0 on outside of a closed set contained in A . \square

(e) *Proof.* Since h is C^∞ , it is continuous, and hence achieves a minimum value on C . Since h is positive on C , this minimum value $\varepsilon = \min_{x \in C} h(x)$ is positive. Let $s_\varepsilon : \mathbb{R} \rightarrow [0, 1]$ be as defined in part (b). Then the function

$$s_\varepsilon \circ h : \mathbb{R}^n \rightarrow [0, 1]$$

is still C^∞ (since the composition of C^∞ functions is C^∞ using repeated applications of the chain rule, similarly to the lemma in part (c)). Letting B be as defined previously, if $y \notin B$ then $h(y) = 0$, so $s_\varepsilon(h(y)) = s_\varepsilon(0) = 0$. Thus $s_\varepsilon \circ h$ is still of the form in part (d).

Moreover, whenever $x \in C$, $h(x) \geq \varepsilon$ so $s_\varepsilon(h(x)) = 1$. \square

Exercise 2-27 Define $g, h : \{x \in \mathbb{R}^2 : |x| \leq 1\} \rightarrow \mathbb{R}^3$ by

$$\begin{aligned} g(x, y) &= (x, y, \sqrt{1 - x^2 - y^2}) \\ h(x, y) &= (x, y, -\sqrt{1 - x^2 - y^2}) \end{aligned}$$

Let $f : \{x \in \mathbb{R}^3 : |x| = 1\} : \mathbb{R}$. Show that the maximum of f is either the maximum of $f \circ g$ or the maximum of $f \circ h$ on $\{x \in \mathbb{R}^2 : |x| \leq 1\}$.

Proof. Let $D_2 = \{x \in \mathbb{R}^2 : |x| \leq 1\}$ and $C_3 = \{x \in \mathbb{R}^3 : |x| = 1\}$. Then supposing that f has a maximum $m = \max_{x \in D_2} f(x)$, then there exists at least one point $x = (x_1, x_2, x_3)$ such that $f(x) = m$. Then we have the cases $x_3 \geq 0$ and $x_3 < 0$.

Case 1: Since $|x| = 1$, $x_1^2 + x_2^2 + x_3^2 = 1$, and hence

$$x_3 = \sqrt{1 - x_1^2 - x_2^2}$$

Thus we have $g(x_1, x_2) = x$, so $(f \circ g)(x_1, x_2) = m$. $(f \circ g)$ certainly cannot achieve a higher value, or else it would contradict m being the maximum of f , so m is also the maximum of g .

Case 2: Similar to Case 1, but we use $h(x_1, x_2)$ instead, and we find that $(f \circ h)$ achieves the maximum m .

Thus we see that m is the maximum of at least one of $f \circ g$ or $f \circ h$ on D_2 . □

Exercise 2-28 Find expressions for the partial derivatives of the following functions:

(a) $F(x, y) = f(g(x)k(y), g(x) + h(y))$

(b) $F(x, y, z) = f(g(x + y), h(y + z))$

(c) $F(x, y, z) = f(x^y, y^z, z^x)$

(d) $F(x, y) = f(x, g(x), h(x, y))$

(a) Let $f(*) = f(g(x)k(y), g(x) + h(y))$. Using the chain rule for partial derivatives,

$$\begin{aligned} D_1F(x, y) &= D_1f(*)D_x[g(x)k(y)] + D_2f(*)D_x[g(x) + h(y)] \\ &= k(y)g'(x)D_1f(*) + g'(y)D_2f(*) \\ D_2F(x, y) &= D_1f(*)D_y[g(x)k(y)] + D_2f(*)D_y[g(x) + h(y)] \\ &= g(x)k'(y)D_1f(*) + h'(y)D_2f(*) \end{aligned}$$

(b) Let $f(*) = f(g(x + y), h(y + z))$. Then

$$\begin{aligned} D_1F(x, y, z) &= D_1f(*)D_xg(x + y) + D_2f(*)D_xh(y + z) \\ &= g'(x + y)D_1f(*) \\ D_2F(x, y, z) &= D_1f(*)D_yg(x + y) + D_2f(*)D_yh(y + z) \\ &= g'(x + y)D_1f(*) + h'(y + z)D_2f(*) \\ D_3F(x, y, z) &= D_1f(*)D_zg(x + y) + D_2f(*)D_zh(y + z) \\ &= h'(y + z)D_2f(*) \end{aligned}$$

(c) Let $f(*) = f(x^y, y^z, z^x)$. Omitting zero terms,

$$\begin{aligned} D_1F(x, y, z) &= yx^{y-1}D_1f(*) + z^x \ln z D_3f(*) \\ D_2F(x, y, z) &= x^y \ln x D_1f(*) + zy^{z-1}D_2f(*) \\ D_3F(x, y, z) &= y^z \ln y D_2f(*) + xz^{x-1}D_3f(*) \end{aligned}$$

(c) Let $f(*) = f(x, g(x), h(x, y))$. Then

$$\begin{aligned} D_1F(x, y) &= D_1f(*) + g'(x)D_2f(*) + D_1h(x, y)D_3f(*) \\ D_2F(x, y) &= D_2h(x, y)D_3f(*) \end{aligned}$$

Exercise 2-29 Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$. For $\vec{x} \in \mathbb{R}^n$, if the limit

$$\lim_{t \rightarrow 0} \frac{f(a + t\vec{x}) - f(a)}{t}$$

exists, it is called the **directional derivative** of f at a in the direction \vec{x} , denoted $D_{\vec{x}}f(a)$.

(a) Show that $D_{e_i}f(a) = D_i f(a)$.

(b) Show that $D_{t\vec{x}}f(a) = tD_{\vec{x}}f(a)$.

(c) If f is differentiable at a , show that $D_{\vec{x}}f(a) = Df(a)(\vec{x})$ and therefore $D_{\vec{x}+\vec{y}}f(a) = D_{\vec{x}}f(a) + D_{\vec{y}}f(a)$.

(a) *Proof.* Immediate from the definitions. □

(b) *Proof.* Fix $t \in \mathbb{R}$. Then

$$\begin{aligned} \lim_{s \rightarrow 0} \frac{f(a + s(t\vec{x})) - f(a)}{s} &= t \lim_{s \rightarrow 0} \frac{f(a + st\vec{x}) - f(a)}{st} \\ &= t \lim_{st \rightarrow 0} \frac{f(a + (st)\vec{x}) - f(a)}{(st)} \\ &= tD_{\vec{x}}f(a) \end{aligned} \quad \square$$

(c) *Proof.* Since the derivative exists, we know that

$$\lim_{t\vec{x} \rightarrow 0} \frac{f(a + t\vec{x}) - f(a) - Df(a)(t\vec{x})}{t|\vec{x}|} = 0$$

We can multiply both sides by $|\vec{x}|$ to clear the denominator, and apply linearity of $Df(a)$ to see that

$$\begin{aligned} \lim_{t \rightarrow 0} \frac{f(a + t\vec{x}) - f(a) - tDf(a)(\vec{x})}{t} &= 0 \\ \implies \lim_{t \rightarrow 0} \frac{f(a + t\vec{x}) - f(a)}{t} &= Df(a)(\vec{x}) \end{aligned}$$

and thus $Df(a)(\vec{x}) = D_{\vec{x}}f(a)$. Since $Df(a)$ is linear,

$$\begin{aligned} D_{\vec{x}+\vec{y}}f(a) &= Df(a)(\vec{x} + \vec{y}) \\ &= Df(a)(\vec{x}) + Df(a)(\vec{y}) \\ &= D_{\vec{x}}f(a) + D_{\vec{y}}f(a) \end{aligned} \quad \square$$

Exercise 2-30 Let f be defined as in Exercise 2-4. Show that $D_{\vec{x}}f(0, 0)$ exists for all x , but if $g \neq 0$, then $D_{\vec{x}+\vec{y}}f(0, 0) = D_{\vec{x}}f(0, 0) + D_{\vec{y}}f(0, 0)$ is not true for all x, y .

Proof. The result of Exercise 2-4 part (a) says that for $x \in \mathbb{R}^2$, defining $h_x(t) = f(tx)$ means that h_x is differentiable at $(0, 0)$. This means that $D_{\vec{x}}f(0, 0)$ exists for all \vec{x} . Similarly, as the result in part (b) shows, $D_{e_1}f(0) = D_{e_2}f(0) = 0$. However, if g is nonzero, then we can take a directional derivative in some direction which is a linear combination of e_1 and e_2 , so the linearity condition fails. \square

Exercise 2-31 Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined as in Exercise 1-26. Show that $D_x f(0, 0)$ exists for all x , even though f is not continuous at $(0, 0)$.

Proof. As we showed in the proof of Exercise 1-26 part (b), f is 0 in an interval about $(0, 0)$ in each direction, and is thus differentiable. \square

Exercise 2-32

(a) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

Show that f is differentiable at 0 but f' is not continuous at 0.

(b) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} (x^2 + y^2) \sin \frac{1}{\sqrt{x^2 + y^2}}, & (x, y) \neq 0 \\ 0, & (x, y) = 0 \end{cases}$$

Show that f is differentiable at $(0, 0)$ but $D_i f$ is not continuous at $(0, 0)$.

(a) *Proof.* Let $\varepsilon > 0$. Then whenever $|x - 0| < \delta = \varepsilon$, we have

$$\left| \frac{f(x) - f(0)}{x} - 0 \right| = \left| \frac{f(x)}{x} \right| = \left| x \sin \frac{1}{x} \right| < \varepsilon$$

Thus f is differentiable at 0 with $f'(0) = 0$.

If we differentiate f elsewhere, we find that

$$f'(x) = \begin{cases} 2x \sin \frac{1}{x} - \cos \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

But

$$\lim_{x \rightarrow 0} 2x \sin \frac{1}{x} - \cos \frac{1}{x} = - \lim_{x \rightarrow 0} \cos \frac{1}{x}$$

which doesn't exist. Thus f' is not continuous at 0 (it has an oscillating discontinuity). \square

(b) *Proof.* Let $\varepsilon > 0$. Then whenever $|(x, y)| = \sqrt{x^2 + y^2} < \delta = \varepsilon$, we have

$$\left| \frac{f(x, y) - f(0, 0)}{|(x, y)|} \right| = \left| \frac{f(x, y)}{\sqrt{x^2 + y^2}} \right| = \left| \sqrt{x^2 + y^2} \sin \frac{1}{\sqrt{x^2 + y^2}} \right| \leq \sqrt{x^2 + y^2} < \varepsilon$$

Thus

$$\lim_{(x, y) \rightarrow (0, 0)} \frac{|f(x, y) - f(0, 0) - 0(x, y)|}{|(x, y)|} = 0$$

so $Df(0, 0)$ exists and is the zero transformation. But in the directions e_1, e_2 , f is simply the single variable case considered in part (a), so we know $D_i f$ is not continuous at $(0, 0)$. \square

Exercise 2-33 Show that the continuity of $D_1 f^j$ at a may be eliminated from the hypothesis of Theorem 2-8.

Proof. In the proof of Theorem 2-8, we attempted to prove that

$$\lim_{\vec{h} \rightarrow 0} \frac{|f(\vec{a} + [\vec{h}]^j) - f(\vec{a} + [\vec{h}]^{j-1}) - D_j f(\vec{a})h_j|}{|\vec{h}|} = 0$$

for all j . We did this by using the continuity of $D_j f$ at a to extend its differentiability nearby. However, in the case of the first partial derivative $D_1 f$, the continuous differentiability condition already shows us that

$$\lim_{\vec{h} \rightarrow 0} \frac{|f(\vec{a} + h_1 e_1) - f(\vec{a}) - D_1 f(\vec{a})h_1|}{|\vec{h}|} = 0$$

so we can omit continuity. (Obviously, any other direction would also work.) \square

Exercise 2-34 A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is **homogeneous** of degree m if $f(tx) = t^m f(x)$ for all x . If f is also differentiable, show that

$$\sum_{i=1}^n x^i D_i f(x) = m f(x)$$

Proof. Define $g(t) = f(tx)$. Then $D_x f(x) = g'(1)$. Moreover, we showed in Exercise 2-30 that D_* is linear, so

$$D_x f(x) = \sum_{i=1}^n x_i D_i f(x)$$

At the same time, we know that $g(t) = f(tx) = t^m f(x)$. Differentiating with respect to t ,

$$g'(x) = mt^{m-1} f(x)$$

so

$$\sum_{i=1}^n x_i D_i f(x) = D_x f(x) = g'(1) = mf(x)$$

□

Exercise 2-35 If $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is differentiable and $f(0) = 0$, prove that there exist $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

$$f(x) = \sum_{i=1}^n x_i g_i(x)$$

Proof. Since f is differentiable, the directional derivative $D_x f(tx)$ exists for all t, x . Define $h_x(t) = f(tx)$. Then $h'_x(t) = D_x f(tx)$. Thus h_x is differentiable. Then by the fundamental theorem of calculus,

$$f(x) = f(1x) = \int_0^1 h'_x(t) dt = \int_0^1 D_x f(tx) dt$$

Since D_* is linear with respect to direction, we then have

$$f(x) = \int_0^1 \sum_{i=1}^n x_i D_i f(tx) dt = \sum_{i=1}^n x_i \int_0^1 D_i f(tx) dt$$

Then defining $g_i(x) = \int_0^1 D_i f(tx) dt$, we have found g_i satisfying

$$f(x) = \sum_{i=1}^n x_i g_i(x)$$

□

Exercise 2-36 Let $A \subseteq \mathbb{R}^n$ be an open set and $f : A \rightarrow \mathbb{R}^n$ a continuously differentiable one-to-one function such that $\det f'(x) \neq 0$ for all x . Show that $f(A)$ is an open set and $f^{-1} : f(A) \rightarrow A$ is differentiable. Show also that $f(B)$ is open for any open set $B \subseteq A$.

Proof. Let $y \in f(A)$. Then since f is one-to-one, there exists a unique $x \in A$ such that $f(x) = y$. Since f is continuously differentiable at x and $\det f'(x) \neq 0$, the Inverse Function Theorem tells us there exist open sets $V \subseteq A$ containing x and $W \subseteq \mathbb{R}^n$ such that $f : V \rightarrow W$

has an inverse. Thus $W \subseteq f(A)$ and $y \in W$, so $f(A)$ is open. Moreover, the Inverse Function Theorem also says f^{-1} is differentiable at y . But this is true for every $y \in f(A)$, so f^{-1} is differentiable. Lastly, let $B \subseteq A$ be open. Then the restriction $\bar{f} : B \rightarrow \mathbb{R}^n$ is also continuously differentiable and one-to-one, so $f(B)$ is open. \square

Exercise 2-37

- (a) Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a continuously differentiable function. Show that f is **not** one-to-one.
- (b) Generalize this result to the case of a continuously differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ with $m < n$.

- (a) *Proof.* If $D_1f(x, y) = 0$ for all $(x, y) \in \mathbb{R}^2$, then f is independent of the first variable and is not one-to-one. So suppose there exists some $(x_1, y_1) \in \mathbb{R}^2$ with $D_1f(x_1, y_1) \neq 0$. Since f is continuously differentiable, there exists an open set A containing (x_1, y_1) such that $D_1f(x, y) \neq 0$ for any $(x, y) \in A$. Then define $g : A \rightarrow \mathbb{R}^2$ by $g(x, y) = (f(x, y), y)$. Then the derivative is given by

$$g'(x, y) = \begin{bmatrix} D_1f(x, y) & D_2f(x, y) \\ 0 & 1 \end{bmatrix} \implies \det g'(x, y) = D_1f(x, y) \neq 0$$

In particular, $\det g'(x_1, y_1) \neq 0$. Then by the Inverse Function Theorem, there exists an open set V containing (x_1, y_1) and an open set W containing $(f(x_1, y_1), y_1)$ such that $g : V \rightarrow W$ has a continuous, differentiable inverse $g^{-1} : W \rightarrow V$. Then pick some $y_2 \neq y_1$ such that $(f(x_1, y_1), y_2) \in W$. Then we have

$$g(g^{-1}(f(x_1, y_1), y_2))) = (f(x_1, y_1), y_2)$$

but by definition,

$$g(g^{-1}(f(x_1, y_1), y_2))) = (f(g^{-1}(f(x_1, y_1), y_2)), g_2^{-1}(f(x_1, y_1), y_2)))$$

So

$$f(x_1, y_1) = f(g^{-1}(f(x_1, y_1), y_2)))$$

While the x coordinate of $g^{-1}(f(x_1, y_1), y_2)$ is unknown, the y coordinate is certainly y_2 . Thus we have

$$f(x_1, y_1) = f(*, y_2)$$

But we mandated that $y_1 \neq y_2$, so $(x_1, y_1) \neq (*, y_2)$. So f is not one-to-one. \square

- (b) *Proof.* \square

Exercise 2-38

- (a) If $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies $f'(a) \neq 0$ for all $a \in \mathbb{R}$, show that f is one-to-one (on all of \mathbb{R}).
- (b) Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ by $f(x, y) = (e^x \cos y, e^x \sin y)$. Show that $\det f'(x, y) \neq 0$ for all (x, y) but f is not one-to-one.

(a) *Proof.* Suppose without loss of generality that $f'(a) > 0$ for some $a \in \mathbb{R}$. One can prove in single variable analysis that if $g = f'$ for some function f , then g satisfies the intermediate value property. If $f'(b) < 0$ for some $b \in \mathbb{R}$, then there exists c between a and b such that $f'(c) = 0$, contradicting the assumption. So we must have $f'(x) > 0$ for all x . Thus f is strictly increasing (or decreasing), so it is one-to-one. \square

(b) *Proof.* The Jacobian matrix is given by

$$f'(x, y) = \begin{bmatrix} e^x \cos y & e^x \sin y \\ -e^x \sin y & e^x \cos y \end{bmatrix}$$

so

$$\det f'(x, y) = e^x(\sin^2 y + \cos^2 y) = e^x \neq 0$$

But for any (x, y) , we have

$$f(x, y) = f(x, y + 2\pi)$$

so f is not one-to-one. \square

Exercise 2-39 Use the function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} \frac{x}{2} + x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$$

To show that continuity of the derivative cannot be eliminated from the hypothesis of the Inverse Function Theorem.

First, we verify that f is differentiable at 0. We have

$$\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{1}{2} + h \sin \frac{1}{h} = \frac{1}{2} + \lim_{h \rightarrow 0} h \sin \frac{1}{h} = \frac{1}{2}$$

and by the formula f is clearly differentiable everywhere else. So f is differentiable in an open set around 0. However, I claim that for any open set V around 0, f is **not** injective onto $f(V)$.

To see this, let V be an open set around 0. Then pick n large enough that

$$a = \frac{1}{2\pi n} \in V$$

Now, we have

$$f'(a) = \frac{1}{2} + 2a \sin \frac{1}{a} - \cos \frac{1}{a} = \frac{1}{2} - 1 = -\frac{1}{2} < 0$$

Thus there exists $b < a$ with $f(b) > f(a)$ and $b > 0$. Now, pick m large enough that

$$c = \frac{1}{2\pi m} < b$$

Then we have

$$f(c) = \frac{c}{2} < \frac{a}{2} = f(a)$$

So $f(c) < f(a) < f(b)$, and $b \in [c, a]$. Pick some y with $f(a) < y < f(b)$. By the Intermediate Value Theorem, there exists $x_1 \in (c, b)$ with $f(x_1) = y$, and $x_2 \in (b, a)$ with $f(x_2) = y$, so f is not one-to-one onto $f(V)$. Thus the Inverse Function Theorem is false for f .

Exercise 2-40 Use the implicit function theorem to redo Problem 2-15 (c). For reference, this problem is reprinted here:

If $a_{ij} : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable, let $A(t)$ be the matrix such that $A(t)_{ij} = a_{ij}(t)$. If $\det(A(t)) \neq 0$ for all t and $b_1, \dots, b_n : \mathbb{R} \rightarrow \mathbb{R}$ are differentiable, let $s_1, \dots, s_n : \mathbb{R} \rightarrow \mathbb{R}$ be the functions such that $s_1(t), \dots, s_n(t)$ are the solutions of the equations

$$\sum_{j=1}^n a_{ji}(t)s_j(t) = b_i(t)$$

Show that s_i is differentiable and find $s'_i(t)$.

Proof. Define $F : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that the component functions are given by

$$F^i(t, x) = -b_i(t) + \sum_{j=1}^n a_{ji}(t)x_j$$

Then F^i can alternately be written as

$$F^i = -b_i \circ (\pi^2, \dots, \pi^n) + \sum_{j=1}^n (a_{ji} \circ \pi^1)\pi^j$$

which makes it clear that it can be written as sums, products, and compositions of differentiable functions. If we assume that the a_{ij} and b_i are additionally continuously differentiable, then F is also continuously differentiable.

Now, fix t_1 . Let $M(t, x)$ be the matrix with ij th entry given by $D_{j+1}F^i(t, x)$. To calculate the matrix of partial derivatives, for $k \geq 2$ we have

$$\begin{aligned} D_k F^i(t_1, x) &= D_k \left(\sum_{j=1}^n a_{ji}(t_1)x_j \right) \\ &= \sum_{j=1}^n a_{ji}(t_1)e_j \delta_{ik} \\ &= a_{ki}(t_1)e_k \end{aligned}$$

Thus $M(t_1, x)$ is simply the matrix $[A(t_1)]^T$, where $A(t_1)$ has ij -th entry given by $a_{ij}(t_1)$. By assumption, $\det[A(t_1)]^T = \det A(t_1) \neq 0$, so $\det M(t_1, x) \neq 0$ and the Implicit Function Theorem applies. Then there exists an open set $A \subseteq \mathbb{R}$ containing t and a function $g : A \rightarrow \mathbb{R}^n$ such that

$$F(t, g(t)) = 0$$

But this happens precisely when each component function is zero, so for each component we have

$$-b_i(t) + \sum_{j=1}^n a_{ji}(t)g^j(t) = 0 \iff \sum_{j=1}^n a_{ji}(t)g^j(t) = b_i(t)$$

Thus we may let $s_j = g^j$. Since $\det A(t) \neq 0$ for all t we are able to "patch" the local definitions of g^j into a global function without issue. Moreover, the Implicit Function Theorem tells us that g is differentiable at t_1 , so each s_j is everywhere.

To calculate s'_i , we know that $F^i(t, \vec{s}(t)) = 0$. Taking partial derivatives on both sides, we have

$$\begin{aligned} D_1 F^i(t, \vec{s}(t)) &= 0 \\ D_2 F^i(t, \vec{s}(t))s'_1(t) &= 0 \\ D_3 F^i(t, \vec{s}(t))s'_2(t) &= 0 \\ &\vdots \\ D_{n+1} F^i(t, \vec{s}(t))s'_n(t) &= 0 \end{aligned}$$

which we can combine as

$$D_1 F^i(t, \vec{s}(t)) + \sum_{j=1}^n D_{j+1} F^i(t, \vec{s}(t))s'_j(t) = 0$$

Consider the system of equations this forms. We can rewrite it in matrix-vector multiplication using our definition of $M(t, x)$ from above as

$$M(t, \vec{s}(t))s'(t) = -(D_1 F^i(t, \vec{s}(t)))$$

Moreover, the i th coordinate of the vector $(D_1 F^i(t, \vec{s}(t)))$ is given by

$$-b'_i(t) + \sum_{j=1}^n a'_{ji}(t)s_j(t)$$

Since $M(t, \vec{s}(t))$ is invertible by assumption, we find that

$$s'(t) = [M(t, \vec{s}(t))]^{-1} \begin{bmatrix} b'_1(t) - \sum_{j=1}^n a'_{j1}(t)s_j(t) \\ \vdots \\ b'_n(t) - \sum_{j=1}^n a'_{jn}(t)s_j(t) \end{bmatrix} \quad \square$$

Exercise 2-41 Let $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ be differentiable. For each $x \in \mathbb{R}$ define $g_x : \mathbb{R} \rightarrow \mathbb{R}$ by $g_x(y) = f(x, y)$. Suppose that for each x there is a unique y with $g'_x(y) = 0$. Then let $c(x)$ be this y .

(a) If $D_{2,2}f(x, y) \neq 0$ for all (x, y) , show that c is differentiable and

$$c'(x) = -\frac{D_{2,1}f(x, c(x))}{D_{2,2}f(x, c(x))}$$

(b) Show that if $c'(x) = 0$, then for some y we have

$$\begin{aligned} D_{2,1}f(x, y) &= 0 \\ D_{2,2}f(x, y) &= 0 \end{aligned}$$

(c) Let $f(x, y) = x(y \ln y - y) - y \ln x$. Find

$$\max_{\frac{1}{2} \leq x \leq 2} \left(\min_{\frac{1}{3} \leq y \leq 1} f(x, y) \right)$$

Note: Spivak does not include this, but we must assume that f is twice continuously differentiable.

(a) *Proof.* Note that by our definition, $D_2f(x, y) = g'_x(y)$. So $y = c(x)$ precisely when $D_2f(x, y) = 0$. Note that D_2f is a function $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, and the matrix $M = (D_{j+1}(D_2f)^i(x, y))$ is simply the matrix with sole entry $D_{2,2}f(x, y)$. By assumption, $D_{2,2}f(x, y) \neq 0$, so $\det M \neq 0$ and the Implicit Function Theorem applies to D_2f , and we conclude that c is differentiable.

Now, the function $x \mapsto D_2f(x, c(x))$ is a function $\mathbb{R} \rightarrow \mathbb{R}$ and is 0 everywhere, so we can differentiate it:

$$D_{2,1}f(x, c(x)) + D_{2,2}f(x, c(x))c'(x) = 0$$

which we can rearrange as

$$c'(x) = -\frac{D_{2,1}f(x, c(x))}{D_{2,2}f(x, c(x))}$$

□

(b) *Proof.* Pick $y = c(x)$. Then by definition, $g'_x(c(x)) = 0$, and $D_2f(x, c(x)) = g'_x(c(x))$, so $D_2f(x, c(x)) = 0$. Moreover, from part (a),

$$D_{2,1}f(x, c(x)) = -c'(x)D_{2,2}f(x, c(x)) = 0$$

so this choice of y works. □

(c) For any fixed x ,

$$\min_{\frac{1}{3} \leq y \leq 1} f(x, y) = \min_{\frac{1}{3} \leq y \leq 1} g_x(y)$$

We already know that $g'_x(c(x)) = 0$, so it is a critical point. If we calculate $g''_x(y) = D_{2,2}f(x, y)$ for any y , we get

$$\begin{aligned} D_2f(x, y) &= x(\ln y + 1 - 1) - \ln x = x \ln y - \ln x \\ D_{2,2}f(x, y) &= \frac{x}{y} \end{aligned}$$

which is strictly positive (as both x, y must be positive for this function to be defined). Thus g_x is concave upward, and the critical point at $c(x)$ is in fact a global minimum.¹ So if $c(x) \in [\frac{1}{3}, 1]$, then the minimum is at $c(x)$. If $c(x) < \frac{1}{3}$, then the minimum is at $\frac{1}{3}$, and if $c(x) > 1$, then the minimum is at 1.

If we explicitly calculate $c(x)$, we use the fact that $D_2f(x, c(x)) = 0$ to find

$$\ln c(x) = \frac{\ln x}{x} \implies c(x) = e^{\frac{\ln x}{x}} = \sqrt[x]{x}$$

and the derivative of this is positive, so c is strictly increasing. Thus there exists a unique α with $c(\alpha) = \frac{1}{3}$, and $x < \alpha \implies c(x) < \frac{1}{3}$. Similarly, $x > 1 \implies c(x) > 1$. So we can explicitly find the minimum of g_x :

$$\begin{aligned} \min_{\frac{1}{3} \leq y \leq 1} g_x(y) &= \begin{cases} f(x, \frac{1}{3}), & x < \alpha \\ f(x, c(x)), & \alpha \leq x \leq 1 \\ f(x, 1), & x > 1 \end{cases} \\ &= \begin{cases} x(\frac{\ln \frac{1}{3}}{3} - \frac{1}{3}) - \frac{\ln x}{3}, & x < \alpha \\ x(\sqrt[x]{x} \frac{\ln x}{x} - \sqrt[x]{x}) - \sqrt[x]{x} \ln x, & \alpha \leq x \leq 1 \\ -x - \ln x, & x > 1 \end{cases} \\ &= \begin{cases} \frac{-x \ln 3 - x - \ln x}{3}, & x < \alpha \\ -x \sqrt[x]{x}, & \alpha \leq x \leq 1 \\ -x - \ln x, & x > 1 \end{cases} \end{aligned}$$

Call the above function $h(x)$. Then

$$\begin{aligned} h'(x) &= \begin{cases} \frac{-\ln 3 - 1}{3} - \frac{1}{3x}, & x < \alpha \\ \frac{d}{dx}(-xc(x)), & \alpha < x < 1 \\ -1 - \frac{1}{x}, & x > 1 \end{cases} \\ &= \begin{cases} \frac{-\ln 3 - 1}{3} - \frac{1}{3x}, & x < \alpha \\ -c(x) - xc'(x), & \alpha < x < 1 \\ -1 - \frac{1}{x}, & x > 1 \end{cases} \end{aligned}$$

¹Credit for work past this part to the solution presented here

Now, since $D_{2,2}f(x, y) \neq 0$ for all x, y , part a) applies and

$$c'(x) = -\frac{D_{2,1}f(x, c(x))}{D_{2,2}f(x, c(x))} = -\frac{\ln c(x) - \frac{1}{x}}{\frac{x}{c(x)}} = -\frac{\frac{\ln x - 1}{x}}{\frac{x}{c(x)}} = -c(x) \frac{\ln x - 1}{x^2}$$

Thus

$$h'(x) = \begin{cases} \frac{-\ln 3 - 1}{3} - \frac{1}{3x}, & x < \alpha \\ -c(x) \frac{x+1-\ln x}{x}, & \alpha < x < 1 \\ -1 - \frac{1}{x}, & x > 1 \end{cases}$$

Note that $x > \ln x$, so $\frac{x+1-\ln x}{x} > 0$ and $c(x) > 0$, so $h'(x)$ is negative everywhere (except possibly the boundary points $\alpha, 1$, but it is continuous there). Thus the minimum of h on $[\frac{1}{2}, 2]$ is given when $x = \frac{1}{2}$. To check whether $\frac{1}{2} < \alpha$, simply note that $c(\frac{1}{2}) = \frac{1}{4} < \frac{1}{3}$, so $\frac{1}{2} < \alpha$. Thus

$$\max_{\frac{1}{2} \leq x \leq 2} \left(\min_{\frac{1}{2} \leq y \leq 2} f(x, y) \right) = h\left(\frac{1}{2}\right) = \frac{-\ln 3 - 1 - 2 \ln \frac{1}{2}}{6} = \frac{\ln \frac{3}{4} - 1}{6}$$

A.3 Chapter 3 Exercises

Exercise 3-1 Let $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} 0, & x \in [0, \frac{1}{2}) \\ 1, & x \in [\frac{1}{2}, 1] \end{cases}$$

Show that f is integrable and $\int_{[0,1] \times [0,1]} f = \frac{1}{2}$.

Proof. Let $\varepsilon > 0$. Choose a partition \mathcal{P} with subrectangles given by

$$\begin{aligned} A &= \left[0, \frac{1}{2} - \frac{\varepsilon}{2}\right] \times [0, 1] \\ B &= \left[\frac{1}{2} - \frac{\varepsilon}{2}, \frac{1}{2} + \frac{\varepsilon}{2}\right] \times [0, 1] \\ C &= \left[\frac{1}{2} + \frac{\varepsilon}{2}, 1\right] \times [0, 1] \end{aligned}$$

Then

$$\begin{aligned} m_A(f) &= M_A(f) = 0 \\ m_B(f) &= 0, M_B(f) = 1 \\ m_C(f) &= M_C(f) = 1 \end{aligned}$$

and

$$v(B) = \varepsilon$$

So

$$\begin{aligned} U(f, \mathcal{P}) - L(f, \mathcal{P}) &= v(B)(M_B(f) - m_B(f)) \\ &= v(B) = \varepsilon \end{aligned}$$

So f is integrable by the alternate criterion for integrability. Moreover,

$$U(f, \mathcal{P}) = v(A)M_A(f) + v(B)M_B(f) + v(C)M_C(f) = v(B \sqcup C) = \frac{1}{2} + \frac{\varepsilon}{2}$$

and similarly

$$L(f, \mathcal{P}) = \frac{1}{2} - \frac{\varepsilon}{2}$$

So $L \geq \frac{1}{2} - \frac{\varepsilon}{2}$ and $U \leq \frac{1}{2} + \frac{\varepsilon}{2}$, but we know that $U = L$ so $\int_A f = \frac{1}{2}$. □

Exercise 3-2 Let $f : A \rightarrow \mathbb{R}$ be integrable and let $g = f$ except at finitely many points. Show that g is integrable and $\int_A f = \int_A g$.

Proof. Refer to Exercise 3-3. Its proof does not depend on this problem, and we will use the fact that $\int_A f + g = \int_A f + \int_A g$ when f, g are integrable.

Let $\varepsilon > 0$ be arbitrary. We aim to show that $g - f$ is integrable with $\int_A g - f = 0$. Since $g \neq f$ at only finitely many points, it is bounded. Let $\mu = \max\{|f - g|\}$. Let p_1, \dots, p_k be those points where $g - f \neq 0$. Let S_1, \dots, S_k be the subrectangles they are in for a given partition (pick them small enough that they are distinct). Then choose \mathcal{P} such that

$$\sum_{i=1}^k v(S_i) < \varepsilon$$

Then

$$\begin{aligned} U(g - f, \mathcal{P}) - L(g - f, \mathcal{P}) &= \sum_{S \in \mathcal{P}} [M_S(g - f) - m_S(g - f)]v(S) \\ &= \sum_{i=1}^k [M_{S_i}(g - f) - m_{S_i}(g - f)]v(S_i) \\ &= \sum_{i=1}^k v(S_i) \\ &< \varepsilon \end{aligned}$$

So $g - f$ is integrable and a similar argument shows $\int_A g - f = 0$. So $\int_A g = \int_A g - f + f = \int_A g - f + \int_A f = \int_A f$. □

Exercise 3-3 Let $f, g : A \rightarrow \mathbb{R}$ be integrable.

(a) For any partition \mathcal{P} of A and subrectangle $S \in \mathcal{P}$, show that

$$m_S(f) + m_S(g) \leq m_S(f + g)$$

and

$$M_S(f + g) \leq M_S(f) + M_S(g)$$

so that

$$L(f, \mathcal{P}) + L(g, \mathcal{P}) \leq L(f + g, \mathcal{P})$$

and

$$U(f + g, \mathcal{P}) \leq U(f, \mathcal{P}) + U(g, \mathcal{P})$$

(b) Show that $f + g$ is integrable and $\int_A f + g = \int_A f + \int_A g$.

(c) For any constant c , show that $\int_A cf = c \int_A f$.

(a) *Proof.* Let $S \in \mathcal{P}$. Then for any point $x \in S$, we have

$$(f + g)(x) = f(x) + g(x) \geq m_S(f) + m_S(g)$$

Thus

$$m_S(f) + m_S(g) \leq m_S(f + g)$$

Similarly,

$$M_S(f + g) \leq M_S(f) + M_S(g)$$

Thus we have

$$\begin{aligned} L(f, \mathcal{P}) + L(g, \mathcal{P}) &= \sum_{S \in \mathcal{P}} v(S)[m_S(f) + m_S(g)] \\ &\leq \sum_{S \in \mathcal{P}} v(S)m_S(f + g) \\ &= L(f + g, \mathcal{P}) \end{aligned}$$

Similarly,

$$U(f + g, \mathcal{P}) \leq U(f, \mathcal{P}) + U(g, \mathcal{P})$$

□

(b) *Proof.* Let $\varepsilon > 0$ be arbitrary. Pick $\mathcal{P}_1, \mathcal{P}_2$ such that

$$\begin{aligned} U(f_1, \mathcal{P}_1) - L(f_1, \mathcal{P}_1) &< \frac{\varepsilon}{2} \\ U(f_2, \mathcal{P}_2) - L(f_2, \mathcal{P}_2) &< \frac{\varepsilon}{2} \end{aligned}$$

Let \mathcal{Q} be the common refinement of $\mathcal{P}_1, \mathcal{P}_2$. Then

$$\begin{aligned}
 U(f_1 + f_2, \mathcal{Q}) - L(f_1 + f_2, \mathcal{Q}) &= \sum_{S \in \mathcal{Q}} v(S)[M_S(f_1 + f_2) - m_S(f_1 + f_2)] \\
 &\leq \sum_{S \in \mathcal{Q}} v(S)[M_S(f_1) + M_S(f_2) - m_S(f_1) - m_S(f_2)] \\
 &= U(f_1, \mathcal{Q}) + U(f_2, \mathcal{Q}) - L(f_1, \mathcal{Q}) - L(f_2, \mathcal{Q}) \\
 &\leq U(f_1, \mathcal{P}_1) - L(f_1, \mathcal{P}_1) + U(f_2, \mathcal{P}_2) - L(f_2, \mathcal{P}_2) \\
 &< \varepsilon
 \end{aligned}$$

So $f_1 + f_2$ is integrable and a similar argument shows $\int_A f_1 + f_2 = \int_A f_1 + \int_A f_2$. \square

(c) *Proof.* Let \mathcal{P} be a partition and let $S \in \mathcal{P}$. Since S is a closed rectangle, it is compact, so there exists $x \in S$ with $f(x) = M_S(f)$. Then $(cf)(x) = cM_S(f)$ so $M_S(cf) \geq cM_S(f)$. But for any $y \in S$, we also have $(cf)(y) = cf(y) \leq cM_S(f)$ so $M_S(cf) = cM_S(f)$. Similarly, $m_S(cf) = cm_S(f)$.

Now, let $\varepsilon > 0$. Then there exists a partition \mathcal{P} with

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \frac{\varepsilon}{c}$$

Then we have

$$\begin{aligned}
 U(cf, \mathcal{P}) - L(cf, \mathcal{P}) &= \sum_{S \in \mathcal{P}} v(S)[M_S(cf) - m_S(cf)] \\
 &= \sum_{S \in \mathcal{P}} cv(S)[M_S(f) - m_S(f)] \\
 &= c[U(f, \mathcal{P}) - L(f, \mathcal{P})] \\
 &< \varepsilon
 \end{aligned}$$

So that cf is integrable. Now, let $\varepsilon > 0$ be arbitrary. Then there exists a partition \mathcal{P} such that

$$U(f, \mathcal{P}) \leq \int_A f + \frac{\varepsilon}{c}$$

Then we have

$$U(cf, \mathcal{P}) \leq c \int_A f + \varepsilon$$

So $\int_A cf = c \int_A f$. \square

Exercise 3-4 Let $f : A \rightarrow \mathbb{R}$ and let \mathcal{P} be a partition of A . Show that f is integrable if and only if, for each subrectangle $S \in \mathcal{P}$ the restriction $f|_S$ of f to S is integrable, and in this case $\int_A f = \sum_S \int_S f|_S$.

Proof. (\implies) Suppose that f is integrable on A , and let \mathcal{P} be given. Let $\varepsilon > 0$. Then there exists a partition \mathcal{P}' of A with

$$U(f, \mathcal{P}') - L(f, \mathcal{P}') < \varepsilon$$

Now let \mathcal{Q} be the common refinement of \mathcal{P} and \mathcal{P}' . Then each subrectangle of \mathcal{Q} is entirely contained within a subrectangle of \mathcal{P} . In other words, for any $S \in \mathcal{P}$, we may enumerate $S_1, \dots, S_k \in \mathcal{Q}$ such that $S_1 \sqcup \dots \sqcup S_k = S$, which means that $\mathcal{S} = \{S_1, \dots, S_k\}$ is a partition of S . Thus

$$\begin{aligned} U(f|_S, \mathcal{S}) - L(f|_S, \mathcal{S}) &= \sum_{S' \in \mathcal{S}} v(S') [M_{S'}(f|_S) - m_{S'}(f|_S)] \\ &\leq v(S) [M_S(f) - m_S(f)] \\ &\leq \sum_{S'' \in \mathcal{P}} v(S'') [M_{S''}(f) - m_{S''}(f)] \\ &= U(f, \mathcal{P}) - L(f, \mathcal{P}) \\ &< \varepsilon \end{aligned}$$

So $f|_S$ is integrable on S .

(\impliedby) Let \mathcal{P} be given, and suppose each $f|_S$ is integrable on the respective S . Let $\varepsilon > 0$. Then let N be the number of subrectangles in the partition \mathcal{P} . For each S , pick a partition \mathcal{P}^S such that

$$U(f|_S, \mathcal{P}^S) - L(f|_S, \mathcal{P}^S) < \frac{\varepsilon}{N}$$

Now, suppose that $\mathcal{P}^S = (\mathcal{P}_1^S, \dots, \mathcal{P}_n^S)$. Then

$$\mathcal{Q}_1 := \bigcup_{S \in \mathcal{P}} \mathcal{P}_1^S$$

is a partition of $[a_1, b_1]$. Let $\mathcal{Q} := (\mathcal{Q}_1, \dots, \mathcal{Q}_n)$. Then \mathcal{Q} is a refinement of \mathcal{P} , and moreover, for any $S \in \mathcal{P}$, \mathcal{Q}^S (which is the collection of subrectangles in \mathcal{Q} which are contained in S) is a refinement of \mathcal{P}^S . Thus

$$\begin{aligned} U(f, \mathcal{Q}) - L(f, \mathcal{Q}) &= \sum_{S' \in \mathcal{Q}} v(S') [M_{S'}(f) - m_{S'}(f)] \\ &= \sum_{S \in \mathcal{P}} \sum_{S'' \in \mathcal{Q}^S} v(S'') [M_{S''}(f) - m_{S''}(f)] \\ &\leq \sum_{S \in \mathcal{P}} \sum_{S'' \in \mathcal{P}^S} v(S'') [M_{S''}(f) - m_{S''}(f)] \\ &= \sum_{S \in \mathcal{P}} [U(f|_S, \mathcal{P}^S) - L(f|_S, \mathcal{P}^S)] \\ &< \sum_{S \in \mathcal{P}} \frac{\varepsilon}{N} \\ &= \varepsilon \end{aligned}$$

So f is integrable on A . A similar argument shows that $\int_A f = \sum_S \int_S f|_S$. \square

Exercise 3-5 Let $f, g : A \rightarrow \mathbb{R}$ be integrable and suppose $f \leq g$. Show that $\int_A f \leq \int_A g$.

Proof. Let \mathcal{P} be a partition of A . Then for any $S \in \mathcal{P}$, $M_S(f) \leq M_S(g)$. Thus

$$U(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} v(S)M_S(f) \leq \sum_{S \in \mathcal{P}} v(S)M_S(g) = U(g, \mathcal{P})$$

Since we know f and g are integrable, we conclude that

$$\int_A f = \inf U(f, \mathcal{P}) \leq \inf U(g, \mathcal{P}) = \int_A g \quad \square$$

Exercise 3-6 If $f : A \rightarrow \mathbb{R}$ is integrable, show that $|f|$ is integrable and $|\int_A f| \leq \int_A |f|$.

Proof. Let $\varepsilon > 0$. Let \mathcal{P} be a partition such that

$$U(f, \mathcal{P}) - L(f, \mathcal{P}) < \varepsilon$$

Let $S \in \mathcal{P}$. If $M_S(f) \geq m_S(f) \geq 0$, then $M_S(|f|) = M_S(f)$ and $m_S(|f|) = m_S(f)$. If $m_S(f) \leq M_S(f) \leq 0$, then $M_S(|f|) = -m_S(f)$ and $m_S(|f|) = -M_S(f)$. If $M_S(f) > 0$ and $m_S(f) < 0$, then I claim that $M_S(|f|) \leq \max\{|M_S(f)|, |m_S(f)|\}$.

To see this, note that for any $x \in S$, if $f(x) < 0$ then $|f(x)| = -f(x) \leq -m_S(f) = |m_S(f)|$. If $f(x) > 0$, then $|f(x)| = f(x) \leq M_S(f) = |M_S(f)|$. So $M_S(|f|) \leq \max\{|M_S(f)|, |m_S(f)|\}$. Using the fact that $m_S(|f|) \geq 0$, we have

$$\begin{aligned} M_S(|f|) - m_S(|f|) &\leq M_S(|f|) \\ &\leq \max\{|M_S(f)|, |m_S(f)|\} \\ &= \begin{cases} M_S(f), & |M_S(f)| \geq |m_S(f)| \\ -m_S(f), & |m_S(f)| > |M_S(f)| \end{cases} \\ &\leq M_S(f) - m_S(f) \end{aligned}$$

As a result, we have the following:

$$\begin{aligned} M_S(|f|) - m_S(|f|) &\leq \begin{cases} M_S(f) - m_S(f), & M_S(f) \geq m_S(f) \geq 0 \\ -m_S(f) - (-M_S(f)), & m_S(f) \leq M_S(f) \leq 0 \\ M_S(f) - m_S(f), & M_S(f) > 0, m_S(f) < 0 \end{cases} \\ &= M_S(f) - m_S(f) \end{aligned}$$

Thus, we have

$$\begin{aligned}
 U(|f|, \mathcal{P}) - L(|f|, \mathcal{P}) &= \sum_{S \in \mathcal{P}} v(S)[M_S(|f|) - m_S(|f|)] \\
 &\leq \sum_{S \in \mathcal{P}} v(S)[M_S(f) - m_S(f)] \\
 &= U(f, \mathcal{P}) - L(f, \mathcal{P}) \\
 &< \varepsilon
 \end{aligned}$$

So $|f|$ is integrable.

For any partition \mathcal{P} , and any $S \in \mathcal{P}$, we showed that $M_S(|f|) \leq \max\{|M_S(f)|, |m_S(f)|\}$. However, we can make a stronger statement, that $M_S(|f|) = \max\{|M_S(f)|, |m_S(f)|\}$. Indeed, since S is compact there exists $x, y \in S$ with $f(x) = M_S(f)$ and $f(y) = m_S(f)$. Then $|f|(x) = |M_S(f)|$ and $|f|(y) = |m_S(f)|$ so $|f|$ attains the value of $\max\{|M_S(f)|, |m_S(f)|\}$. Thus $|M_S(f)| \leq M_S(|f|)$. So

$$\left| \int_A f \right| \leq |U(f, \mathcal{P})| = \left| \sum_{S \in \mathcal{P}} v(S)M_S(f) \right| \leq \sum_{S \in \mathcal{P}} v(S)|M_S(f)| \leq \sum_{S \in \mathcal{P}} v(S)M_S(|f|) = U(|f|, \mathcal{P})$$

So for any partition \mathcal{P} , $U(|f|, \mathcal{P}) \geq |\int_A f|$ so $\int_A |f| \geq |\int_A f|$. □

Exercise 3-7 Let $f : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ be defined by

$$f(x, y) = \begin{cases} 0, & x \notin \mathbb{Q} \text{ or } y \notin \mathbb{Q} \\ \frac{1}{q}, & x \in \mathbb{Q}, y = \frac{p}{q} \in \mathbb{Q} \end{cases}$$

where we assume that $y = \frac{p}{q}$ is given in lowest terms. Show that f is integrable and $\int_{[0,1] \times [0,1]} f = 0$.

Proof. First, note that for any partition \mathcal{P} the density of \mathbb{Q} implies that $L(f, \mathcal{P}) = 0$. So it suffices to show that $U = 0$.

Let $\varepsilon > 0$. Pick a partition \mathcal{P} as follows: Choose N large enough that

$$\frac{1}{N} < \frac{\varepsilon}{2}$$

Then there are finitely many $y = p/q \in \mathbb{Q}$ such that $q < N$. Denote them by y_1, \dots, y_k . Then pick intervals I_1, \dots, I_k about each such that the total length of the intervals is less than $\varepsilon/2$ (and such that the I_i are disjoint). Let \mathcal{P}_2 be the partition of $[0, 1]$ given by these intervals, with the gaps filled in appropriately.

Let \mathcal{P}_1 be the single partition $\{0, 1\}$. Then $\mathcal{P} = (\mathcal{P}_1, \mathcal{P}_2)$ consists of subrectangles of the form $[0, 1] \times I$, where I is either one of the I_i we defined previously, or it is not (in this case,

it is a gap between them). Let \mathcal{L} denote the set of all subrectangles of the form $[0, 1] \times I_i$, and let \mathcal{R} denote the set of all other subrectangles. Then

$$U(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} v(S)M_S(f) = \sum_{S \in \mathcal{L}} v(S)M_S(f) + \sum_{S \in \mathcal{R}} v(S)M_S(f)$$

Now, if $S \in \mathcal{L}$, then f attains a value of at most 1 on S , so $M_S(f) \leq 1$. But if $M_S(f) \in \mathcal{R}$, then by construction there is no point $(x, y) \in S$ with $y = p/q$ and $q < N$. Thus

$$f(x, y) = \frac{1}{q} < \frac{1}{N} < \frac{\varepsilon}{2}$$

so $M_S(f) \leq \frac{\varepsilon}{2}$. Thus

$$\sum_{S \in \mathcal{L}} v(S)M_S(f) + \sum_{S \in \mathcal{R}} v(S)M_S(f) \leq \sum_{S \in \mathcal{L}} v(S) + \frac{\varepsilon}{2} \sum_{S \in \mathcal{R}} v(S) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

Thus $U = 0$. So f is integrable and $\int_{[0,1] \times [0,1]} f = 0$. □

Exercise 3-8 Prove that $A = [a_1, b_1] \times \dots \times [a_n, b_n]$ does not have content zero if $a_i < b_i$ for each i .

Proof. Let \mathcal{O} be a finite cover of A by closed rectangles. Without loss of generality we may assume that each rectangle is contained within A . Then let $T_i = \{t_0^i, \dots, t_{k_i}^i\}$ be the set of endpoints of the rectangles in the i th direction (that is, if $O = \{[c_1, d_1] \times \dots \times [c_n, d_n]\} \in \mathcal{O}$, then $c_1, d_1 \in T_1$ and $c_i, d_i \in T_i$ for any i). Without loss of generality we may order them so that $a_i = t_0^i \leq \dots \leq t_{k_i}^i = b_i$. Then each $v(O_i)$ for $O_i \in \mathcal{O}$ is the sum of $v(A_i)$ for A_i of the form $[t_{j_1-1}^1, t_{j_1}^1] \times \dots \times [t_{j_n-1}^n, t_{j_n}^n]$. Moreover, each of those rectangles is contained within some O_i . So

$$\sum_{i=1}^n v(O_i) \geq \sum_{j=1}^{k_1 \times \dots \times k_n} v(A_j) = \prod_{j=1}^n (b_j - a_j)$$

So A does not have content zero. □

Exercise 3-9

- (a) Show that an unbounded set cannot have content zero.
- (b) Give an example of a closed set of measure zero which does not have content zero.

- (a) *Proof.* Let A be an unbounded set and \mathcal{O} a finite cover of A by closed rectangles. Then there exists a closed rectangle M such that

$$\bigcup_{O \in \mathcal{O}} O \subseteq M$$

But since A is unbounded it contains points outside M . So \mathcal{O} cannot be a cover of A , contradiction. Thus A is in fact not covered by any finite set of closed (or open) rectangles, so it cannot have content zero. □

- (b) *Proof.* \mathbb{Q} is closed and has measure zero (this follows from the fact that it is countable). However, it is unbounded, and thus does not have content zero by part a). \square

Exercise 3-10

- (a) If C is a set of content zero, show that the boundary of C has content zero.
 (b) Give an example of a bounded set C of measure zero such that the boundary of C does not have measure zero.

- (a) *Proof.* Let \mathcal{O} be a finite cover of C by closed rectangles. I claim that \mathcal{O} contains ∂C . To see this, suppose that there exists a point $x \in \partial C$ such that $x \notin O$ for each $O \in \mathcal{O}$. Then

$$x \in \bigcap_{i=1}^k \mathbb{R}^n \setminus O_i$$

But since each O_i is closed, $\mathbb{R}^n \setminus O_i$ is open, and this is a finite intersection of open sets, which is open. Then since $x \in \partial C$, there exists a point $y \in C$ with

$$y \in \bigcap_{i=1}^k \mathbb{R}^n \setminus O_i$$

But this contradicts the assumption that \mathcal{O} is a cover of C . Thus \mathcal{O} covers ∂C . So any closed cover of C is a cover of ∂C . Then let $\varepsilon > 0$. We may produce a finite cover of C by closed rectangles with total volume less than ε . This cover works for ∂C as well. Thus ∂C has content zero. \square

- (b) Pick $\mathbb{Q} \cap [0, 1]$. This is a bounded set of measure zero. But $\partial(\mathbb{Q} \cap [0, 1]) = [0, 1]$, which does not have measure zero.

Exercise 3-11 Let A be the union of open intervals (a_i, b_i) such that each rational number in $(0, 1)$ is contained in some (a_i, b_i) , as in Exercise 1-18. If

$$\sum_{i=1}^{\infty} b_i - a_i < 1$$

show that ∂A does not have measure zero.

Proof. Suppose that ∂A has measure zero. Pick a cover \mathcal{O} of ∂A by open intervals such that

$$\sum_{O \in \mathcal{O}} v(O) < 1 - \sum_{i=1}^{\infty} b_i - a_i$$

which we rewrite as

$$1 > \sum_{O \in \mathcal{O}} v(O) + \sum_{i=1}^{\infty} b_i - a_i$$

From Exercise 1-18, we know that $\partial A = [0, 1] \setminus A$. So the collection of intervals in \mathcal{O} combined with the open intervals which make up A form a cover of $[0, 1]$ by open intervals. Call this cover \mathcal{O}' . Then we know

$$\sum_{O \in \mathcal{O}'} v(O) \geq 1$$

But we also have

$$\sum_{O \in \mathcal{O}} v(O) + \sum_{i=1}^{\infty} b_i - a_i \geq \sum_{O \in \mathcal{O}'} v(O)$$

So

$$1 > \sum_{O \in \mathcal{O}} v(O) + \sum_{i=1}^{\infty} b_i - a_i \geq \sum_{O \in \mathcal{O}'} v(O) \geq 1$$

and we conclude that $1 > 1$, contradiction. So ∂A does not have measure zero. \square

Exercise 3-12 Let $f : [a, b] \rightarrow \mathbb{R}$ be an increasing function. Show that $\{x : f \text{ is discontinuous at } x\}$ has measure zero.

Proof. I claim that for any n , there are at most $n(f(b) - f(a))$ points with $o(f, x) > \frac{1}{n}$.

To prove this, suppose there are more than $n(f(b) - f(a))$ such points, x_1, \dots, x_k . Then we may pick y_0, \dots, y_k with $a = y_0 < x_1 < y_1 < \dots < x_k < y_k = b$. Then because f is increasing, for each x_i we have

$$o(f, x_i) \leq f(y_i) - f(y_{i-1})$$

Then by a telescoping argument,

$$\sum_{i=1}^k o(f, x_i) \leq f(y_k) - f(y_0) = f(b) - f(a)$$

But we also have

$$\sum_{i=1}^k o(f, x_i) \geq \sum_{i=1}^k \frac{1}{n} = \frac{k}{n} > \frac{n(f(b) - f(a))}{n} = f(b) - f(a)$$

contradiction. Thus there are at most $n(f(b) - f(a))$ such points. Recall that f is discontinuous at x precisely when $o(f, x) > 0$. But

$$\{x : o(f, x) > 0\} = \bigcup_{n=1}^{\infty} \{x : o(f, x) > \frac{1}{n}\}$$

So $\{x : f \text{ is discontinuous at } x\}$ is the countable union of finite sets and thus has measure zero. \square

Exercise 3-13

- (a) Show that the collection of all rectangles $[a_1, b_1] \times \dots \times [a_n, b_n]$ with all a_i and b_i rational can be arranged in a sequence.
- (b) If $A \subseteq \mathbb{R}^n$ is any set and \mathcal{O} is an open cover of A , show that there is a sequence O_1, O_2, \dots of members of \mathcal{O} which also cover A .

(a) *Proof.* This collection may be placed in bijection with \mathbb{Q}^{2n} , which is a finite Cartesian product of countable sets, so it is countable. \square

(b) *Proof.* For each point $x \in A$, $x \in O$ for some $O \in \mathcal{O}$, and O is open, so there exists an open rectangle $R_x \subseteq O$ containing x . Moreover, we demand that each endpoint of R_x is rational. Then the set of $R = \{R_x : x \in A\}$ is a subset of the set of all rectangles with rational endpoints, which we showed is countable. Thus R is countable, so we may order its elements as R_1, R_2, \dots

We then pick a countable subcover \mathcal{O}' of \mathcal{O} by picking O'_1 such that $R_1 \subseteq O'_1$, and so on. We may skip terms if R_i is already contained in a previously chosen open set. This gives a countable subcover of R , and R covers A , so this is a countable subcover of A . \square

Exercise 3-14 Show that if $f, g : A \rightarrow \mathbb{R}$ are integrable, then fg is as well.

Proof. Since f and g are both integrable, they are discontinuous on sets $C_1, C_2 \subseteq A$ of measure zero. For any x such that $x \notin C_1$ and $x \notin C_2$, f, g are both continuous at x so fg is continuous at x . Thus C_3 , the set of points where fg is discontinuous, is a subset of $C_1 \cup C_2$ and has measure zero. So fg is integrable. \square

Exercise 3-15 Show that if C has content zero, then $C \subseteq A$ for some closed rectangle A and C is Jordan measurable with $\int_A \chi_C = 0$

Proof. We showed in Exercise 3-9 part a) that any unbounded set does not have content zero. So $C \subseteq A$ for a closed rectangle A . We showed in Exercise 3-10 part a) that ∂C has content zero whenever C has content zero. So C is Jordan-measurable.

Now pick a partition \mathcal{P} of A . For every subrectangle S of \mathcal{P} , we cannot have $S \subseteq C$, since otherwise C would not have content zero. So $m_S(\chi_C) = 0$ for each S and thus $L(f, \mathcal{P}) = 0$. This is true for all partitions \mathcal{P} , so

$$\int_A \chi_C = L = 0 \quad \square$$

Exercise 3-16 Give an example of a bounded set C of measure zero such that $\int_A \chi_C$ does not exist.

Set $C = \mathbb{Q} \cap [0, 1]$. Then χ_C is the Dirichlet function, which is discontinuous on $[0, 1]$ (since both irrationals and rationals are dense in $[0, 1]$). So χ_C is not discontinuous on a set of measure zero, so $\int_A \chi_C$ does not exist.

Exercise 3-17 If C is a bounded set of measure zero and $\int_A \chi_C$ exists, show that $\int_A \chi_C = 0$.

Proof. See the second paragraph of the argument from Exercise 3-15. □

Exercise 3-18 If $f : A \rightarrow \mathbb{R}$ is nonnegative and $\int_A f = 0$, show that $\{x : f(x) \neq 0\}$ has measure zero.

Proof. Consider the set $B_n = \{x : f(x) \geq \frac{1}{n}\}$ for any n . I claim that B_n has content zero. Suppose it does not. Then there exists $\varepsilon > 0$ such that any cover of B_n has total volume no less than ε . Then let \mathcal{P} be any partition. If \mathcal{S} is the collection of subrectangles which intersect B_n , then $M_S(f) \geq \frac{1}{n}$ for any $S \in \mathcal{S}$. So

$$U(f, \mathcal{P}) = \sum_{S \in \mathcal{P}} v(S)M_S(f) \geq \sum_{S \in \mathcal{S}} v(S)M_S(f) \geq \frac{1}{n} \sum_{S \in \mathcal{S}} v(S) \geq \frac{\varepsilon}{n}$$

So $U \geq \frac{\varepsilon}{n} > 0$, but this contradicts the assumption that $\int_A f = 0$. So B_n has content zero. Thus

$$\{x : f(x) \neq 0\} = \bigcup_{n=1}^{\infty} B_n$$

has measure zero. □

Exercise 3-19 Let U be the union of open intervals (a_i, b_i) such that each rational number in $(0, 1)$ is contained in some (a_i, b_i) , and

$$\sum_{i=1}^{\infty} b_i - a_i < 1$$

as in Exercise 3-11. Show that if $f = \chi_U$ except on a set of measure zero, then f is not integrable on $[0, 1]$.

Proof. In Exercise 3-11 we showed that $\partial U = [0, 1] \setminus U$ does not have measure zero. χ_U is discontinuous on ∂U , so it is discontinuous on a set that is not of measure zero, and thus not integrable. Then we need to show that f is also discontinuous on a set not of measure zero.

Let $x \in \partial U$, and suppose that $f(x) = \chi_U(x)$. Suppose for contradiction, suppose that f is continuous at x . Since $x \in \partial U$ and $\partial U = [0, 1] \setminus U$, $x \notin U$. Thus $f(x) = \chi_U(x) = 0$. If f is continuous at x , then for any $\varepsilon > 0$ there exists a neighborhood around x such that $|f(y)| < \varepsilon$ for y in the neighborhood. We will show that this is not the case.

Let $\varepsilon = \frac{1}{2}$. Let V be any neighborhood around x contained in $[0, 1]$. Then there exists a rational $q \in V$. $q \in U$ which is open, so there exists an open rectangle R containing q contained in $U \cap V$. So $\chi_U = 1$ on an open rectangle within V . So if $|f(y)| < \varepsilon$ for any $y \in V$, we must have $f \neq \chi_U$ on R . But R is not a set of measure zero, so this contradicts the assumption that $f = \chi_U$ on a set of measure zero. So f is not continuous at x .

We have shown that for any $x \in \partial U$ such that $f(x) = \chi_U(x)$, f is discontinuous at x . Then we must show that the set of $x \in \partial U$ with $f(x) = \chi_U(x)$ does not have measure zero.

Suppose that it does. Let $\varepsilon > 0$. Then there exists a cover \mathcal{U} of $\{x \in \partial U : f(x) = \chi_U(x)\}$ by open intervals with total length less than $\varepsilon/2$. We also know that $\{x \in [0, 1] : f(x) \neq \chi_U(x)\}$ has measure zero by assumption, so $\{x \in \partial U : f(x) \neq \chi_U(x)\}$ also has measure zero and we may cover it by an open cover \mathcal{O} with total length less than $\varepsilon/2$.

Now for any $x \in \partial U$, we must have $f(x) = \chi_U(x)$ or $f(x) \neq \chi_U(x)$, so $\mathcal{U} \cup \mathcal{O}$ covers ∂U . Now we have

$$\sum_{(c_i, d_i) \in \mathcal{U} \cup \mathcal{O}} d_i - c_i \leq \sum_{(c_i, d_i) \in \mathcal{U}} d_i - c_i + \sum_{(c_i, d_i) \in \mathcal{O}} d_i - c_i < \varepsilon$$

So ∂U has measure zero. But in Exercise 3-11 we showed that this is not the case. So the assumption that $\{x \in \partial U : f(x) = \chi_U(x)\}$ has measure zero is incorrect. But we showed that f is discontinuous on this set, and it does not have measure zero, so f is not integrable on $[0, 1]$. \square

Exercise 3-20 Show that an increasing function $f : [a, b] \rightarrow \mathbb{R}$ is integrable on $[a, b]$.

Proof. In Exercise 3-12 we showed that f is discontinuous on a set of measure zero. So it is integrable on $[a, b]$. \square

Exercise 3-21 If A is a closed rectangle, show that $C \subseteq A$ is Jordan-measurable if and only if for every $\varepsilon > 0$ there is a partition \mathcal{P} of A such that

$$\sum_{S \in \mathcal{S}_1} v(S) - \sum_{S \in \mathcal{S}_2} v(S) < \varepsilon$$

where \mathcal{S}_1 consists of all subrectangles intersecting C and \mathcal{S}_2 all subrectangles contained in C .

We first prove the following fact:

Lemma

If $A \subseteq \mathbb{R}^n$ and $x \in \text{int } A$, $y \in \text{ext } A$, then there exists $z = tx + (1-t)y$ with $0 \leq t \leq 1$ such that $z \in \partial A$. (Intuitively, this z lies along the line segment between x and y).

Proof. To see this, first note for sufficiently small $t > 0$, $tx + (1-t)y \in A$ since $x \in \text{int } A$. Thus the set $\{0 \leq t \leq 1 : tx + (1-t)y \in A\}$ is nonempty. Moreover, it is clearly bounded. Then let

$$t' = \sup\{0 \leq t \leq 1 : tx + (1-t)y \in A\}$$

Now, first note that $t' < 1$. This is because $y \in \text{ext } A$, so there exists a ball around y entirely contained in $\mathbb{R}^n \setminus A$.

I claim that $z = t'x + (1-t')y \in \partial A$. To see this, let $B_r(z)$ be any open ball around z . $B_r(z)$ contains a point in A , as we can simply pick $tx + (1-t)y$ for $t \leq t'$ such that $t' - t < r$. Then we need to show that $B_r(z)$ contains a point in $\mathbb{R}^n \setminus A$.

Let $z' = (t' + \varepsilon)x + (1 - t' - \varepsilon)y$, where $\varepsilon < r$ and $t' + \varepsilon < 1$ (possible because $t' < 1$). Then $|z - z'| = \varepsilon < r$, so $z' \in B_r(z)$. But $t' + \varepsilon > t'$, so $t' + \varepsilon \notin \{0 \leq t \leq 1 : tx + (1-t)y \in A\}$. Since we provided that $t' + \varepsilon < 1$, we conclude that $z' \notin A$. So $z \in \partial A$. \square

Now, continuing to the main proof:

Proof. (\implies) Suppose that $C \subseteq A$ is Jordan-measurable. ∂C has measure zero, and is compact, so we may pick a finite collection of closed rectangles \mathcal{O} whose interiors cover ∂C with total volume is less than ε . Then apply Lemma 3.9 to pick a partition \mathcal{P} such that every subrectangle $S \in \mathcal{P}$ is either contained in some $O \in \mathcal{O}$ or does not intersect ∂C . If $S \in \mathcal{P}$ and S intersects C but is not contained in C , I claim that there exists $z \in S$ with $z \in \partial C$.

Indeed, we can pick $x, y \in S$ such that $x \in C$ and $y \notin C$. Then if either of these points is in ∂C , then we are done. Otherwise, $x \in \text{int } C$ and $y \in \text{ext } C$. By the Lemma, there exists $z = tx + (1-t)y$ with $0 \leq t \leq 1$ such that $z \in \partial C$. Since S is convex, $z \in S$. So the claim is proved. Then S intersects ∂C , so we must have $S \subseteq O$ for some $O \in \mathcal{O}$. Thus

$$\sum_{S \in \mathcal{S}_1} v(S) - \sum_{S \in \mathcal{S}_2} v(S) \leq \sum_{O \in \mathcal{O}} v(O) < \varepsilon$$

(\impliedby) Suppose that $C \subseteq A$ satisfies the condition that for every $\varepsilon > 0$ there is a partition \mathcal{P} such that

$$\sum_{S \in \mathcal{S}_1} v(S) - \sum_{S \in \mathcal{S}_2} v(S) < \varepsilon$$

I claim that $\mathcal{S}_1 \setminus \mathcal{S}_2$ covers ∂C . To see this, let $x \in \partial C$. Then $x \in S$ for some $S \in \mathcal{P}$. Then $S \in \mathcal{S}_1$ or $S \notin \mathcal{S}_1$. But if $S \notin \mathcal{S}_1$, then there exists an open rectangle (S) around x entirely contained in $\text{ext } C$, contradicting $x \in \partial C$. So $S \in \mathcal{S}_1$. But similarly, if $S \in \mathcal{S}_2$ then that contradicts $x \in \partial C$. So $S \in \mathcal{S}_1 \setminus \mathcal{S}_2$.

Thus $\mathcal{S}_1 \setminus \mathcal{S}_2$ covers ∂C , and by assumption it can be made as small as required. So ∂C has measure zero and C is Jordan-measurable. \square

Exercise 3-22 If A is Jordan-measurable and $\varepsilon > 0$, show that there exists a compact Jordan-measurable set $C \subseteq A$ such that $\int_{A \setminus C} 1 < \varepsilon$.

Proof. Let A be Jordan-measurable and let $\varepsilon > 0$. Then by Exercise 3-21, we may pick a partition \mathcal{P} such that

$$\sum_{S \in \mathcal{S}_1} v(S) - \sum_{S \in \mathcal{S}_2} v(S) < \varepsilon$$

where \mathcal{S}_1 is the collection of subrectangles intersecting A and \mathcal{S}_2 is the collection of subrectangles contained in A . Then $C = \bigcup \mathcal{S}_2$ is a union of finite closed rectangles and thus closed. Moreover, $C \subseteq A$. Since A is bounded, C is also bounded and thus compact. So we need to show that it is Jordan-measurable.

I claim that $\partial C \subseteq \bigcup_{S \in \mathcal{S}_2} \partial S$. Let $x \in \partial C$. Then consider the sequence of open balls (B_n) , where $B_n = B_{1/n}(x)$. Then for each B_n , there exists some point $y_n \in C$. Each $y_n \in S$ for some $S \in \mathcal{S}_2$, but there are only finitely many such S , so there is some S' such that $y_n \in S'$ for infinitely many n . Moreover, each B_n contains a point not contained in C , which is thus also not contained in S' . So $x \in \partial S'$. Thus the claim is proved.

We take without proof the fact that a rectangle is Jordan-measurable. Then ∂S has measure zero for each $S \in \mathcal{S}_2$, so the finite union $\bigcup_{S \in \mathcal{S}_2} \partial S$ also has measure zero, and thus ∂C has measure zero and C is Jordan measurable.

Now, because $C \subseteq A$, we have $\int_{A \setminus C} 1 = \int_A 1 - \int_C 1$. Moreover, \mathcal{S}_1 covers A . So

$$\int_A 1 \leq \int_{\bigcup \mathcal{S}_1} 1$$

and thus

$$\int_{A \setminus C} 1 = \int_A 1 - \int_C 1 \leq \int_{\bigcup \mathcal{S}_1} 1 - \int_C 1 = \sum_{S \in \mathcal{S}_1} v(S) - \sum_{S \in \mathcal{S}_2} v(S) < \varepsilon \quad \square$$

Exercise 3-23 Let $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$. Let $C \subseteq A \times B$ be a set of $n + m$ -dimensional content zero. Let $A' \subseteq A$ be the set of all $x \in A$ such that $\{y \in B : (x, y) \in C\}$ is not of m -dimensional content zero. Show that A' is a set of n -dimensional measure zero.

Proof. First, because C has content zero, ∂C has content zero so χ_C is integrable on $A \times B$ and $\int_{A \times B} \chi_C = 0$. Let $\mathcal{U}(x) = \int_B \chi_C(x, y) dy$. Then by Fubini's Theorem,

$$\int_{A \times B} \chi_C = \int_A \mathcal{U} = 0$$

Now, fix some $x \in A$, and let \mathcal{P} be a partition of B .

If $x \in A'$, then there exists some $\varepsilon_x > 0$ such that any finite cover of $\{y \in B : (x, y) \in C\}$ by closed rectangles has total length at least ε_x . Let \mathcal{S}_1 be the collection of subrectangles S in \mathcal{P} that intersect $\{y \in B : (x, y) \in C\}$. Because \mathcal{S}_1 is a finite cover of $\{y \in B : (x, y) \in C\}$,

$$U(\chi_C, \mathcal{P}) = \sum_{S \in \mathcal{S}_1} M_S(\chi_C) v(S) = \sum_{S \in \mathcal{S}_1} v(S) \geq \varepsilon_x$$

Then $\mathcal{U}(x) = \mathbf{U} \int_B \chi_C \geq \varepsilon_x$.

Now, \mathcal{U} is clearly nonnegative, and we know that $\int_A \mathcal{U} = 0$. So by Exercise 3-18, $\{x : \mathcal{U}(x) \neq 0\}$ has measure zero. But we just showed that $A' \subseteq \{x : \mathcal{U}(x) \neq 0\}$, so A' has measure zero. \square

Exercise 3-24 Let $C \subseteq [0, 1] \times [0, 1]$ be the union of all $\{p/q\} \times [0, 1/q]$, where p/q is a rational in $[0, 1]$ in lowest terms. Show that it is not true that the set A' in Exercise 3-23 has content zero.

Proof. First we show that C has content zero. Let $\varepsilon > 0$. Then let

$$R_0 = [0, 1] \times \left[0, \frac{\varepsilon}{2}\right]$$

Then there a finite number of rationals p/q such that $\{p/q\} \times [0, 1/q]$ is not contained in R_0 . Call these $r_1, \dots, r_k = p_1/q_1, \dots, p_k/q_k$. Then for $1 \leq i \leq k$, let

$$R_i = \left[\frac{p_i}{q_i} - \frac{q_i \varepsilon}{2^{i+1}}, \frac{p_i}{q_i} + \frac{q_i \varepsilon}{2^{i+1}}\right] \times \left[0, \frac{1}{q_i}\right]$$

Letting $\mathcal{R} = \{R_0, R_1, \dots, R_k\}$, \mathcal{R} is a finite cover of C by closed rectangles with

$$\sum_{R \in \mathcal{R}} v(R) = v(R_0) + \sum_{i=1}^k v(R_i) = \frac{\varepsilon}{2} + \sum_{i=1}^k \frac{\varepsilon}{2^{i+1}} \leq \frac{\varepsilon}{2} + \sum_{i=1}^{\infty} \frac{\varepsilon}{2^{i+1}} = \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

So C has content zero.

But for each rational $p/q \in [0, 1]$, the set $\{y \in [0, 1] : (p/q, y) \in C\}$ is simply the set $[0, 1/q]$, which does not have content zero. So $A' = \mathbb{Q} \cap [0, 1]$, which does not have content zero. \square

Exercise 3-25 Use induction on n to show that $[a_1, b_1] \times \dots \times [a_n, b_n]$ is not a set of measure zero (or content zero) if $a_i < b_i$.

Proof. In the base case, $n = 1$, let \mathcal{U} be a cover of $[a_1, b_1]$ by open intervals. Since $[a_1, b_1]$ is compact, we can assume \mathcal{U} is finite. From here the base case proceeds as in Exercise 3-8.

Now suppose the theorem is true for n , and we will prove it for $n + 1$. Then $[a_1, b_1] \times \dots \times [a_{n+1}, b_{n+1}] = ([a_1, b_1] \times \dots \times [a_n, b_n]) \times [a_{n+1}, b_{n+1}]$. Let $A = [a_1, b_1] \times \dots \times [a_n, b_n]$ and $B = [a_{n+1}, b_{n+1}]$. By Fubini's Theorem²

$$\int_{A \times B} 1 = \int_A \left(\int_B 1 \, dy \right) dx = \left(\int_A 1 \, dx \right) \left(\int_B 1 \, dy \right)$$

²Credit for work past this point to <https://hidenori-shinohara.github.io/2019/12/23/measure-zero-ex-3-25.html>

Now, the constant function 1 is a nonnegative function, and Exercise 3-18 shows that if $\int_A 1 \, dx = 0$, then 1 is nonzero on a set of measure zero. But 1 is nonzero on A , which is not a set of measure zero by the inductive hypothesis. So

$$\int_A 1 \, dx > 0$$

and similarly

$$\int_B 1 \, dy > 0$$

so

$$\int_{A \times B} 1 > 0$$

Now $A \times B$ is bounded. If it has measure zero, then Exercise 3-18 says that $\int_{A \times B} \chi_{A \times B} = \int_{A \times B} 1 = 0$. But this is not the case, so $A \times B$ does not have measure zero. \square

Exercise 3-26 Let $f : [a, b] \rightarrow \mathbb{R}$ be integrable and nonnegative and let $A_f = \{(x, y) : a \leq x \leq b, 0 \leq y \leq f(x)\}$. Show that A_f is Jordan-measurable and has area $\int_a^b f$.

Proof. Since $f : [a, b] \rightarrow \mathbb{R}$ is integrable and nonnegative, there exists $M > 0$ such that $M > f(x)$ for any x .

Claim A.1

Let

$$\begin{aligned} B &= ([a, b] \times \{0\}) \\ C &= \{(x, f(x)) : x \in [a, b]\} \\ D &= \{a\} \times [0, M] \\ E &= \{b\} \times [0, M] \\ F &= \{x : f \text{ is discontinuous at } x\} \times [0, M] \end{aligned}$$

Then

$$\partial A_f \subseteq B \cup C \cup D \cup E \cup F$$

To prove this, note that any (x, y) satisfies exactly one of the following conditions³:

1. $(x, y) \notin [a, b]$.
2. $x = a$.
3. $x = b$.

³Strictly speaking, conditions 5 and 8 are both filled by $(x, 0)$ for $x : f(x) = 0$, but this does not detract from the overall argument.

4. $(x, y) \in (a, b), y < 0$.
5. $(x, y) \in (a, b), y = 0$.
6. $(x, y) \in (a, b), 0 < y < f(x)$, f is continuous at x .
7. $(x, y) \in (a, b), 0 < y < f(x)$, f is not continuous at x .
8. $(x, y) \in (a, b), y = f(x)$.
9. $(x, y) \in (a, b), y > f(x)$, f is continuous at x .
10. $(x, y) \in (a, b), f(x) < y \leq M$, f is not continuous at x .
11. $(x, y) \in (a, b), y > M$, f is not continuous at x .

For cases 2, 3, 5, 7, 8, 10, $(x, y) \in B \cup C \cup D \cup E \cup F$. Thus we must show that $(x, y) \notin \partial A_f$ whenever conditions 1, 4, 6, 9, or 11 are met.

Case 1: We can pick an open rectangle R containing (x, y) such that $(x_1, y_1) \in R \implies x_1 \notin [a, b]$. So $(x, y) \in \text{ext } A_f$.

Case 4: We can pick an open rectangle R containing (x, y) such that $(x_1, y_1) \in R \implies y_1 < 0$. So $(x, y) \in \text{ext } A_f$.

Case 6: Since f is continuous at x , there exists an interval $(x - \delta, x + \delta)$ such that $f(x_1) > y + \varepsilon$ whenever $x_1 \in (x - \delta, x + \delta)$, for $\varepsilon > 0$ sufficiently small (where δ is chosen small enough that this makes sense). Then the rectangle $R = (x - \delta, x + \delta) \times (0, y + \varepsilon)$ is an open rectangle containing (x, y) which is contained in A_f . So $(x, y) \in \text{int } A_f$.

Case 9: Similarly to Case 4, since f is continuous at x , there exists an interval $(x - \delta, x + \delta)$ such that $f(x_1) < y - \varepsilon$ whenever $x_1 \in (x - \delta, x + \delta)$ for $\varepsilon > 0$ sufficiently small. Then the rectangle $R = (x - \delta, x + \delta) \times (y - \varepsilon, M)$ shows that $(x, y) \in \text{ext } A$.

Case 11: Similarly to Case 2, we may pick an open rectangle R containing (x, y) such that $(x_1, y_1) \in R \implies y_1 > M \implies (x_1, y_1) \notin A_f$. So $(x, y) \in \text{ext } A_f$.

Thus Claim 1 is proved.

Claim A.2

The sets B, C, D, E, F each have measure zero.

The line interval $[a, b] \times \{0\}$ has measure zero, as for any $\varepsilon > 0$ we cover it by

$$R_\varepsilon = [a, b] \times \left[-\frac{\varepsilon}{2(b-a)}, \frac{\varepsilon}{2(b-a)} \right]$$

which has $v(R_\varepsilon) = \varepsilon$. So B has measure zero. A similar proof holds for the line segments D and E .

The set $\{x : f \text{ is discontinuous at } x\}$ has measure zero since f is integrable. Let $\varepsilon > 0$. Then we may pick a cover \mathcal{I} of $\{x : f \text{ is discontinuous at } x\}$ by open intervals such that

$$\sum_{(c,d) \in \mathcal{I}} d - c < \frac{\varepsilon}{4M}$$

Then the collection \mathcal{U} of rectangles of the form $(c, d) \times (-\frac{M}{2}, \frac{3M}{2})$ for $(c, d) \in \mathcal{I}$ forms a cover of $\{x : f \text{ is discontinuous at } x\} \times [0, M]$. Moreover, consider the remaining set

$$S = [a, b] \setminus \bigcup_{I \in \mathcal{I}} I$$

Since each I is open, S is closed. It is also bounded, so it is compact. Moreover, f is continuous at each $x \in S$. Since f is continuous on S compact, it is uniformly continuous. Thus we may pick $\delta > 0$ such that

$$x, y \in S, |x - y| < \delta \implies |f(x) - f(y)| < \frac{\varepsilon}{4(b-a)}$$

Moreover, pick δ such that $m\delta = b - a$ for some $m \in \mathbb{N}$. Now let $\delta_i = [a + (i-1)\delta, a + i\delta]$. Then the collection $\{\delta_i\}_{i=1}^m$ partitions the interval $[a, b]$. Now for each i , define the rectangle P_i as follows: if $S \cap \delta_i = \emptyset$, then let $P_i = \delta_i \times \{0\}$. Otherwise, pick $x_i \in S \cap \delta_i$. Then let

$$P_i = \delta_i \times \left[f(x_i) - \frac{\varepsilon}{4(b-a)}, f(x_i) + \frac{\varepsilon}{4(b-a)} \right]$$

Let $\mathcal{P} = \{P_1, \dots, P_n\}$, and let $\bar{\mathcal{U}} = \{\bar{U} : U \in \mathcal{U}\}$. I claim that $\mathcal{P} \cup \bar{\mathcal{U}}$ is a cover of $C \cup F$. Indeed, we already showed that \mathcal{U} covers F , so $\bar{\mathcal{U}}$ does as well.

Now, for any $x \in [a, b]$, either $x \in S$ or $x \notin S$. If $x \notin S$, then $x \in I$ for some $I \in \mathcal{I}$ and thus $(x, f(x)) \in U$ for some $U \in \bar{\mathcal{U}}$. On the other hand, if $x \in S$, then $x \in \delta_i$ for some i (this does not require $x \in S$, just $x \in [a, b]$). Then $|x - x_i| < \delta_i$, so

$$|f(x) - f(x_i)| < \frac{\varepsilon}{4(b-a)}$$

so $(x, f(x)) \in P_i$. Thus $\mathcal{P} \cup \bar{\mathcal{U}}$ is a cover of $C \cup F$ by closed rectangles. Lastly, we have

$$\sum_{\bar{U} \in \bar{\mathcal{U}}} v(U) = \sum_{U \in \mathcal{U}} v(U) = \sum_{(c,d) \in \mathcal{I}} v((c,d) \times (-\frac{M}{2}, \frac{3M}{2})) = 2M \sum_{(c,d) \in \mathcal{I}} d - c < \frac{\varepsilon}{2}$$

and

$$\sum_{i=1}^m v(P_i) = \sum_{i=1}^m \delta \cdot \frac{\varepsilon}{2(b-a)} = \frac{\varepsilon}{2(b-a)} m\delta = \frac{\varepsilon}{2}$$

so the total volume of $\mathcal{P} \cup \bar{\mathcal{U}}$ is less than ε . Thus $C \cup F$ has measure zero, and C and F each do.

Thus Claim 2 is proved.

Now, by Claim 2, each of B, C, D, E, F has measure zero. So $B \cup C \cup D \cup E \cup F$ has measure zero, and by Claim 1 $\partial A_f \subseteq B \cup C \cup D \cup E \cup F$, so ∂A_f has measure zero. It is also bounded, so A_f is Jordan-measurable.

The last part of the proof is to show that $v(A_f) = \int_a^b f$. Since A_f is Jordan-measurable, χ_{A_f} is integrable on $[a, b] \times [0, M]$. So by Fubini's Theorem,

$$v(A_f) = \int_{[a,b] \times [0,M]} \chi_{A_f} = \int_a^b \left(\mathbf{L} \int_0^M \chi_{A_f}(x, y) dy \right) dx$$

For each fixed $x \in [a, b]$, $g_x = \chi_{A_f}(x, \cdot)$ is integrable as it is only discontinuous at $f(x)$. Thus

$$\mathbf{L} \int_0^M \chi_{A_f}(x, y) \, dy = \int_0^M \chi_{A_f}(x, y) \, dy$$

Moreover,

$$\int_0^M \chi_{A_f}(x, y) \, dy = \int_0^f(x) 1 \, dy = f(x)$$

So we have

$$v(A_f) = \int_a^b \left(\int_0^M \chi_{A_f}(x, y) \, dy \right) dx - \int_a^b f(x) \, dx = \int_a^b f \, dx \quad \square$$

Exercise 3-27 If $f : [a, b] \times [a, b] \rightarrow \mathbb{R}$ is continuous, show that

$$\int_a^b \int_a^y f(x, y) \, dx \, dy = \int_a^b \int_x^b f(x, y) \, dy \, dx$$

Proof. Define $C = \{(x, y) \in [a, b] \times [a, b] : y \geq x\}$. Then C has boundary $\partial C = (\{a\} \times [a, b]) \cup ([a, b] \times \{b\}) \cup \{(x, x) : x \in [a, b]\}$ which are all line segments, and thus have measure zero. So C is Jordan-measurable and $\chi_C f$ is integrable on $[a, b] \times [a, b]$. By Fubini's Theorem, since f is continuous,

$$\int_{[a, b] \times [a, b]} \chi_C f = \int_a^b \int_a^b \chi_C(x, y) f(x, y) \, dy \, dx = \int_a^b \int_x^b f(x, y) \, dy \, dx$$

But applying it in the opposite order,

$$\int_{[a, b] \times [a, b]} \chi_C f = \int_a^b \int_a^b \chi_C(x, y) f(x, y) \, dx \, dy = \int_a^b \int_a^y f(x, y) \, dx \, dy \quad \square$$

Exercise 3-28 Use Fubini's theorem to prove that $D_{1,2}f = D_{2,1}f$ if both are continuous.

Proof. Suppose that $D_{1,2}f$ and $D_{2,1}f$ both exist and are continuous. Then $D_{1,2}f - D_{2,1}f$ is continuous. Suppose there exists a such that $D_{1,2}f(a) - D_{2,1}f(a) > 0$ (for the case < 0 the proof is analogous). Then there exists a rectangle $A = [a, b] \times [c, d]$ containing a such that

$$D_{1,2}f(x) - D_{2,1}f(x) > \varepsilon$$

for any $x \in A$ and $\varepsilon > 0$ smaller than $D_{1,2}f(a) - D_{2,1}f(a)$. Since $D_{1,2}f - D_{2,1}f$ is continuous, it is integrable on A . So

$$\int_A D_{1,2}f - D_{2,1}f \geq \int_A \varepsilon = \varepsilon \int_A 1 = \varepsilon v(A) > 0$$

But by Fubini's Theorem,

$$\begin{aligned}
 \int_A D_{1,2}f &= \int_a^b \int_c^d D_{1,2}f(x, y) \, dy \, dx \\
 &= \int_a^b \left(\int_c^d \frac{d}{dy} D_1f(x, y) \, dy \right) dx \\
 &= \int_a^b D_1f(x, d) - D_1f(x, c) \, dx \\
 &= f(b, d) - f(b, c) - f(a, d) + f(a, c)
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 \int_A D_{2,1}f &= \int_c^d \int_a^b D_{2,1}f(x, y) \, dx \, dy \\
 &= \int_c^d D_2f(b, y) - D_2f(a, y) \, dy \\
 &= f(b, d) - f(a, d) - f(b, c) + f(a, c)
 \end{aligned}$$

So

$$\begin{aligned}
 \int_A D_{1,2}f - D_{2,1}f &= f(b, d) - f(b, c) - f(a, d) + f(a, c) - f(b, d) + f(a, d) + f(b, c) - f(a, c) \\
 &= 0
 \end{aligned}$$

contradiction. Thus $D_{1,2}f - D_{2,1}f = 0$ and $D_{1,2}f = D_{2,1}f$. □

Exercise 3-29 Use Fubini's theorem to derive an expression for the volume of a set of \mathbb{R}^3 obtained by revolving a Jordan-measurable set in the yz -plane about the z -axis.

Exercise 3-30 Let $C \subseteq [0, 1] \times [0, 1]$ contain at most one point on each horizontal and each vertical line, with $\partial C = [0, 1] \times [0, 1]$, as in Exercise 1-17. Show that

$$\int_{[0,1]} \left(\int_{[0,1]} \chi_C(x, y) \, dx \right) dy = \int_{[0,1]} \left(\int_{[0,1]} \chi_C(x, y) \, dy \right) dx$$

but

$$\int_{[0,1] \times [0,1]} \chi_C$$

does not exist.

Proof. Fix some $y \in [0, 1]$. Then A intersects $[0, 1] \times \{y\}$ at at most one point, so $h_y(x) = \chi_C(x, y)$ is zero everywhere except possibly one point. Thus it is nonzero at a finite number

of points, so

$$\int_{[0,1]} \chi_C(x, y) \, dx = 0$$

so

$$\int_{[0,1]} \left(\int_{[0,1]} \chi_C(x, y) \, dx \right) dy = 0$$

Similarly, for any $x \in [0, 1]$, A intersects $\{x\} \times [0, 1]$ at at most one point, so $g_x(y) = \chi_C(x, y)$ is nonzero at a finite number of points, so

$$\int_{[0,1]} \chi_C(x, y) \, dy = 0$$

and

$$\int_{[0,1]} \left(\int_{[0,1]} \chi_C(x, y) \, dy \right) dx = \int_{[0,1]} \left(\int_{[0,1]} \chi_C(x, y) \, dx \right) dy = 0$$

On the other hand, $\partial A = [0, 1] \times [0, 1]$ by assumption, which does not have measure zero and thus χ_C is not integrable on $[0, 1] \times [0, 1]$. \square

Exercise 3-31 If $A = [a_1, b_1] \times \dots \times [a_n, b_n]$ and $f : A \rightarrow \mathbb{R}$ is continuous, define $F : A \rightarrow \mathbb{R}$ by

$$F(x) = \int_{[a_1, x_1] \times \dots \times [a_n, x_n]} f$$

What is $D_i F(x)$ for $x \in \text{int } A$?

Define $G_1 : \mathbb{R} \rightarrow \mathbb{R}$ by

$$G_1(y) = F(y, x_2, \dots, x_n) = \int_{[a_1, y] \times \dots \times [a_n, x_n]} f$$

and $g_1 : \mathbb{R} \rightarrow \mathbb{R}$ by

$$g_1(y) = f(y, x_2, \dots, x_n)$$

Since f is continuous, we may apply Fubini's theorem to write

$$G_1(y) = \int_{a_1}^y \left(\int_{[a_2, x_2] \times \dots \times [a_n, x_n]} f(y, x^2, \dots, x^n) \, dx \right) dy$$

(where x^i represents a variable being integrated against, as opposed to x_i which is the i th component of x). So by the Fundamental Theorem of Calculus,

$$G_1'(y) = \left(\int_{[a_2, x_2] \times \dots \times [a_n, x_n]} f(y, x^2, \dots, x^n) \, dx \right) = \int_{a_2}^{x_2} \dots \int_{a_n}^{x_n} f(y, x^2, \dots, x^n) \, dx^n \dots dx^2$$

We can make a similar argument for g_i for any i , so that

$$D_i F(x) = g_i'(y) = \int_{a_1}^{x_1} \dots \int_{a_i}^{x_i} \dots \int_{a_n}^{x_n} f(x^1, \dots, x^{i-1}, x_i, x^{i+1}, \dots, x^n) \, dx^n \dots dx^1$$

where the strikethroughs indicate that the i th variables is not integrated against (that is, we integrate against all other variables but hold x_i constant).

Exercise 3-32 Let $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ be continuous and suppose D_2f is continuous. Define $F(y) = \int_a^b f(x, y) dx$. Prove **Leibnitz's rule**:

$$F'(y) = \int_a^b D_2f(x, y) dx$$

Proof. Define $g_x(y) : [c, d] \rightarrow \mathbb{R}$ by

$$g_x(y) = f(x, y)$$

Then by definition,

$$g'_x(y) = D_2f(x, y)$$

Since D_2f is continuous, by the Fundamental Theorem of Calculus,

$$f(x, y) = g_x(y) = g_x(c) + \int_c^y g'_x(t) dt = f(x, c) + \int_c^y D_2f(x, t) dt$$

So

$$F(y) = \int_a^b \left(f(x, c) + \int_c^y D_2f(x, t) dt \right) dx = \int_a^b f(x, c) dx + \int_a^b \int_c^y D_2f(x, t) dt dx$$

Now, by Fubini's Theorem we have

$$\int_a^b \int_c^y D_2f(x, t) dt dx = \int_c^y \int_a^b D_2f(x, t) dx dt$$

so

$$F'(y) = \frac{d}{dy} \int_a^b \int_c^y D_2f(x, t) dt dx = \frac{d}{dy} \int_c^y \int_a^b D_2f(x, t) dx dt = \int_a^b D_2f(x, y) dx \quad \square$$

Exercise 3-33 If $f : [a, b] \times [c, d] \rightarrow \mathbb{R}$ is continuous and D_2f is continuous, define

$$F(x, y) = \int_a^x f(t, y) dt$$

(a) Find D_1F and D_2F .

(b) If $G(x) = \int_a^{g(x)} f(t, x) dt$, find $G'(x)$.

(a) Define $h_y(x) = f(x, y)$. Let $F_y(x) = F(x, y)$, so that $D_1F(x, y) = F'_y(x)$. Then

$$F_y(x) = F(x, y) = \int_a^x f(t, y) dt = \int_a^x h_y(t) dt$$

so

$$D_1F(x, y) = F'_y(x) = \frac{d}{dx} \int_a^x h_y(t) dt = h_y(x) = f(x, y)$$

Now, define $H_x(y) = F(x, y) = \int_a^x f(t, y) dt$, so that $D_2F(x, y) = H'_x(y)$. By Leibnitz's rule from Exercise 3-32,

$$D_2F(x, y) = H'_x(y) = \int_a^x D_2f(t, y) dt$$

(b) Here we have $G(x) = F(g(x), x)$. By the Chain Rule,

$$G'(x) = D_1F(g(x), x)g'(x) + D_2F(g(x), x) = f(g(x), x)g'(x) + \int_a^x D_2f(t, x) dt$$

Exercise 3-34 Let $g_1, g_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$ be continuously differentiable and suppose $D_1g_2 = D_2g_1$. As in Exercise 2-21, let

$$f(x, y) = \int_0^x g_1(t, 0) dt + \int_0^y g_2(x, t) dt$$

Show that $D_1f(x, y) = g_1(x, y)$.

Proof. Differentiating term by term, the Fundamental Theorem of Calculus gives us

$$\frac{d}{dx} \int_0^x g_1(t, 0) dt = g_1(x, 0)$$

Now, since g_2 is continuously differentiable, it is also continuous, so by Leibnitz's Rule (considering D_1 rather than D_2),

$$\frac{d}{dx} \int_0^y g_2(x, t) dt = \int_0^y D_1g_2(x, t) dt$$

By assumption, $D_1g_2 = D_2g_1$, so

$$\int_0^y D_1g_2(x, t) dt = \int_0^y D_2g_1(x, t) dt$$

Then by the Fundamental Theorem of Calculus,

$$D_1f(x, y) = \frac{d}{dx} \int_0^x g_1(t, 0) dt + \frac{d}{dx} \int_0^y g_2(x, t) dt = g_1(x, 0) + \int_0^y D_2g_1(x, t) dt = g_1(x, y) \quad \square$$

Exercise 3-35

(a) Let $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation of one of the following types:

$$\begin{cases} g(e_i) = e_i, & i \neq j \\ g(e_j) = ae_j, \end{cases}$$

$$\begin{cases} g(e_i) = e_i, & i \neq j \\ g(e_j) = e_j + e_k, \end{cases}$$

$$\begin{cases} g(e_k) = e_k, & k \neq i, k \neq j \\ g(e_i) = e_j \\ g(e_j) = e_i \end{cases}$$

If U is a rectangle, show that $v(g(U)) = |\det g|v(U)$.

(b) Prove that $v(g(U)) = |\det g|v(U)$ for any linear transformation $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$.

(a) *Proof.* First note that the scaling factor of g is scale invariant, for any of the above cases. For instance, let $U = [a_1, b_1] \times \dots \times [a_n, b_n]$. Let $\mathbf{x} = (a_1, \dots, a_n)$. Then let $y \in U$. Since g is linear,

$$g(y) = g(y - \mathbf{x} + \mathbf{x}) = g(y - \mathbf{x}) + g(\mathbf{x})$$

So $g(U) = g(U - \mathbf{x}) + g(\mathbf{x})$, and thus $g(U)$ is a translated version of $g(U - \mathbf{x})$, which has the same volume. Thus we may assume that $U = [0, b_1] \times \dots \times [0, b_n]$.

Let $\vec{y}_i = b_i e_i$, so that $\vec{y}_1, \dots, \vec{y}_n$ are the edges of U . Then $g(U)$ is the rectangle with edges given by $g(\vec{y}_1), \dots, g(\vec{y}_n)$.

Case 1: We have

$$g(\vec{y}_i) = b_i g(e_i) = \begin{cases} b_i e_i, & i \neq j \\ ab_i e_i, & i = j \end{cases}$$

so $g(U) = [0, b_1] \times \dots \times [0, ab_j] \times \dots \times [0, b_n]$. Then

$$v(g(U)) = b_1 b_2 \dots ab_j \dots b_n = a(b_1 \dots b_n) = av(U)$$

Now, the matrix of g is

$$[g] = \begin{bmatrix} 1 & 0 & \dots & \dots & 0 \\ 0 & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & a & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 0 & 1 \end{bmatrix}$$

so

$$\det g = \det[g] = a$$

Case 2: Since g is linear, it is continuous. Assume without loss of generality that $j = 1$ and $k = 2$. Then $g(U) = V \times [0, b_3] \times \dots \times [0, b_n]$, where

$$V \subseteq \mathbb{R}^2 = \{(x, y) : 0 \leq x \leq b_1, x \leq y \leq x + b_2\}$$

is a rhombus. Then by Fubini's Theorem, (letting M be any rectangle bounding $g(U)$)

$$\begin{aligned} v(g(U)) &= \int_M \chi_{g(U)} \\ &= \int_0^{b_1} \int_x^{x+b_2} \left(\int_0^{b_3} \dots \int_0^{b_n} dx_n \dots dx_3 \right) dy dx \\ &= b_3 \dots b_n \int_0^{b_1} \int_x^{x+b_2} dy dx \\ &= b_3 \dots b_n \int_0^{b_1} b_2 \\ &= b_1 \dots b_n \\ &= v(U) \end{aligned}$$

The matrix of g is given by

$$[g] = \begin{bmatrix} 1 & 0 & \dots & \dots & 0 \\ 0 & \ddots & \ddots & 1 & \vdots \\ \vdots & \ddots & a & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 0 & 1 \end{bmatrix}$$

(where the off-diagonal 1 is an arbitrary off-diagonal location), which has determinant 1.

Case 3: We have

$$g(U) = [0, b_1] \times \dots \times \underbrace{[0, b_j]}_{i\text{th position}} \times \dots \times \underbrace{[0, b_i]}_{j\text{th position}} \times \dots \times [0, b_n]$$

which has $v(g(U)) = b_1 \dots b_n = v(U)$. The matrix of g is simply the identity matrix with two columns switched, so $\det g = -1$ and $|\det g| = 1$. \square

- (b) *Proof.* If $\det g = 0$, then $g(U)$ has volume zero for any U . If $\det g \neq 0$, then $\text{RREF}([g]) = I_n$. Moreover, note that the elementary row operations correspond

to the following matrices:

$$\left\{ \begin{array}{l} \left[\begin{array}{cccc} 1 & & & \\ & \ddots & & \\ & & a & \\ & & & \ddots \\ & & & & 1 \end{array} \right], & \text{scaling of a row} \\ \left[\begin{array}{cccc} 1 & & & \\ & \ddots & & \\ & & 0 & 1 \\ & & & \ddots \\ & & 1 & 0 \\ & & & & \ddots \\ & & & & & 1 \end{array} \right], & \text{row swap} \\ \left[\begin{array}{cccc} 1 & & & \\ & \ddots & & \\ & & 1 & a \\ & & & \ddots \\ & & & & \ddots \\ & & & & & 1 \end{array} \right], & \text{addition of rows} \end{array} \right.$$

The first two ERO matrices directly correspond to Cases 1 and 3, respectively.

For the third matrix, suppose the ERO in question sends R_i to $R_i + aR_j$. Then this ERO matrix may be written as $[g_1][g_2][g_3]$, where g_3 scales R_j by a (Case 1), g_2 is a Case 2 transformation which sends e_i to $e_i + e_j$, and g_1 scales R_j by $1/a$ (Case 1).

Thus any invertible transformation has a matrix which may be written as

$$[g] = [g_1] \dots [g_k] \text{RREF}([g]) = [g_1] \dots [g_k]$$

where each of the g_k is of one of the three types considered above. By the property of the determinant,

$$\det[g] = \det([g_1] \dots [g_k]) = \det([g_1]) \dots \det([g_k])$$

By applying part a), we have

$$\begin{aligned} v(g(U)) &= v(g_1(\dots(g_k(U)))) \\ &= |\det g_1| v(g_2(\dots(g_k(U)))) \\ &= |\det g_1| \dots |\det g_k| v(U) \\ &= |\det g_1 \dots \det g_k| v(U) \\ &= |\det g| v(U) \end{aligned}$$

□

Exercise 3-36 (Cavalieri's Principle) Let A and B be Jordan-measurable subsets of \mathbb{R}^3 . Let $A_c = \{(x, y) : (x, y, c) \in A\}$ and define B_c similarly. Suppose each A_c and B_c are Jordan-measurable and have the same area. Show that A and B have the same volume.

Proof. Let $M = [a_1, b_1] \times [a_2, b_2] \times [a_3, b_3]$ be a closed rectangle which bounds both A and B . Since A is Jordan-measurable, χ_A is integrable on M , and so is χ_B . By Fubini's Theorem,

$$\int_M \chi_A = \int_{a_3}^{b_3} \left(\int_{[a_1, b_1] \times [a_2, b_2]} \chi_A(x, y) \, dx \right) dy$$

where our use of the integral sign is justified since A_c is Jordan measurable. Then we may write

$$\int_{[a_1, b_1] \times [a_2, b_2]} \chi_A(x, y) \, dx = \int_{[a_1, b_1] \times [a_2, b_2]} \chi_{A_y}$$

This is precisely the area of A_y , which by assumption is the area of B_y . So

$$\begin{aligned} \int_M \chi_A &= \int_{a_3}^{b_3} \left(\int_{[a_1, b_1] \times [a_2, b_2]} \chi_A(x, y) \, dx \right) dy \\ &= \int_{a_3}^{b_3} \left(\int_{[a_1, b_1] \times [a_2, b_2]} \chi_{A_y} \right) dy \\ &= \int_{a_3}^{b_3} v(A_y) \\ &= \int_{a_3}^{b_3} v(B_y) \\ &= \int_{a_3}^{b_3} \left(\int_{[a_1, b_1] \times [a_2, b_2]} \chi_{B_y} \right) dy \\ &= \int_{a_3}^{b_3} \left(\int_{[a_1, b_1] \times [a_2, b_2]} \chi_B(x, y) \, dx \right) dy \\ &= \int_M \chi_B \end{aligned}$$

so $v(A) = v(B)$. □

Exercise 3-37

(a) Suppose that $f : (0, 1) \rightarrow \mathbb{R}$ is a nonnegative continuous function. Show that

$$\text{ext} \int_{(0,1)} f$$

exists if and only if

$$\lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon}^{1-\varepsilon} f$$

exists.

(b) Define

$$A_n := \left[1 - \frac{1}{2^n}, 1 - \frac{1}{2^{n+1}} \right]$$

Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies

$$\int_{A_n} f = \frac{(-1)^n}{n}$$

and $f = 0$ outside of $\bigcup_{i=1}^{\infty} A_i$. Suppose also that f does not change sign on the interiors of any of the A_n . Show that

$$\text{ext} \int_{(0,1)} f$$

does not exist, but

$$\lim_{\varepsilon \rightarrow 0^+} \text{ext} \int_{(\varepsilon, 1-\varepsilon)} f = -\ln 2$$

Note: The hypothesis that f does not change sign is not included in Spivak's original exercise. Spivak's exercise is incorrect as written, but this is not the only possible hypothesis to rectify the issue.

(a) *Proof.* (\implies) Suppose that

$$\text{ext} \int_{(0,1)} f$$

exists. Let Φ be some partition of unity subordinate to an admissible open cover \mathcal{O} of $(0, 1)$. Now, let $\varepsilon > 0$. Then let Φ_{ε} be the finite collection of $\varphi \in \Phi$ which are nonzero on $[\varepsilon, 1 - \varepsilon]$. Then we have

$$\int_{\varepsilon}^{1-\varepsilon} f = \int_{\varepsilon}^{1-\varepsilon} f \sum_{\varphi \in \Phi_{\varepsilon}} \varphi = \sum_{\varphi \in \Phi_{\varepsilon}} \int_{\varepsilon}^{1-\varepsilon} \varphi f$$

Now, since f is nonnegative, we have

$$\sum_{\varphi \in \Phi_\varepsilon} \int_\varepsilon^{1-\varepsilon} \varphi f \leq \sum_{\varphi \in \Phi_\varepsilon} \int_{C_\varphi} \varphi f \leq \sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f = \text{ext} \int_{(0,1)} f$$

So $\int_\varepsilon^{1-\varepsilon} f$ is bounded above. Moreover, let $\varepsilon' < \varepsilon$. Since f is nonnegative, we have

$$\int_\varepsilon^{1-\varepsilon} f \leq \int_{\varepsilon'}^{1-\varepsilon'} f$$

so

$$\lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^{1-\varepsilon} f$$

exists.

(\Leftarrow) Suppose that

$$\lim_{\varepsilon \rightarrow 0^+} \int_\varepsilon^{1-\varepsilon} f$$

exists. For any $n \in \mathbb{N}$, let

$$A_n := \left[\frac{1}{2^{n+1}}, \frac{1}{2^n} \right] \cup \left[1 - \frac{1}{2^n}, 1 - \frac{1}{2^{n+1}} \right]$$

By Exercise 2-26 there exists a C^∞ function φ_n such that $\varphi_n > 0$ on A_n but $\varphi_n = 0$ outside of some closed set contained in

$$\left(\frac{1}{2^{n+2}}, \frac{1}{2^{n-1}} \right) \cup \left(1 - \frac{1}{2^{n-1}}, 1 - \frac{1}{2^{n+2}} \right)$$

which can be smoothly extended to have domain $(-1, 2)$. Now, $(0, 1) = \bigcup_{i=1}^{\infty} A_i$, so for any $x \in (0, 1)$ at least one φ_n is nonzero at x . Moreover, it is clear that only finitely many are nonzero at x . So

$$\sum_{i=1}^{\infty} \varphi_i(x) > 0$$

and we may define the C^∞ function $\psi_n : (-1, 2) \rightarrow \mathbb{R}$ by

$$\psi_n(x) = \frac{\varphi_n(x)}{\sum_{i=1}^{\infty} \varphi_i(x)}$$

Then $\Psi = \{\psi_1, \psi_2, \dots\}$ is a partition of unity subordinate to the open cover

$$\mathcal{O} = \left\{ \left(\frac{1}{2^{n+2}}, \frac{1}{2^{n-1}} \right) \cup \left(1 - \frac{1}{2^{n-1}}, 1 - \frac{1}{2^{n+2}} \right) \right\}_{n=1}^{\infty}$$

Now, let S_k be the partial sum

$$S_k := \sum_{n=1}^k \int_{C_{\varphi_n}} \varphi_n |f| = \sum_{n=1}^k \int_{C_{\varphi_n}} \varphi_n |f|$$

For each φ_i we have

$$C_{\varphi_i} \subseteq \left(\frac{1}{2^{k+2}}, 1 - \frac{1}{2^{k+2}} \right)$$

so

$$S_k = \sum_{n=1}^k \int_{\frac{1}{2^{k+2}}}^{1 - \frac{1}{2^{k+2}}} \varphi_i f = \int_{\frac{1}{2^{k+2}}}^{1 - \frac{1}{2^{k+2}}} \sum_{i=1}^k \varphi_i f \leq \int_{\frac{1}{2^{k+2}}}^{1 - \frac{1}{2^{k+2}}} f \leq \lim_{\varepsilon \rightarrow 0^+} \int_{\varepsilon}^{1-\varepsilon} f$$

where the last inequality follows since f is nonnegative. Moreover, since f is nonnegative we have

$$\int_{C_{\varphi_i}} \varphi_i f \geq 0$$

so we have an increasing, bounded above series which thus converges. So f is extended integrable on $(0, 1)$. \square

(b) *Proof.* To show that

$$\text{ext} \int_{(0,1)} f$$

does not exist, we will exhibit a partition of unity Φ subordinate to an admissible open cover \mathcal{O} of $(0, 1)$ such that

$$\text{ext}_{\Phi} \int_{(0,1)} f = \sum_{\varphi \in \Phi} \int_{C_{\varphi}} \varphi |f|$$

does not converge. Define

$$O_n = \left(\frac{1}{2^{n+2}}, \frac{1}{2^{n-1}} \right) \cup \left(1 - \frac{1}{2^{n-1}}, 1 - \frac{1}{2^{n+2}} \right)$$

for each n , and let $\mathcal{O} = \{O_n\}_{n \in \mathbb{N}}$ be our open cover. By Exercise 2-26, pick ψ_n so that $\psi_n > 0$ on A_n but $\psi_n = 0$ outside of some closed set contained in O_n . Then only finitely many (but at least one) ψ_i are nonzero at any given point $x \in (0, 1)$, so write

$$\varphi_n(x) = \frac{\psi_n(x)}{\sum_{i=1}^{\infty} \psi_i(x)}$$

$\Phi = \{\varphi_1, \varphi_2, \dots\}$ is our desired partition of unity subordinate to \mathcal{O} .

Since $\bigcup_{i=1}^{\infty} A_i = [1/2, 1)$ and $f = 0$ outside of $\bigcup_{i=1}^{\infty} A_i$, we have

$$\text{supp}(\varphi_n |f|) \subseteq \left(1 - \frac{1}{2^{n-1}}, 1 - \frac{1}{2^{n+2}} \right)$$

so that

$$\int_{C_{\varphi_n}} \varphi_n |f| = \int_{\text{supp} \varphi_n |f|} \varphi_n |f| = \int_{A_{n-1}} \varphi_n |f| + \int_{A_n} \varphi_n |f| + \int_{A_{n+1}} \varphi_n |f|$$

(for $n = 1$ the first term is omitted). Letting

$$S_k = \sum_{i=1}^k \int_{C_{\varphi_i}} \varphi_i |f|$$

we have

$$\begin{aligned} & \sum_{i=1}^{\infty} \int_{C_{\varphi_i}} \varphi_i |f| \geq S_k \\ &= \sum_{i=1}^k \left(\int_{A_{i-1}} \varphi_i |f| + \int_{A_i} \varphi_i |f| + \int_{A_{i+1}} \varphi_i |f| \right) \\ &= \sum_{i=1}^{k-1} \int_{A_i} \varphi_{i+1} |f| + \sum_{i=1}^k \int_{A_i} \varphi_i |f| + \sum_{i=2}^{k+1} \int_{A_i} \varphi_{i-1} |f| \\ &\geq \sum_{i=1}^{k-1} \int_{A_i} \varphi_{i+1} |f| + \sum_{i=1}^k \int_{A_i} \varphi_i |f| + \sum_{i=2}^k \int_{A_i} \varphi_{i-1} |f| \\ &= \int_{A_1} |f|(\varphi_2 + \varphi_1) + \sum_{i=2}^{k-1} \left(\int_{A_i} |f|(\varphi_{i+1} + \varphi_i + \varphi_{i-1}) \right) + \int_{A_k} |f|(\varphi_k + \varphi_{k-1}) \\ &\geq \int_{A_1} |f|(\varphi_2 + \varphi_1) + \sum_{i=2}^{k-1} \left(\int_{A_i} |f|(\varphi_{i+1} + \varphi_i + \varphi_{i-1}) \right) \end{aligned}$$

Note that by construction, φ_1 and φ_2 are the only nonzero φ on A_1 , and $\varphi_{i-1}, \varphi_i, \varphi_{i+1}$ are the only nonzero φ on A_i for $i \geq 2$. Thus this simplifies to

$$\int_{A_1} |f| + \sum_{i=2}^{k-1} \int_{A_i} |f| \geq \sum_{i=1}^{k-1} \left| \int_{A_i} f \right| = \sum_{i=1}^{k-1} \frac{1}{n}$$

so (S_k) is the sequence of partial sums of the harmonic series, which diverges. Thus $\text{ext}_{\Phi} \int_{(0,1)} f$ does not exist.

But in contrast, we have

$$\text{ext} \int_{(\varepsilon, 1-\varepsilon)} f = \sum_{i=1}^{M-1} \int_{A_i} f + \int_{(1-1/2^M, 1-\varepsilon)} f$$

where M is the largest integer such that $1 - 1/2^M \leq 1 - \varepsilon$. If M is even then we have

$$\sum_{i=1}^{M-1} \int_{A_i} f \leq \text{ext} \int_{(\varepsilon, 1-\varepsilon)} f \leq \sum_{i=1}^M \int_{A_i} f$$

and if M is odd then

$$\sum_{i=1}^{M-1} \int_{A_i} f \geq \text{ext} \int_{(\varepsilon, 1-\varepsilon)} f \geq \sum_{i=1}^M \int_{A_i} f$$

so

$$\lim_{\varepsilon \rightarrow 0} \text{ext} \int_{(\varepsilon, 1-\varepsilon)} f = \sum_{i=1}^{\infty} \int_{A_i} f = \sum_{i=1}^M \frac{(-1)^i}{i} = -\ln 2 \quad \square$$

Exercise 3-38 Let A_n be a closed set contained in $(n, n+1)$. Suppose that $f : \mathbb{R} \rightarrow \mathbb{R}$ satisfies

$$\int_{A_i} f = \frac{(-1)^i}{i}$$

and $f = 0$ outside of $\bigcup_{i=1}^{\infty} A_i$. Find two partitions of unity Φ, Ψ for \mathbb{R} such that

$$\sum_{\varphi \in \Phi} \int_{C_\varphi} \varphi f$$

and

$$\sum_{\psi \in \Psi} \int_{C_\psi} \psi f$$

both converge absolutely, but to different values.

Proof. First, pick C^∞ functions $g_1, g_2, \dots : \mathbb{R} \rightarrow [0, 1]$ such that $g_i = 1$ on A_i and $g_i = 0$ outside of a closed set contained in $(i, i+1)$. Now, let $\varphi_n = g_{2n-1} + g_{2n}$. Then the collection $\Phi = \{\varphi_1, \varphi_2, \dots\}$, together with appropriately chosen functions a partition of unity for \mathbb{R} . We have

$$\int_{C_{\varphi_n}} \varphi_n f = \int_{C_{g_{2n-1}}} f + \int_{C_{g_{2n}}} f = \int_{A_{2n-1}} f + \int_{A_{2n}} f = \frac{-1}{2n-1} + \frac{1}{2n} = -\frac{1}{4n^2 - 2n}$$

Thus

$$\text{ext} \int_{\mathbb{R}} f = \sum_{i=1}^{\infty} \int_{C_{\varphi_i}} \varphi_i f = \sum_{i=1}^{\infty} -\frac{1}{4n^2 - 2n} = -\ln 2$$

If we instead pick $\psi_1 = g_1$ and $\psi_n = g_{2n} + g_{2n+1}$, then $\Psi = \{\psi_1, \psi_2, \dots\}$ (with appropriately chosen functions) forms a partition of unity and we similarly have

$$\text{ext} \int_{\mathbb{R}} f = \int_{A_1} f + \sum_{i=2}^{\infty} \left(\int_{A_{2n}} f + \int_{A_{2n+1}} f \right) = -1 + \sum_{i=2}^{\infty} \frac{1}{4n^2 + 2n} = -\frac{1}{6} - \ln 2$$

Both of the series indicated converge absolutely since they converge, and do not change sign. \square

Exercise 3-39 Prove Theorem 3.19 without the assumption $\det u'(x) \neq 0$ using Sard's Theorem.

Proof. Suppose $u : A \rightarrow \mathbb{R}^n$ is injective and continuously differentiable, with A open. Let C be the set of points $x \in A$ such that $\det u'(x) = 0$. $\det u'(x)$ is composed of products

and sums of the partial derivatives, which are continuous, so $x \mapsto \det u'(x)$ is continuous. So C is a closed set in A , which means that $A \setminus C$ is open in A and thus in \mathbb{R}^n . Then the restriction of u to $A \setminus C$ is an injective, continuously differentiable function defined on an open set with $\det u'(x) \neq 0$ for $x \in A \setminus C$. By Theorem 3.19, we have

$$\text{ext} \int_{u(A \setminus C)} f = \text{ext} \int_{A \setminus C} (f \circ u) |\det u'|$$

Since u is injective, $u(A \setminus C) = u(A) \setminus u(C)$. By Sard's Theorem, $u(C)$ has measure zero so

$$\text{ext} \int_{u(A)} f = \text{ext} \int_{u(A) \setminus u(C)} f + \text{ext} \int_{u(C)} f = \text{ext} \int_{u(A) \setminus u(C)} f$$

Now, since $(f \circ u) |\det u'| = 0$ on C , and

$$\text{ext} \int_{A \setminus C} |\det u'| = \text{ext} \int_A |\det u'|$$

By Sard's Theorem, $u(C)$ has measure zero. So we have

$$\text{ext} \int_{u(A)} 1 = \text{ext} \int_{u(A) \setminus u(C)} 1 = \text{ext} \int_{u(A \setminus C)} 1 = \text{ext} \int_{A \setminus C} |\det u'| = \text{ext} \int_A |\det u'|$$

□

Exercise 3-40

- (a) If $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuously differentiable and $\det g'(a) \neq 0$, prove that in some open set containing a we can write $g = T \circ g_n \circ \dots \circ g_1$, where g_i is of the form

$$g_i(x) = (x_1, \dots, f_i(x), \dots, x_n)$$

for some $f_i : \mathbb{R}^n \rightarrow \mathbb{R}$, and where T is a linear transformation.

Note: Spivak failed to require that g be C^1 .

- (b) Show that if f_i does not depend on $x_j, i \neq j$, then we can take $T = I$ if and only if $g'(a)$ is diagonal.

Note: Spivak's original question does not include the stipulation that f_i does not depend on the other variables, but it is incorrect as stated.

- (a) *Proof.* First note that it suffices to prove the case $g'(a) = I$. In the general case, we would consider $(Dg(a))^{-1} \circ g$, and then g may be written as $Dg(a)$ composed with the representation produced in the identity case.

Recursively define the following:

$$g_1(x) = (g^1(x), x_2, \dots, x_n)$$

$$g_2(x) = (x_1, g^2(g_1^{-1}(x)), x_3, \dots, x_n)$$

⋮

$$g_n(x) = (x_1, \dots, x_{n-1}, g^n(g_1^{-1}(\dots(g_{n-1}^{-1}(x))))))$$

The fact that each g_i^{-1} exists is by the Inverse Function Theorem, since each has $g'_i(a) = I$ and thus there is an open set around a where all g_i are invertible. It follows that

$$g = g_n \circ \dots \circ g_1$$

□

(b) (\implies) Suppose $T = I$. Then if $j \neq i$, we have

$$\begin{aligned} D_j g_i(a) &= D_j(g^i \circ (g_1^{-1} \circ \dots \circ g_{i-1}^{-1})(a)) \\ &= \underbrace{D_j g^i(g_1^{-1} \circ \dots \circ g_{i-1}^{-1})(a)}_{=0} D_j(g_1 \circ \dots \circ g_{i-1})(a) \\ &= 0 \end{aligned}$$

so $g'(a)$ is diagonal.

(\impliedby) Suppose

$$g'(a) = \begin{bmatrix} a_1 & & \\ & \ddots & \\ & & a_n \end{bmatrix}$$

where each a_i is nonzero. Then $g \circ [Dg(a)]^{-1}$ satisfies

$$(g \circ [Dg(a)]^{-1})'(a) = g'(Dg(a)^{-1}(a))[g'([Dg(a)]^{-1}(a))]^{-1} = I$$

So we have $g = g_n \circ \dots \circ g_1 \circ Dg(a)$. Since $Dg(a)$ is of the form

$$Dg(a) = f_1 \circ \dots \circ f_n$$

we can write

$$g = g_n \circ \dots \circ g_1 \circ f_1 \circ \dots \circ f_n$$

Since f_i only depends on and changes the i th coordinate, and the same is true for g_i , we can freely interchange them so long as the relative order of g_i, f_i is preserved for each i . So this becomes

$$g = (g_n \circ f_n) \circ \dots \circ (g_1 \circ f_1)$$

Define $f : \{r : r > 0\} \times (0, 2\pi) \rightarrow \mathbb{R}^2$ by $f(r, \theta) = (r \cos \theta, r \sin \theta)$.

- (a) Show that f is injective, compute $f'(r, \theta)$, and show that $\det f'(r, \theta) \neq 0$ for all (r, θ) . Show that $f(\{r : r > 0\} \times (0, 2\pi))$ is the set $A = \{x < 0 \text{ or } x \geq 0, y \neq 0\}$, as in Exercise 2-23.
- (b) If $P = f^{-1}$, show that $P(x, y) = (r(x, y), \theta(x, y))$, where

$$r(x, y) = \sqrt{x^2 + y^2}$$

$$\theta(x, y) = \begin{cases} \arctan \frac{y}{x}, & x > 0, y > 0 \\ \pi + \arctan \frac{y}{x}, & x < 0 \\ 2\pi + \arctan \frac{y}{x}, & x > 0, y < 0 \\ \frac{\pi}{2}, & x = 0, y > 0 \\ \frac{3\pi}{2}, & x = 0, y < 0 \end{cases}$$

Find $P'(x, y)$. P is called the **polar coordinate system** on A .

- (c) Let $C \subseteq A$ be the region between the circles of radius r_1 and r_2 and the half-lines through 0 which make angles of θ_1 and θ_2 with the x -axis. If $h : C \rightarrow \mathbb{R}$ is integrable and $h(x, y) = g(r(x, y), \theta(x, y))$, show that

$$\int_C h = \int_{r_1}^{r_2} \int_{\theta_1}^{\theta_2} r g(r, \theta) \, d\theta \, dr$$

If $B_r = \{(x, y) : x^2 + y^2 \leq r^2\}$, show that

$$\int_{B_r} h = \int_0^r \int_0^{2\pi} r g(r, \theta) \, d\theta \, dr$$

- (c) If $C_r = [-r, r] \times [-r, r]$, show that

$$\int_{B_r} e^{-(x^2+y^2)} \, dx \, dy = \pi(1 - e^{-r^2})$$

and

$$\int_{C_r} e^{-(x^2+y^2)} \, dx \, dy = \left(\int_{-r}^r e^{-x^2} \, dx \right)^2$$

- (e) Prove that

$$\lim_{r \rightarrow \infty} \int_{B_r} e^{-(x^2+y^2)} \, dx \, dy = \lim_{r \rightarrow \infty} \int_{C_r} e^{-(x^2+y^2)} \, dx \, dy$$

and conclude that

$$\int_{-\infty}^{\infty} e^{-x^2} = \sqrt{\pi}$$

(a) *Proof.* Suppose $r_1 \cos \theta_1 = r_2 \cos \theta_2$ and $r_1 \sin \theta_1 = r_2 \sin \theta_2$. Then

$$r_1^2 = r_1^2(\cos^2 \theta_1 + \sin^2 \theta_1) = r_2^2(\cos^2 \theta_2 + \sin^2 \theta_2) = r_2^2$$

so $r_1 = r_2$. So $\sin \theta_1 = \sin \theta_2$ and $\cos \theta_1 = \cos \theta_2$, and we conclude that $\theta_1 = \theta_2$. We have

$$\det f'(r, \theta) = \det \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix} = r \cos^2 \theta + r \sin^2 \theta = r > 0$$

$f(r, \theta) \in \mathbb{R}^2 \setminus A$ only if $y = 0$ and $x \geq 0$, which implies $\sin \theta = 0$ and $\cos \theta > 0$ and thus $\theta = 0$, or $\sin \theta = \cos \theta = 0$ which is impossible. So $f(\{r : r > 0\} \times (0, 2\pi)) \subseteq A$. Let $A = (x, y)$. Then take

$$r = \sqrt{x^2 + y^2}$$

$$\theta = \begin{cases} \arctan \frac{y}{x}, & x > 0, y > 0 \\ \pi + \arctan \frac{y}{x}, & x < 0 \\ 2\pi + \arctan \frac{y}{x}, & x > 0, y < 0 \\ \frac{\pi}{2}, & x = 0, y > 0 \\ \frac{3\pi}{2}, & x = 0, y < 0 \end{cases}$$

So $A \subseteq f(\{r : r > 0\} \times (0, 2\pi))$ and we have equality. \square

(b) *Proof.* It suffices to check that $r(f(r, \theta)) = r$ and $\theta(f(r, \theta)) = \theta$. The first equality is easy:

$$r(f(r, \theta)) = \sqrt{r^2 \cos^2 \theta + r^2 \sin^2 \theta} = r$$

For the second:

$$\begin{cases} 0 < \theta < \frac{\pi}{2} \implies \cos \theta > 0, \sin \theta > 0 \\ \frac{\pi}{2} < \theta < \frac{3\pi}{2} \implies \cos \theta < 0 \\ \frac{3\pi}{2} < \theta < 2\pi \implies \cos \theta > 0, \sin \theta < 0 \\ \theta = \frac{\pi}{2} \implies \cos \theta = 0, \sin \theta > 0 \\ \theta = \frac{3\pi}{2} \implies \cos \theta = 0, \sin \theta < 0 \end{cases}$$

Since $r > 0$, all of the following remain true when $\cos \theta$ is replaced by f_1 and $\sin \theta$ by f_2 . So we have

$$\begin{cases} 0 < \theta < \frac{\pi}{2} \implies \theta(f(r, \theta)) = \arctan \tan \theta = \theta \\ \frac{\pi}{2} < \theta < \frac{3\pi}{2} \implies \theta(f(r, \theta)) = \pi + \arctan \tan \theta = \theta \\ \frac{3\pi}{2} < \theta < 2\pi \implies \theta(f(r, \theta)) = 2\pi + \arctan \tan \theta = \theta \\ \theta = \frac{\pi}{2} \implies \theta(f(r, \theta)) = \frac{\pi}{2} = \theta \\ \theta = \frac{3\pi}{2} \implies \theta(f(r, \theta)) = \frac{3\pi}{2} = \theta \end{cases}$$

We have

$$\begin{aligned} D_1 P^1(x, y) &= D_1 r(x, y) = \frac{x}{\sqrt{x^2 + y^2}} \\ D_2 P^1(x, y) &= D_2 r(x, y) = \frac{y}{\sqrt{x^2 + y^2}} \\ D_1 P^2(x, y) &= D_1 \theta(x, y) = \frac{1}{1 + \frac{y^2}{x^2}} \left(-\frac{y}{x^2} \right) = -\frac{y}{x^2 + y^2} \\ D_2 P^2(x, y) &= D_2 \theta(x, y) = \begin{cases} \frac{1}{1 + \frac{y^2}{x^2}} \frac{1}{x} = \frac{x}{x^2 + y^2}, & x \neq 0 \\ 0, & x = 0 \end{cases} \end{aligned}$$

so

$$P'(x, y) = \begin{bmatrix} \frac{x}{\sqrt{x^2 + y^2}} & \frac{y}{\sqrt{x^2 + y^2}} \\ -\frac{y}{x^2 + y^2} & \frac{x}{x^2 + y^2} \end{bmatrix}$$

□

(c) *Proof.* Let $C' = P(C) = (r_1, r_2) \times (0, 2\pi)$, so that $C = P^{-1}C'$. Note also that $h = g \circ P$. P^{-1} is continuously differentiable by the Inverse Function Theorem, so by the Change of Variables theorem,

$$\int_C h = \int_{C'} (h \circ P^{-1}) |\det(P^{-1})'| = \int_{C'} (h \circ P^{-1}) \frac{1}{|\det P'|} = \int_{C'} g \frac{1}{|\det P'|}$$

We can calculate,

$$\det P'(x, y) = \frac{x^2 + y^2}{\sqrt{x^2 + y^2}^3} = \frac{1}{\sqrt{x^2 + y^2}} = \frac{1}{r}$$

So

$$\int_{C'} rg = \int_{(r_1, r_2) \times (0, 2\pi)} rg$$

By Fubini's Theorem, this becomes

$$\int_C h = \int_{r_1}^{r_2} \int_0^{2\pi} rg(r, \theta) d\theta dr$$

Similarly, let $B'_r = P(B_r) = (0, r) \times (0, 2\pi)$. By similar logic,

$$\int_{B_r} h = \int_{B'_r} (h \circ P^{-1}) \frac{1}{|\det P'|} = \int_{B'_r} gr = \int_{(0, r) \times (0, 2\pi)} gr = \int_0^r \int_0^{2\pi} rg(r, \theta) d\theta dr \quad \square$$

(d) *Proof.* Using the result from part c),

$$\int_{B_r} e^{-(x^2 + y^2)} dx dy = \int_0^r \int_0^{2\pi} re^{-r^2} d\theta dr = \int_0^r 2\pi re^{-r^2} dr = -\pi e^{-r^2} \Big|_0^r = \pi(1 - e^{-r^2})$$

By Fubini's Theorem,

$$\begin{aligned}
 \int_{C_r} e^{-(x^2+y^2)} dx dy &= \int_{-r}^r \left(\int_{-r}^r e^{-x^2} e^{-y^2} dy \right) dx \\
 &= \int_{-r}^r e^{-x^2} \left(\int_{-r}^r e^{-y^2} dy \right) dx \\
 &= \left(\int_{-r}^r e^{-x^2} dx \right)^2 \quad \square
 \end{aligned}$$

(e) *Proof.* The quantity $e^{-(x^2+y^2)}$ is positive everywhere. So for any r , there exists $r' > r$ such that $C_r \subseteq B_{r'}$, giving

$$\int_{C_r} e^{-(x^2+y^2)} dx dy \leq \int_{B_{r'}} e^{-(x^2+y^2)} dx dy$$

But we can also pick r'' so that $B_r \subseteq C_{r''}$ so that the other direction is true. This shows that

$$\lim_{r \rightarrow \infty} \int_{B_r} e^{-(x^2+y^2)} dx dy = \lim_{r \rightarrow \infty} \int_{C_r} e^{-(x^2+y^2)} dx dy$$

Then we have

$$\begin{aligned}
 \int_{-\infty}^{\infty} e^{-x^2} dx &= \lim_{r \rightarrow \infty} \int_{-r}^r e^{-x^2} dx \\
 &= \lim_{r \rightarrow \infty} \sqrt{\int_{C_r} e^{-x^2+y^2} dx dy} \\
 &= \sqrt{\lim_{r \rightarrow \infty} \int_{C_r} e^{-x^2+y^2} dx dy} \\
 &= \sqrt{\lim_{r \rightarrow \infty} \int_{B_r} e^{-x^2+y^2} dx dy} \\
 &= \sqrt{\lim_{r \rightarrow \infty} \pi(1 - e^{-r^2})} \\
 &= \sqrt{\pi} \quad \square
 \end{aligned}$$

Definitions

- k -tensor, 60
- admissible, 49
- alternating, 65
- angle preserving, 76
- boundary, 6
- Cartesian product, 5
- Cavalieri's Principle, 157
- characteristic function, 40
- closed, 5
- closed k -cell, 5
- closed rectangle, 5
- common refinement, 32
- compact, 6
- component functions, 9
- composition, 8
- content, 41
- content zero, 35
- continuous, 9
- continuously differentiable, 21
- derivative, 12
- differentiable, 12
- directional derivative, 120
- Euclidean n -space, 3
- extended integrable, 50, 51
- exterior, 6
- functional limit, 9
- image, 8
- inner product, 4, 64
- inner product preserving, 75
- integrable, 33
- integral, 33
- interior, 6
- inverse, 8
- iterated integrals, 42
- Jacobian matrix, 13
- Jordan-measurable, 40
- Leibnitz's rule, 152
- level curve, 28
- lower integral, 42
- measure zero, 34
- multilinear, 60
- norm, 3
- norm preserving, 75
- Note:, 46
- open, 5
- open k -cell, 5
- open cover, 6
- open rectangle, 5
- orthogonal, 80
- orthonormal, 64
- oscillation, 10
- partial derivative, 19
- partition, 31
- partition of unity, 46
- polar coordinate system, 165
- preimage, 8
- projection function, 9
- pullback, 63
- Pythagorean Identity, 80
- refinement, 32

sign, 65

standard basis, 5

subcover, 6

subordinate, 46

subrectangles, 31

Taylor polynomial, 93

tensor product, 61

upper integral, 42

vector valued functions, 8

volume, 41

wedge product, 67